

Original Research Article

**Effect of Boron and Lime on Boron Transformation in a Fluvaquentic Humaquepts Soil
Concerning Uptake and Yield of Green Gram**

Abstract

Understanding of different fractions and availability of boron (B) is essential while studying the response of crops towards B. Fractionation provides information about the chemistry of B and quantifies its bioavailability. Therefore, a pot experiment was performed during 2019 pre-kharif season in acid soil to investigate the effect of different levels of boron and lime on boron fractions (viz., readily soluble, specifically adsorbed, oxide bound, organically bound, residual and total boron), uptake and yield of green gram (*Vigna radiata* L.) as a test crop. The plant growth parameters were recorded at a regular intervals of 15 days after sowing, and at harvest. All the experiments were conducted with three-re in triplicates following factorial randomized block design (FRBD). The result revealed that the changes in different fractions of boron, its plant uptake and yield of green gram are significantly affected by the application of boron and lime. -All the fractions contribute towards plant available boron form. Inter conversion between boron fractions was also observed. Application of boron and lime significantly influenced different boron pools, boron uptake and yield of green gram.

Methodology not complete!!! How the samples analysed for B?

How was the data analysed?

Was the same amount of lime added to all? What is the role of lime?

Keywords: lime, boron, boron fraction, uptake and yield.

Introduction

Micronutrients such as B play an important role in achieving the potential yield of green grams. Boron has a close relationship with calcium both in soil and in plant. Calcium increases the B requirement of plants due to similarity in function (Golakya and Patel, 1986). Boron exists in the soil in five fractions. They are easily soluble, with specific adsorbed, oxide bound, organically bound, and residual boron properties (Padbhushan and Kumar 2017). The proportions of boron in soil are as follows: residual boron > organically bound boron > specifically adsorbed boron > readily soluble boron > oxide bound boron. The relative proportion of boron in the fractions and the transformation among these fractions are affected by a variety of factors such as soil pH, clay mineral, Fe and Al

Formatted: Indent: First line: 0"

oxide, CaCO₃ content ~~among others, and so on~~ (Raina *et al.*, 2006; Anitha *et al.*, 2013; Patra *et al.*, 2018).

Among the micronutrients, the deficiency of boron (B) is most widespread in acid soils. As under acid soil conditions, boron is more water soluble and therefore leaches below the root-zones of plants by rainfall, coupled with adsorption by aluminium (Al) and iron (Fe) oxide minerals. The difference between deficiency and toxicity is very narrow and, therefore, boron requires careful fertility management. Depending on the crop, it has a relatively low quantitative need. Micronutrients like B also have the significant effect to achieve the potential yield of green gram. Thus, the application of B not only enhanced the boron concentration and dry matter yield of green gram but also improved the quality of green gram (Das *et al.*, 2020). The effect of liming on the availability of boron in soil is usually assessed by plant uptake data, which requires elaborate experimentation involving plants.

Liming, which reduces soil acidity, also reduces B availability due to adsorption on freshly precipitated Al and Fe hydroxides (Tsadilas and Kassioti, 2005). However, there is little information available on the effect of liming, particularly at lower levels, on nutrient availability in acid soils. Liming has been shown to have both negative and positive effects on nutrient availability in acid soils and subsequent crop uptake (Quaggio *et al.*, 2004). The transformation of added B in lime-amended acid soils is poorly understood. A better understanding of the distribution of B in various soil fractions and their relationships with plant response, particularly in lime-amended soils, would provide a foundation for assessing soil B availability to plants and developing appropriate management strategies to combat B deficiency in acid soils.

Depending on the crop, its quantitative requirement is very low. **Thus, the use of B not only increased yield but also improved green gram quality. This is conclusion!** The effect of liming on boron availability in soil is typically assessed using plant uptake data, which necessitates extensive plant experimentation. Proper micronutrient management in acid soils has been discovered to be extremely important in sustaining higher crop yields (Bhupalraj *et al.*, 2005). The present investigation was undertaken to study the effect of boron and lime on boron fractions, its uptake, and yield of green gram (variety DGG-4) in a Fluvaquentic Humaquepts soil. However, only a few **M** micronutrients forms are available to plants and their determination is important for estimation of its availability to plants. An attempt has been made in this study to estimate the application of boron and lime on different forms of boron. With this preview, the present investigation was programmed to quantify the different forms of B in soil.

What were the objectives of the study?

Formatted: Highlight

2. Materials and Methods

2.1. Sampling location and soil characteristic

A pot experiments was conducted during pre-Kharif 2019, at the Department of Soil Science and Agricultural Chemistry, College of Agriculture, Iroisemba, Central Agricultural University, Imphal. The experiment was carried out under factorial randomized block design (FRBD) with total ten treatments replicated three times. The general characteristics were presented in **Table 1**.

Table 1: General characteristics of the soil used in the experiment

Soil Characteristics	Results	Methods
Textural Class	Clayey soil	
Sand (%)	24.32	Bouyoucos, 1962
Silt (%)	21.01	
Clay (%)	54.67	
pH (1:2.5 Soil: water ratio)	5.2	Jackson, 1973
EC (1:2.5 Soil: water ratio, dSm ⁻¹)	0.15	Jackson, 1973
CEC [cmol(p ⁺) kg ⁻¹]	17.2	Borah et al., 1987
Lime requirement (ton/ha)	4.62	Shoemaker <i>et al.</i> , 1961
Organic carbon (%)	1.18	Walkley and Black, 1934
Available nitrogen (kg ha ⁻¹)	274.59	Subbiah and Asija, 1956
Available phosphorus (kg ha ⁻¹)	19.20	Bray and Kurtz, 1945
Available potassium (kg ha ⁻¹)	208.37	Jackson, 1973
Calcium (mg kg ⁻¹)	215.6	Jackson, 1973
Magnesium (mg kg ⁻¹)	34.10	Jackson, 1973
Sulfur (mg kg ⁻¹)	17.48	Jackson, 1973
Boron (mg kg ⁻¹)	0.18	John <i>et al.</i> , 1975
Readily soluble boron (mg kg ⁻¹)	0.62	Datta <i>et al.</i> , 2002
Specifically adsorbed boron (mg kg ⁻¹)	2.21	
Oxide-bound boron (mg kg ⁻¹)	6.32	
Organically bound boron (mg kg ⁻¹)	8.55	
Residual boron (mg kg ⁻¹)	110.04	
Total boron (mg kg ⁻¹)	127.71	

Results should go to chapter 4.

Since all analysis was done in triplicate, the results should be mean of the three and sd should also be given.

3.2. Incubation experiment

Five kg each of air-dried soil was taken in a series of pots. The required quantity of N @ 20 kg ha⁻¹, P₂O₅ @ 40 kg ha⁻¹, and K₂O @ 20 kg ha⁻¹ was applied to all the pots as basal dose through urea, SSP, and MOP and also mixed with the soil. Lime was added 7 days before sowing and boron with borax (B-11%) as source material for basal application. Lime requirement was determined by the method described by Shoemaker *et al.*, (1961). For the current study five boron levels; i.e. B₁=0

kg ha⁻¹, B₂=0.5 kg ha⁻¹, B₃=1.0 kg ha⁻¹, B₄=1.5 kg ha⁻¹, B₅=2.0 kg ha⁻¹ and two levels of lime L₁= ~~un~~
~~no~~ lime, L₂= lime. A stock solution of the known concentration was prepared using mineralized water.
From the stock solution, a series of working solutions were prepared and sprayed on the soil in
accordance with the different treatment levels and mixed thoroughly The green gram var. DGG-4
was sown ~~ing~~ in each pot. The pots were irrigated with water throughout the crop growth period while
maintaining 60 % water-holding capacity (WHC). The whole plants were collected on the 15th, 30th,
45th, and 60th days after sowing (DAS) and at harvest by destructive sampling.

Laboratory procedures

Different B fractions in soil samples were determined according to Datta *et al.* (2002).
Readily soluble B was extracted by 0.01 M CaCl₂, specifically adsorbed B with 0.05 M KH₂PO₄, oxide
bound B with 0.175 M ammonium oxalate (pH 3.25), organically bound B with 0.5 M NaOH, and
residual B with soil digestion in H₂SO₄, HF, and HClO₄. Boron in the extracts was determined
colorimetrically by azomethine-H method (John *et al.*, 1975) depending on the extraction medium. All
above fractions ~~i.e.~~, readily soluble B to residual B were summed- up to get the total B content of the
soil. Green gram yield data were collected as per the standard method.

The processed plant samples were subjected to dry ashing at 550 – 600 °C in a muffle
furnace for (time ~~indicate~~) followed by the addition of 0.1 N HCl, and the total B in the extracts was
determined colorimetrically by using azomethine-H method at 420 nm wavelengths.

Statistical analysis

ANOVA was performed on all the data of the investigation to test the statistical
significance of the effects of the treatment. The significance of various effects was tested at 5% level
of probability.

Results and Discussion

Readily soluble boron

Present the results before discussing them.

Among all the fractions, readily soluble boron (RS-B) is the first and most easily
accessible form for plant uptake, as it is in soil solution or is weakly adsorbed by soil particles (Jin *et al.*, 1988).
When lime was added to the soil, there was a significant decrease in readily soluble boron
content in soil. ~~H-~~however, there was no effect to boron content in unlimed soil (**Figure 1**). According
to Barman *et al.* (2017), readily soluble boron decreases with lime application due to increased B
sorption caused by elevated level of calcite in the soil. Application of lime raises soil pH; boron

fixation generally increases in soils (Goldberg 1997).- In comparison to the control, the B applied system had a higher RS-B in soil. When comparing different B applications, soil applied with B₅ (2 kg B ha⁻¹) had significantly higher Readily Soluble-B, which was comparable to B₄ (1.5 kg B ha⁻¹) and B₃ (1 kg B ha⁻¹) at different stages of crop growth. -The application of boron at 0, 0.5, 1, 1.5, and 2 kg B ha⁻¹ with or without lime had no significant effect on readily soluble boron accumulation in soil at different stages of crop growth, according to an interaction study.

Specifically adsorbed boron (SA-B)

These fractions of boron are referred as so due to its specificity to get adsorbed may be specifically adsorbed onto clay surfaces or associated with organic matter in soil (Jin *et al.*, 1987). It is primarily determined by the clay content of the soil. This fraction is in plant available form after RS-B. (Tsalidas *et al.*, 1994). Unlimed soil had significantly higher content of SA-B fraction than limed soil (**Figure 2**). Kanprath (1971) reported that limed soils with high oxide coatings increased B by clays and decreased B availability. As a result, liming reduces the availability of B in soils. SA-B content in soil treated with 2 kg B ha⁻¹ was found to be significantly higher compared to other treatment except soil treated with 1.5 kg B ha⁻¹ throughout the crop growth period. This is in corroboration with the findings of Karthikeyan and Shukla (2011); Sathya *et al.*, (2013); Padbhushan and Kumar (2015b). An interaction study revealed that liming and boron had no effect on this fraction at different stages of crop growth.

Oxide-bound boron

The oxide bound boron (Ox-B) fraction is associated with Fe and Al oxides and hydroxides. Manganese (Mn) is incorporated into the structure via isomorphous substitution (McLaren and Crawford 1973; Tessier *et al.*, 1979; Shuman 1985). It is a less labile form of B i.e. it is in inaccessible forms (Jin *et al.*, 1988). **Figure 3** shows that during the entire crop growth stages, soil treated with boron had significantly higher oxide bound boron than control. On the 45th and 60th DAS and at harvest, soil applied with B₅ (2 kg B ha⁻¹) had a statistically higher oxide bound boron content than soil applied with B₄ (1.5 kg B ha⁻¹) followed by B₃ (1 kg B ha⁻¹) and B₂ (0.5 kg B ha⁻¹). Similar reports on increased oxide bound boron with increasing levels of boron were presented by Karthikeyan and Shukla (2011), Barman *et al.*, (2017), and Sidhu and Kumar (2018). Regardless of liming, boron application had a positive effect on the oxide bound boron fraction in soil. Lime addition increased the oxide bound boron content in soil significantly more than no lime addition during the entire study period. The effect of liming with applied boron on the oxide bound boron fraction in soil was found to be significant throughout the entire growth stages, according to an interaction study. On the 60th DAS

and at harvest, limed soil applied with B₅ (2 kg B ha⁻¹) had statistically higher oxide bound boron than limed soil applied with B₄ (1.5 kg B ha⁻¹).

Organically bound boron

Organic matter adsorbs boron (Goldberg, 1997) thus reducing its availability for plant uptake, accounts for 2-8 % of total-B content in soil (Padbushan and Kumar, 2015a). The conversion of other boron fractions to this boron fraction is responsible for the increase in organically bound boron (Karthikeyan and Shukla, 2011) from **Figure 4**, it is evident that addition of lime to the soil significantly lowers the organically bound boron content in soil compared to a no lime applied system. Among the various boron applications, B₂ (0.5 kg B ha⁻¹) had statistically higher organically bound boron, compared to B₅ (2 kg B ha⁻¹) during the entire growth period. Hou *et al.*, (1994) and Barman *et al.*, (2017) made similar observations regarding the decline in organically bound boron with increasing boron levels. An interaction study revealed that boron and lime application had a significant effect on organically bound boron at different growth stages, with the exception of the 60th DAS, where no significant difference was observed.

Residual boron

This is the largest fraction among all the other fractions of B, which has nothing to do with plant available B (Jin *et al.*, 1987; Tsalidas *et al.*, 1994; Padbushan and Kumar 2015b). Residual boron gradually decreased until harvest. At different stages of green gram growth, the boron-treated system had significantly higher residual boron concentrations than the untreated control. It was observed that throughout the experimental period soil treated with lime had significantly elevated level of residual boron in soil compared to unlimed soil. During the entire growth stages of green gram, a significant interaction effect of boron and lime application on residual boron content was observed. Among the various treatment combinations, limed soil with 2 kg B ha⁻¹ (B₅L₂) had a statistically higher residual boron content than limed soil with 1.5 kg B ha⁻¹ (B₄L₂) and 1 kg B ha⁻¹ (B₃L₂) throughout the entire stages of green gram growth (**Figure 5**).

Total boron

The concentration of total boron (T-B) in soil varies depending on its parent material and degree of weathering (Barber, 1995). A decreasing pattern of total boron with crop growth was observed until crop harvest (Sathya *et al.*, 2013). As shown in **Figure 6**, The addition of lime resulted in a statistically higher amount of total boron in the soil. Again, a significant interaction effect of boron and lime addition levels on total boron concentration was observed. On the 60th DAS and at harvest,

interaction data revealed that total boron concentration was comparatively higher in limed soil applied with 2 kg B ha⁻¹ (B₅L₂), followed by the same limed soil applied with 1.5 kg B ha⁻¹ (B₄L₂).

Uptake

The results of B uptake are presented in Figure 7. From figure 7, Boron uptake in green gram increased until harvest. A critical examination of the results revealed that at various stages of crop growth, green gram grown in boron-applied systems absorbed significantly more boron than the control (Figure 7). Among the various B applications, soil applied with B₅ (2 kg B ha⁻¹) had the highest boron uptake, followed by B₄ (1.5 kg B ha⁻¹), B₃ (1kg B ha⁻¹) and B₂ (0.5 kg B ha⁻¹) at different sampling stages, with a significant difference. On the 15th and 45th DAS, the results show that lime application significantly reduced boron uptake by green gram compared to the no lime applied system. This is supported by the finding that liming acid soil reduced available soil boron and its uptake by green gram, resulting in B deficiency (Tsadilas and Kassioti 2005). On the 30th and 60th DAS, as well as at harvest, limed soil had statistically higher boron uptake than unlimed soil. The interaction effect revealed significant differences in plant boron uptake across all growth stages when boron and lime were combined. On the 15th, 30th, and 45th DAS, soil treated with B₅L₁ had significantly higher boron uptake.

Seed yield of Green gram

Green gram seed yield (seed weight, g plant⁻¹) was higher in the boron and lime-treated system than in the control. The seed yield of green gram increased significantly as the boron level increased up to 2 kg B ha⁻¹ (Figure 8). The yield in the boron-treated soil was significantly higher at 2 kg ha⁻¹, followed by 1.5 kg ha⁻¹, 1 kg ha⁻¹, and 0.5 kg ha⁻¹.

It was also discovered that, regardless of boron application, the incorporation of lime increased seed yield significantly than without lime. A detailed analysis of the interaction effect revealed that limed soil applied with 2 kg B ha⁻¹ (B₅L₂) produced significantly higher seed yield than 1.5 kg B ha⁻¹ + lime (B₄L₂), which was statistically equivalent to 2 kg B ha⁻¹ + unlime (B₅L₁). Boron-treated soil combined with lime produced higher seed yield. Hirpara *et al.*, (2017) and Kamboj and Malik (2018) reported similar findings.

Conclusion

Liming significantly reduced the fraction of readily soluble B, specifically adsorbed B, and organically bound B in soil. However, the use of B had a positive effect on various fractions. Soil residual and total boron content decreased gradually as growth stages progressed until harvest. In terms of plant boron uptake, the interaction effect revealed significant differences between boron and

lime combinations. Green gram (variety DGGS-4) seed yield was significantly higher in limed soil applied with 2 kg B ha⁻¹ than in acid soil. Boron's agronomic effectiveness increased when applied at a higher dose in combination with lime.


References

- Anitha, M. S., K. S., Kumar, K. M., Nair, C. R., Shivaprasad, L. G. K., Naidu, & D, Sarkar. 2013. Soil boron and its fractions in agro-climatic zones of Karnataka. *Clay Research*, 32(1):- 25-33.
- Barber, S. A. (1995). *Soil Nutrient Bioavailability: a Mechanistic Approach*, 2nd ed, New York, NY: John Wiley & Sons. [Pp?](#)
- Barman, P., A. Sen, A. Phonglosa, & K. Bhattacharyya. (2017). Depth wise distribution of boron in some soils of red and laterite zone of West Bengal, India. *International Journal of Current Microbiology and Applied Sciences*, 6(12): 4126-4137.
- Bhupalraj, G., M. C. Patnaik, & K. M. Khadke. (2005). Molybdenum status in soils of Andhra Pradesh. Progress report of All India coordinated research project on micro and secondary nutrients and pollutant elements in soils and plants of Andhra Pradesh. ANGRAU, Hyderabad, 25:1-123???
- Bouyoucos, G. J. (1962). Hydrometer method improved for making particle size analyses of soils 1. *Agronomy journal*, 54(5), 464-465.
- Bray, R. H., & Kurtz, L. T. (1945). Determination of total, organic, and available forms of phosphorus in soils. *Soil science*, 59(1), 39-46.
- Das, H., Devi, N. S., Singh, R. K., Singh, A. H., Singh, L. N., & Singh, N. G., (2020). Influence of boron and lime on boron concentration and dry matter yield of green gram (*Vigna radiata* L.). *International Journal of Chemical Studies*, 8(6): 631-634.
- Datta. S. P., Rattan, R. K., Suribabu, K., & Datta, S. C., (2002). Fractionation and colorimetric determination of boron in soils. *Journal of Plant Nutrition and Soil Science*, 165(2): 179-184.
- Goldberg, S. (1997). Reactions of boron with soils. *Plant and soil*, 193(1-2): 35-48.
- Hirpara, D.V., Sakarvadia, H.L., Savaliya, C.M., Ranpariya, V. S. & Modhavadiya, V. L. (2017). Effect of different levels of boron and molybdenum on growth and yield of summer groundnut (*Arachis hypogaea* L.) under medium black calcareous soils of south Saurashtra region of Gujarat. *International Journal of Chemical Studies*, 5(5): 1290-1293.
- Hou, J., Evans, L. J. & Spiers. G. A., (1994). Boron fractionation in soils. *Communications in Soil Science and Plant Analysis*, 25(9-10): 1841-1853.
- Hou, J., Evans, L. J. & Spiers. G. A., (1996). Chemical fractionation of soil boron: I. Method development. *Canadian Journal of Soil Science*, 76(4): 485-491.
- Jackson, M. L., (1973). *Soil Chemical Analysis*. Prentice Hall of India Pvt. Ltd., New Delhi.
- Jin. J. Y., Martens, D. C. & Zelazny, L. W., (1987). Distribution and plant availability of soil; boron fractions. *Soil Science Society of America Journal*, 51(5): 1228-1231.
- Jin. J. Y., Martens, D. C. & Zelazny, L. W. (1988). Plant availability of applied and native boron in soils with diverse properties. *Plant and Soil*, 105: 127-132.
- John. M. K., Chuah, H. H., & Neufeld, J. H., (1975). Application of improved azomethine-H method to the determination of boron in soils and plants. *Analytical letters*, 8(8): 559-568.

Formatted: Highlight

- Kamboj, N., & Malik, R. S. (2018). Influence of phosphorus and boron application on yield, quality, nutrient content and their uptake by Green Gram (*Vigna radiata* L.). *International Journal of Current Microbiology and Applied Sciences*, 7(3), 1451-1458.
- Kanprath, N., (1971). Effect of lime on nutrient content of soils, yield and nutrient content of potato and infestation by leafminers. Fifth Annual Meeting of Agricultural Scientists (p. 139).
- Karthikeyan, K., & Shukla, L. M., (2011). Different forms of boron and sulphur and their effect on B and S contents in mustard (*Brassica juncea* L.) and sunflower (*Helianthus annuus* L.). *Agropedology*, 21(1): 28-34.
- McLaren, R.G., & Crawford, D.V., (1973). Studies on soil copper I. The fractionation of copper in soils. *Journal of Soil Science*, 24(2): 172-181.
- Padbhushan, R., & Kumar, D., (2015a). Distribution of boron in different fractions in some alkaline calcareous soils. *Communications in Soil Science and Plant Analysis*, 46(8): 939-953.
- Padbhushan, R., & Kumar, D., (2015b). Soil boron fractions and response of green gram in calcareous soils. *Journal of Plant Nutrition*, 38(8): 1143-1157.
- Padbhushan, R., & Kumar, D., (2017). Fractions of soil boron: A review. *The Journal of Agricultural Science*, 155(7): 1023-1032.
- Patra, A., Sinha, A. K., Rakesh, S., Biswas, S., & Mukhopadhyay, P., (2018). Different fractions of boron in soils of Alfisol and Entisol of West Bengal. *Journal of Pharmacognosy and Phytochemistry*, 7(1): 510-513.
- Quaggio, J. A., Gallo, P. B., Gerroh, C. O. Abreu, M. F. & Cantarella, H. 2004. Peanut response to lime and molybdenum application in low pH soils. *Revista Brasileira de Ciencia do Solo* 28: 659-664.
- Raina, J. N., Singh, U., & Bhandari A. R., (2006). Status, forms and transformation of boron in apple orchard soils of Himachal Pradesh. *Journal of the Indian Society of Soil Science*, 54(1): 45-49.
- Sathya, S., Mahendran, P. P. & Arulmozhiselvan, K., (2013). Influence of soil and foliar application of borax on fractions of boron under tomato cultivation in boron deficient soil of typic Haplustalf. *African Journal of Agricultural Research*, 8(21): 2567-2571.
- Shoemaker, H.E., McLean, E.O. & Pratt, P. F., (1961). Buffer methods for determining lime requirement of soils with appreciable amounts of extractable aluminum. *Soil Science Society of America Journal*, 25(4): 274-277.
- Shuman, L. M. (1985). Fractionation method for soil microelements. *Soil Science*, 140(1): 11-22.
- Subbiah B. V., & Asija G. L., (1956). A rapid procedure for estimation of available N in soils. *Current Science* 25: 259-260.
- Tessier, A., Campbell, P. G., & Bisson, M., (1979). Sequential extraction procedure for the speciation of particulate traces metals. *Analytical Chemistry*, 51(7): 844-851.
- Tsadiras, C. D., & Kassioti, T. (2005). Influence of liming and nitrogen forms on boron uptake by tobacco. *Communications in Soil Science and Plant Analysis*, 36(4-6): 701-708.
- Tsadiras, C. D., Yassoglou, N., Kosmas, C. S., & Kallianou, C. H., (1994). The availability of soil boron fractions to olive trees and barley and their relationships to soil properties. *Plant and Soil*, 162: 211-217.
- Walkley, A. & Black, I. A. (1934). An examination of the Degtjareff method for determining soil organic matter and a proposed modification of the chromic acid titration method. *Soil Science*, 37, 29-38.

UNDER PEER REVIEW



FIGURES

Indicate figure numbers and captions!!!!





Figure 1. Effect of boron and lime on readily soluble boron (ppm) content in soil grown with green gram (Error bar showing Standard Error of Mean)

(A) ~~Un~~ No limed

(B) Limed





Figure 2. Effect of boron and lime on specifically adsorbed boron (ppm) content in soil grown with green gram (Error bar showing Standard Error of Mean)

(A) Unlimed

(B) Limed





Figure 3. Effect of boron and lime on oxide bound boron (ppm) content in soil grown with green gram (Error bar showing Standard Error of Mean)

(A) Unlimed

(B) Limed





Figure 4. Effect of boron and lime on organically bound boron (ppm) content in soil grown with green gram (Error bar showing Standard Error of Mean)

(A) Unlimed

(B) Limed





Figure 5. Effect of boron and lime on residual boron (ppm) content in soil grown with green gram (Error bar showing Standard Error of Mean)

(A) Unlimed

(B) Limed





Figure: 6. Effect of boron and lime on total boron (ppm) content in soil grown with green gram (Error bar showing Standard Error of Mean)
(A) Unlimed
(B) Limed





Figure: 7. Effect of boron and lime on boron uptake ($\mu\text{g pot}^{-1}$) by green gram (Error bar showing Standard Error of Mean)

(A) Unlimed

(B) Limed




Figure 8. Effect of boron and lime on seed yield of green gram (g pot^{-1}):

L1 - Unlimed

L2 - Limed

Use No lime instead of unlimed all through.

Use color graphs for proper follow up of results.

Have tables also to supplement the graphs.

Any photographs from the study?

Boron levels:

$$B_1 = B @ 0 \text{ Kg ha}^{-1}$$

$$B_2 = B @ 0.5 \text{ Kg ha}^{-1}$$

$$B_3 = B @ 1.0 \text{ Kg ha}^{-1}$$

$B_4 = B @ 1.5 \text{ Kg ha}^{-1}$

$B_5 = B @ 2.0 \text{ Kg ha}^{-1}$

Liming:

$L_1 = \text{Unlimed}$

$L_2 = \text{Limed}$

UNDER PEER REVIEW