

# Effect of Boron and Lime on Boron Transformation in a Fluvaqueptic Humaquepts Soil Concerning Uptake and Yield of Green Gram

## Abstract

Understanding different fractions and availability of boron (B) is essential while studying the response of crops towards B. Fractionation provides information about the chemistry of B and quantifies its bioavailability. Therefore, a pot experiment was performed during the 2019 pre-kharif season in acid soil to study soil B fractions and response of green gram (*Vigna radiata* L.) to B application. Lime is applied to such soils with the primary objective of increasing the productivity of crops by enhancing the availability of native and applied plant nutrients. On the contrary, availability of B in soil was increase due to liming. The treatments comprised of five levels of soil applied B (0, 0.5, 1.0, 1.5 and 2.0 mg B kg<sup>-1</sup>) and two levels of lime, lime and no lime was used. The experiment was laid out in factorial randomized block design (FRBD) with three replications. The soil and plant growth parameters were recorded at a regular interval of 15 days after sowing, and at harvest. All the result revealed that the changes in different fractions of B (viz., readily soluble, specifically adsorbed, oxide bound, organically bound, residual and total B), its plant uptake and yield of green gram are significantly affected by the application of B and lime. All the fractions contribute towards plant available boron form. Inter conversion between B fractions was also observed. The application of B and lime significantly influenced different B pools, boron uptake and yield of green gram.

**Keywords:** lime, boron, boron fraction, uptake and yield.

## Introduction

“Micronutrients such as B play an important role in achieving the potential yield of green grams. Boron exists in the soil in five fractions. They are soluble, with specific adsorbed, oxide bound, organically bound, and residual boron properties” (Padbhushan and Kumar 2017). “The proportions of boron in soil are as follows: residual boron > organically bound boron > specifically adsorbed boron > readily soluble boron > oxide bound boron. The relative proportion of boron in the fractions and the transformation among these fractions are affected by a variety of factors such as soil pH, clay mineral, Fe and Al oxide, CaCO<sub>3</sub> content, and so on” (Raina *et al.*, 2006; Anitha *et al.*, 2013; Patra *et al.*, 2018).

“Among the micronutrients, the B is most widespread in acid soils. Under acid soil conditions, boron is more water soluble therefore leaches below the root-zones of plants by rainfall, coupled with adsorption by aluminium (Al) and iron (Fe) oxide minerals. The difference between

deficiency and toxicity is very narrow and, therefore, boron requires careful fertility management. Depending on the crop, it has a relatively low quantitative need. Micronutrients like B also have a significant effect on achieving the potential yield of green gram. Thus, applying B not only enhanced the boron concentration and dry matter yield of green gram but also improved the quality of green gram” (Das *et al.*, 2020). The effect of liming on the availability of boron in soil is usually assessed by plant uptake data, which requires elaborate experimentation involving plants.

“Liming, reduces soil acidity, B availability due to adsorption on freshly precipitated Al and Fe hydroxides” (Tsadilas and Kassioti, 2005). “Boron has a close relationship with calcium both in soil and in plant. Ca increases the B requirement of plants due to similarity in function” (Golakya and Patel, 1986). “However, little information available on the effect of liming, particularly at lower levels, on nutrient availability in acid soils. Liming has been shown to have both negative and positive effects on nutrient availability in acid soils and subsequent crop uptake” (Quaggio *et al.*, 2004). The transformation of added B in lime-amended acid soils is poorly understood. A better understanding of the distribution of B in various soil fractions and their relationships with plant response, particularly in lime-amended soils, would provide a foundation for assessing soil B availability to plants and developing appropriate management strategies to combat B deficiency in acid soils.

“Depending on the crop, its quantitative requirement is very low. Thus, the use of B not only increased yield but also improved green gram quality. The effect of liming on boron availability in soil is typically assessed using plant uptake data, which necessitates extensive plant experimentation. Proper micronutrient management in acid soils has been discovered to be extremely important in sustaining higher crop yields” (Bhupalraj *et al.*, 2005). The present investigation was undertaken to study the effect of boron and lime on boron fractions, its uptake, and yield of green gram (variety DGG-4) in a Fluvaquentic Humaquepts soil. However, only a few Micronutrients forms are available to plants and their determination is important for estimation of its availability to plants. This study has attempted to estimate the application of boron and lime on different levels of boron. With this preview, the present investigation was programmed to quantify the different forms of B in soil.

## **2. Materials and Methods**

### ***2.1. Sampling location and soil characteristic***

A pot experiment was conducted during pre-*Kharif* 2019, at the Department of Soil Science and Agricultural Chemistry, College of Agriculture, Iroisemba, Central Agricultural University, Imphal. The experiment was carried out under factorial randomized block design (FRBD) with ten treatments replicated three times. The general characteristics were presented in **Table 1**.

**Table 1: General characteristics of the soil used in the experiment**

Soil Characteristics	Results	Methods
Textural Class	Clayey soil	Bouyoucos, 1962
Sand (%)	24.32	
Silt (%)	21.01	
Clay (%)	54.67	
pH (1:2.5 Soil: water ratio)	5.2	Jackson, 1973
EC (1:2.5 Soil: water ratio, dSm <sup>-1</sup> )	0.15	Jackson, 1973
CEC [cmol(p <sup>+</sup> ) kg <sup>-1</sup> ]	17.2	Borah et al., 1987
Lime requirement (ton/ha)	4.62	Shoemaker <i>et al.</i> , 1961
Organic carbon (%)	1.18	Walkley and Black, 1934
Available nitrogen (kg ha <sup>-1</sup> )	274.59	Subbiah and Asija, 1956
Available phosphorus (kg ha <sup>-1</sup> )	19.20	Bray and Kurtz, 1945
Available potassium (kg ha <sup>-1</sup> )	208.37	Jackson, 1973
Calcium (mg kg <sup>-1</sup> )	215.6	Jackson, 1973
Magnesium (mg kg <sup>-1</sup> )	34.10	Jackson, 1973
Sulfur (mg kg <sup>-1</sup> )	17.48	Jackson, 1973
Boron (mg kg <sup>-1</sup> )	0.18	John <i>et al.</i> , 1975
Readily soluble boron (mg kg <sup>-1</sup> )	0.62	Datta <i>et al.</i> , 2002
Specifically adsorbed boron (mg kg <sup>-1</sup> )	2.21	
Oxide-bound boron (mg kg <sup>-1</sup> )	6.32	
Organically bound boron (mg kg <sup>-1</sup> )	8.55	
Residual boron (mg kg <sup>-1</sup> )	110.04	
Total boron (mg kg <sup>-1</sup> )	127.71	

### 3.2. Incubation experiment

Five kg each of air-dried soil was taken in a series of pots. The required quantity of N @ 20 kg ha<sup>-1</sup>, P<sub>2</sub>O<sub>5</sub> @ 40 kg ha<sup>-1</sup>, and K<sub>2</sub>O @ 20 kg ha<sup>-1</sup> was applied to all the pots as basal dose through urea, SSP, and MOP and also mixed with the soil. Lime was added 7 days before sowing and boron with borax (B-11%) as source material for basal application. The lime requirement was determined by the method described by Shoemaker *et al.*, (1961). For the current study five boron levels i.e. B<sub>1</sub>=0 kg ha<sup>-1</sup>, B<sub>2</sub>=0.5 kg ha<sup>-1</sup>, B<sub>3</sub>=1.0 kg ha<sup>-1</sup>, B<sub>4</sub>=1.5 kg ha<sup>-1</sup>, B<sub>5</sub>=2.0 kg ha<sup>-1</sup> and two levels of lime L<sub>1</sub>= unlime, L<sub>2</sub>= lime. A stock solution of the known concentration was prepared using mineralized water. From the stock solution, a series of working solutions were prepared and sprayed on the soil by the different treatment levels and mixed thoroughly. The green gram *var.* DGGS-4 was sowing in each pot. The pots were irrigated with water throughout the crop growth period maintaining 60 % water-holding capacity (WHC). The whole plants were collected on the 15<sup>th</sup>, 30<sup>th</sup>, 45<sup>th</sup>, and 60<sup>th</sup> days after sowing (DAS) and at harvest by destructive sampling.

### Laboratory procedures

Different B fractions in soil samples were determined according to Datta *et al.*, (2002). Readily soluble B was extracted by 0.01 M CaCl<sub>2</sub>, specifically adsorbed B with 0.05 M KH<sub>2</sub>PO<sub>4</sub>, oxide bound B with 0.175 M ammonium oxalate (pH 3.25), organically bound B with 0.5 M NaOH, and residual B with soil digestion in H<sub>2</sub>SO<sub>4</sub>, HF, and HClO<sub>4</sub>. Boron in the extracts was determined colorimetrically by the azomethine-H method (John *et al.*, 1975) depending on the extraction medium. All above fractions i.e., readily soluble B to residual B were summed- up to get the total B content of the soil. Green gram yield data were collected as per the standard method.

The processed plant samples were subjected to dry ashing at 550 – 600°C in a muffle furnace for (time) followed by the addition of 0.1 N HCl, and the total B in the extracts was determined colorimetrically by using the azomethine-H method at 420 nm wavelengths.

### **Statistical analysis**

ANOVA was performed on all the data to test the statistical significance of the effects of the treatment. The significance of various effects was tested at 5% level of probability.

## **Results and Discussion**

### **Readily soluble boron**

Readily soluble B fraction in soil significantly decreased with increasing levels of applied lime (**Figure 1**). Application of lime raises soil pH; boron fixation generally increases in soils (Goldberg 1997). Compared to the control, the B applied system had a higher RS-B in soil. When comparing different B applications, soil applied with B<sub>5</sub> (2 kg B ha<sup>-1</sup>) had significantly higher Readily Soluble-B, which was comparable to B<sub>4</sub> (1.5 kg B ha<sup>-1</sup>) and B<sub>3</sub> (1 kg B ha<sup>-1</sup>) at different stages of crop growth. According to an interaction study, the application of boron at 0, 0.5, 1.0, 1.5, and 2.0 kg B ha<sup>-1</sup> with or without lime had no significant effect on readily soluble boron accumulation in the soil at different crop growth stages. Application of B had a positive impact on this fraction. “Application of lime generally increased the B fixation in soils because it raised the soil pH. In addition to its effect on soil pH, CaCO<sub>3</sub> also acts as an important B adsorbing surface in soil” (Goldberg and Forster, 1991). Reduction in readily soluble B fraction may be responsible for severe reduction in dry matter yield in lime treated pot, particularly where no B was applied. According to Barman *et al.*, (2017), readily soluble boron decreases with lime application due to increased B sorption caused by elevated levels of calcite in the soil.

### **Specifically adsorbed boron (SA-B)**

There was no significant relationship of specifically adsorbed B with lime application, while this fraction of B was positively and significantly affected by the application of B (**Figure 2**).

These fractions of boron are referred to as so due to their specificity to get adsorbed may be specifically adsorbed onto clay surfaces or associated with organic matter in the soil (Jin *et al.*, 1987). It is primarily determined by the clay content of the soil. This fraction is in plant available form after RS-B. (Tsalidas *et al.*, 1994). Unlimed soil had significantly higher content of SA-B fraction than limed soil (Figure 2). Kanprath (1971) reported that limed soils with high oxide coatings increased B by clays and decreased B availability. As a result, liming reduces the availability of B in soils. SA-B content in soil treated with 2 kg B ha<sup>-1</sup> was significantly higher than other treatment except soil treated with 1.5 kg B ha<sup>-1</sup> throughout the crop growth period. Datta *et al.* (2002) inferred that “specifically adsorbed B originates from the weakly binding sites of both organic and inorganic constituents of soil and none of these constituents contributes exclusively towards this B fraction. This may partially explain the non-significant effect of liming on specifically adsorbed B in soil”.

### **Oxide-bound boron**

Effect of liming on oxide bound fraction of B was irregular. Whereas, effect of B application had a positive effect on oxide bound B fraction in soil. It is a less labile form of B i.e. it is in inaccessible forms (Jin *et al.*, 1988). **Figure 3** showed, soil treated with boron had significantly higher oxide bound boron than the control. On the 45<sup>th</sup> and 60<sup>th</sup> DAS and at harvest, soil applied with B<sub>5</sub> (2 kg B ha<sup>-1</sup>) had a statistically higher oxide bound boron content than soil applied with B<sub>4</sub> (1.5 kg B ha<sup>-1</sup>) followed by B<sub>3</sub> (1 kg B ha<sup>-1</sup>) and B<sub>2</sub> (0.5 kg B ha<sup>-1</sup>). Similar reports on increased oxide bound boron with increasing levels of boron were presented by Karthikeyan and Shukla (2011), Barman *et al.*, (2017), and Sidhu and Kumar (2018). On the 60<sup>th</sup> DAS and at harvest, limed soil with B<sub>5</sub> (2 kg B ha<sup>-1</sup>) had statistically higher oxide bound boron than limed soil with B<sub>4</sub> (1.5 kg B ha<sup>-1</sup>). It appears that substantial amount of applied B goes to oxide bound fraction as a result of liming. Regardless of liming, boron application positively affect the oxide bound boron fraction in soil. Lime addition increased the oxide bound boron content in soil significantly more than no lime addition during the entire study period.

### **Organically bound boron**

There was no significant relation of organically bound B fraction in soil with both lime and B application (**Figure 4**). Among the various boron applications, B<sub>2</sub> (0.5 kg B ha<sup>-1</sup>) had statistically higher organically bound boron, compared to B<sub>5</sub> (2 kg B ha<sup>-1</sup>) during the entire growth period. Hou *et al.* (1994) and Barman *et al.* (2017) made similar observations regarding the decline in organically bound boron with increasing boron levels. An interaction study revealed that boron and lime application had a significant effect on organically bound boron at different growth stages, with the exception of the 60<sup>th</sup> DAS, where no significant difference was observed. Organic matter adsorbs

boron (Goldberg, 1997) thus reducing its availability for plant uptake, accounting for 2-8 % of total-B content in soil (Padbhushan and Kumar, 2015a). The conversion of other boron fractions to this boron fraction is responsible for the increase in organically bound boron (Karthikeyan and Shukla, 2011)

### **Residual boron**

Effect of lime and B application residual fraction of B in soil was not statistically significant. The interactive effect of lime on this fraction was also not significant. It was observed that throughout the experimental period soil treated with lime had significantly elevated levels of residual boron in soil compared to unlimed soil. During the growth stages of green gram, a significant interaction effect of boron and lime application on residual boron content was observed. Among the various treatment combinations, limed soil with 2 kg B ha<sup>-1</sup> (B<sub>5</sub>L<sub>2</sub>) had a statistically higher residual boron content than limed soil with 1.5 kg B ha<sup>-1</sup> (B<sub>4</sub>L<sub>2</sub>) and 1 kg B ha<sup>-1</sup> (B<sub>3</sub>L<sub>2</sub>) throughout the entire stages of green gram growth (**Figure 5**). This is the largest fraction among all the other fractions of B, which has nothing to do with plant available B (Jin *et al.*, 1987; Tsalidas *et al.*, 1994; Padbhushan and Kumar 2015b). Residual boron gradually decreased until harvest. At different stages of green gram growth, the boron-treated system had significantly higher residual boron concentrations than the untreated control.

### **Total boron**

A decreasing pattern of total boron with crop growth was observed until crop harvest (Sathya *et al.*, 2013). As shown in **Figure 6**. Adding lime resulted in a statistically higher total boron in the soil. Again, a significant interaction effect of boron and lime addition levels on total boron concentration was observed. On the 60<sup>th</sup> DAS and at harvest, interaction data revealed that total boron concentration was comparatively higher in limed soil applied with 2 kg B ha<sup>-1</sup> (B<sub>5</sub>L<sub>2</sub>), followed by the same limed soil applied with 1.5 kg B ha<sup>-1</sup> (B<sub>4</sub>L<sub>2</sub>). The total boron concentration (T-B) varies depending on its parent material and degree of weathering (Barber, 1995).

### **Uptake**

Boron uptake in green gram increased until harvest. A critical examination of the results revealed that green gram grown in boron-applied systems at various stages of crop growth absorbed significantly more boron than the control (**Figure 7**). Among the various B applications, soil applied with B<sub>5</sub> (2 kg B ha<sup>-1</sup>) had the highest boron uptake, followed by B<sub>4</sub> (1.5 kg B ha<sup>-1</sup>), B<sub>3</sub> (1kg B ha<sup>-1</sup>) and B<sub>2</sub> (0.5 kg B ha<sup>-1</sup>) at different sampling stages, with a significant difference. The results of the 15<sup>th</sup> and 45<sup>th</sup> DAS, the results show that lime application significantly reduced boron uptake by green gram compared to the no lime applied system. This is supported by the finding that liming acid soil reduced available soil boron and its uptake by green gram, resulting in B deficiency (Tsalidas and Kassioti

2005). On the 30<sup>th</sup> and 60<sup>th</sup> DAS, as well as at the harvest, limed soil had statistically higher boron uptake than unlimed soil. The interaction effect revealed significant differences in plant boron uptake across all growth stages when combining boron and lime. On the 15<sup>th</sup>, 30<sup>th</sup>, and 45<sup>th</sup> DAS, soil treated with B<sub>5</sub>L<sub>1</sub> had significantly higher boron uptake.

### Seed yield of Green gram

Green gram seed yield (seed weight, g plant<sup>-1</sup>) was higher in the boron and lime-treated system than in the control. The seed yield of green gram increased significantly as the boron level increased up to 2 kg B ha<sup>-1</sup> (**Figure 8**). The yield in the boron-treated soil was significantly higher at 2 kg ha<sup>-1</sup>, followed by 1.5 kg ha<sup>-1</sup>, 1 kg ha<sup>-1</sup>, and 0.5 kg ha<sup>-1</sup>. It was also discovered that, regardless of boron application, the incorporation of lime increased seed yield significantly more than without lime. A detailed analysis of the interaction effect revealed that limed soil applied with 2 kg B ha<sup>-1</sup> (B<sub>5</sub>L<sub>2</sub>) produced significantly higher seed yield than 1.5 kg B ha<sup>-1</sup> + lime (B<sub>4</sub>L<sub>2</sub>), which was statistically equivalent to 2 kg B ha<sup>-1</sup> + unlime (B<sub>5</sub>L<sub>1</sub>). Boron-treated soil combined with lime produced a higher seed yield. Hirpara *et al.*, (2017) and Kamboj and Malik (2018) reported similar findings.

### Conclusion

Liming significantly reduced the fraction of readily soluble B, specifically adsorbed B, and organically bound B in soil. However, the use of B had a positive effect on various fractions. Soil residual and total boron content decreased gradually as growth stages progressed until harvest. Regarding plant boron uptake, the interaction effect revealed significant differences between boron and lime combinations. Green gram (variety DGGS-4) seed yield was significantly higher in limed soil applied with 2 kg B ha<sup>-1</sup> than in acid soil. Boron's agronomic effectiveness increased when applied at a higher dose with lime.

### References

- Anitha, M. S., K. S, Kumar, K. M, Nair, C. R, Shivaprasad, L. G. K, Naidu, & D, Sarkar. 2013. Soil boron and its fractions in agro-climatic zones of Karnataka. *Clay Research*, 32(1): 25-33.
- Barber, S. A. (1995). *Soil Nutrient Bioavailability: a Mechanistic Approach*, 2<sup>nd</sup> ed, New York, NY: John Wiley & Sons.
- Barman, P., A. Sen, A. Phonglosa, & K. Bhattacharyya. (2017). Depth wise distribution of boron in some soils of red and laterite zone of West Bengal, India. *International Journal of Current Microbiology and Applied Sciences*, 6(12): 4126-4137.
- Bhupalraj, G., M. C. Patnaik, & K. M. Khadke. (2005). Molybdenum status in soils of Andhra Pradesh. Progress report of All India coordinated research project on micro and secondary nutrients and pollutant elements in soils and plants of Andhra Pradesh. ANGRAU, Hyderabad, 25:1-123.

- Bouyoucos, G. J. (1962). Hydrometer method improved for making particle size analyses of soils 1. *Agronomy journal*, 54(5), 464-465.
- Bray, R. H., & Kurtz, L. T. (1945). Determination of total, organic, and available forms of phosphorus in soils. *Soil science*, 59(1), 39-46.
- Das, H., Devi, N. S., Singh, R. K., Singh, A. H., Singh, L. N., & Singh, N. G., (2020). Influence of boron and lime on boron concentration and dry matter yield of green gram (*Vigna radiata* L.). *International Journal of Chemical Studies*, 8(6): 631-634.
- Datta. S. P., Rattan, R. K., Suribabu, K., & Datta, S. C., (2002). Fractionation and colorimetric determination of boron in soils. *Journal of Plant Nutrition and Soil Science*, 165(2): 179-184.
- Goldberg. S. (1997). Reactions of boron with soils. *Plant and soil*, 193(1-2): 35-48.
- Hirpara, D.V., Sakarvadia, H.L., Savaliya, C.M., Ranpariya, V. S. & Modhavadiya, V. L. (2017). Effect of different levels of boron and molybdenum on growth and yield of summer groundnut (*Arachis hypogaea* L.) under medium black calcareous soils of south Saurashtra region of Gujarat. *International Journal of Chemical Studies*, 5(5): 1290-1293.
- Hou, J., Evans, L. J. & Spiers. G. A., (1994). Boron fractionation in soils. *Communications in Soil Science and Plant Analysis*, 25(9-10): 1841-1853.
- Hou, J., Evans, L. J. & Spiers. G. A., (1996). Chemical fractionation of soil boron: I. Method development. *Canadian Journal of Soil Science*, 76(4): 485-491.
- Jackson, M. L., (1973). *Soil Chemical Analysis*. Prentice Hall of India Pvt. Ltd., New Delhi.
- Jin. J. Y., Martens, D. C. & Zelazny, L. W., (1987). Distribution and plant availability of soil; boron fractions. *Soil Science Society of America Journal*, 51(5): 1228-1231.
- Jin. J. Y., Martens, D. C. & Zelazny, L. W. (1988). Plant availability of applied and native boron in soils with diverse properties. *Plant and Soil*, 105: 127-132.
- John. M. K., Chuah, H. H., & Neufeld, J. H., (1975). Application of improved azomethine-H method to the determination of boron in soils and plants. *Analytical letters*, 8(8): 559-568.
- Kamboj, N., & Malik, R. S. (2018). Influence of phosphorus and boron application on yield, quality, nutrient content and their uptake by Green Gram (*Vigna radiata* L.). *International Journal of Current Microbiology and Applied Sciences*, 7(3), 1451-1458.
- Kanprath. N., (1971). Effect of lime on nutrient content of soils, yield and nutrient content of potato and infestation by leafminers. Fifth Annual Meeting of Agricultural Scientists (p. 139).
- Karthikeyan. K., & Shukla, L. M., (2011). Different forms of boron and sulphur and their effect on B and S contents in mustard (*Brassica juncea* L.) and sunflower (*Helianthus annuus* L.). *Agropedology*, 21(1): 28-34.
- McLaren. R.G., & Crawford, D.V., (1973). Studies on soil copper I. The fractionation of copper in soils. *Journal of Soil Science*, 24(2): 172-181.
- Padbhushan. R., & Kumar, D., (2015a). Distribution of boron in different fractions in some alkaline calcareous soils. *Communications in Soil Science and Plant Analysis*, 46(8): 939-953.
- Padbhushan. R., & Kumar, D., (2015b). Soil boron fractions and response of green gram in calcareous soils. *Journal of Plant Nutrition*, 38(8): 1143-1157.
- Padbhushan. R., & Kumar, D., (2017). Fractions of soil boron: A review. *The Journal of Agricultural Science*, 155(7): 1023-1032.

- Patra, A., Sinha, A. K., Rakesh, S., Biswas, S., & Mukhopadhyay, P., (2018). Different fractions of boron in soils of Alfisol and Entisol of West Bengal. *Journal of Pharmacognosy and Phytochemistry*, 7(1): 510-513.
- Quaggio, J. A., Gallo, P. B., Gerroh, C. O. Abreu, M. F. & Cantarella, H. 2004. Peanut response to lime and molybdenum application in low pH soils. *Revista Brasileira de Ciencia do Solo* 28: 659-664.
- Raina, J. N., Singh, U., & Bhandari A. R., (2006). Status, forms and transformation of boron in apple orchard soils of Himachal Pradesh. *Journal of the Indian Society of Soil Science*, 54(1): 45-49.
- Sathya, S., Mahendran, P. P. & Arulmozhiselvan, K., (2013). Influence of soil and foliar application of borax on fractions of boron under tomato cultivation in boron deficient soil of typic Haplustalf. *African Journal of Agricultural Research*, 8(21): 2567-2571.
- Shoemaker, H.E., McLean, E.O. & Pratt, P. F., (1961). Buffer methods for determining lime requirement of soils with appreciable amounts of extractable aluminum. *Soil Science Society of America Journal*, 25(4): 274-277.
- Shuman, L. M. (1985). Fractionation method for soil microelements. *Soil Science*, 140(1): 11-22.
- Subbiah B. V., & Asija G. L., (1956). A rapid procedure for estimation of available N in soils. *Current Science* 25: 259-260.
- Tessier, A., Campbell, P. G., & Bisson, M., (1979). Sequential extraction procedure for the speciation of particulate traces metals. *Analytical Chemistry*, 51(7): 844-851.
- Tsadilas, C. D., & Kassioti, T. (2005). Influence of liming and nitrogen forms on boron uptake by tobacco. *Communications in Soil Science and Plant Analysis*, 36(4-6): 701-708.
- Tsalidas, C. D., Yassoglou, N., Kosmas, C. S., & Kallianou, C. H., (1994). The availability of soil boron fractions to olive trees and barley and their relationships to soil properties. *Plant and Soil*, 162: 211-217.
- Walkley, A. & Black, I. A. (1934). An examination of the Degtjareff method for determining soil organic matter and a proposed modification of the chromic acid titration method. *Soil Science*, 37, 29-38.

## FIGURES






Figure 1. Effect of boron and lime on readily soluble boron (ppm) content in soil grown with green gram (Error bar showing Standard Error of Mean)

(A) Unlimed



(B) Limed



**Figure 2. Effect of boron and lime on specifically adsorbed boron (ppm) content in soil grown with green gram (Error bar showing Standard Error of Mean)**

**(A) Unlimed**



**(B) Limed**



**Figure 3. Effect of boron and lime on oxide bound boron (ppm) content in soil grown with green gram (Error bar showing Standard Error of Mean)**

**(A) Unlimed**



**(B) Limed**



**Figure 4. Effect of boron and lime on organically bound boron (ppm) content in soil grown with green gram (Error bar showing Standard Error of Mean)**

**(A) Unlimed**



**(B) Limed**



**Figure 5. Effect of boron and lime on residual boron (ppm) content in soil grown with green gram (Error bar showing Standard Error of Mean)**

**(A) Unlimed**



**(B) Limed**



**Figure: 6. Effect of boron and lime on total boron (ppm) content in soil grown with green gram (Error bar showing Standard Error of Mean)**

**(A) Unlimed**

**(B) Limed**



**Figure: 7. Effect of boron and lime on boron uptake ( $\mu\text{g pot}^{-1}$ ) by green gram (Error bar showing Standard Error of Mean)**

**(A) Unlimed**

**(B) Limed**




Figure 8. Effect of boron and lime on seed yield of green gram (g pot<sup>-1</sup>):

■ Unlimed

■ Limed

**Boron levels:**

B<sub>1</sub> = B @ 0 Kg ha<sup>-1</sup>

B<sub>2</sub> = B @ 0.5 Kg ha<sup>-1</sup>

B<sub>3</sub> = B @ 1.0 Kg ha<sup>-1</sup>

B<sub>4</sub> = B @ 1.5 Kg ha<sup>-1</sup>

B<sub>5</sub> = B @ 2.0 Kg ha<sup>-1</sup>

**Liming:**

L<sub>1</sub> = Unlimed

L<sub>2</sub> = Limed