

Correlation between physiochemical characteristics of soil and the morphological characteristics of *Grewia optiva* Drummond in North western Himalayan region.

Abstract:

Study on Physiochemical properties of soil is critical for the long-term maintenance of agricultural crops and field trees, as well as their economic development. The current study was carried out in the Department of Tree Improvement and Genetic Resources, COF, Nauri, Solan (H.P.) during the period 2020-2022 to quantify the impact of soil nutrient variation on the morphological characteristics of *Grewia optiva* Drummond in different districts with variable climate and heterogeneous soils. The impacts on growth parameters (tree height, crown spread, leaf traits) of selected populations of *Grewia optiva* Drummond statistically analyzed using Karl Pearson correlation coefficient. Two composite soil samples representative of the different population were drawn from the two depths *i.e.*, 0-15cm (Surface layer) and 15-30cm (subsurface layer). These samples were collected underneath the selected populations of *Grewia optiva* Drummond. There was highly positive correlation observed between leaf area and soil N (0.509), leaf area and SOC (0.407), leaf area and soil P (0.728) and leaf area and soil K (0.577). Leaf length showed a highly significant correlation with SOC (0.401), soil N (0.509), soil P (0.710), and soil K (0.592). The tree height (0.385), tree diameter (0.602), crown spread (N-S) (0.629), crown spread (E-W) (0.334), branch nodal length (0.436) and leaf width (0.470) showed a significant positive correlation with soil P. Soil K showed a significant positive correlation with tree height (0.774), tree diameter (0.645), crown spread (N-S) (0.576), crown spread (E-W) (0.314), branch nodal length (0.737) and leaf width (0.592). soil pH showed highly significant correlation with Leaf width (0.449) .

Comment [FKM1]: Describe how data was analysed and which software was used

Please give the recommendations and conclusions

1. Introduction:

Grewia optiva, often known as Biul/Bihul/Bhimal, is a Tiliaceae plant. This species is favoured by mountainous farmers in Uttarakhand, Himachal Pradesh, Nepal, and elsewhere for qualities like as palatability, rapid growth, ease of propagation, and fodder production (Mukherjee et al., 2018). It supplies fodder during the lean season when there is no alternative to green fodder. 2017; Katoch et al. It possesses more than 70 (%) potential DM digestibility and 56.7 per cent effective degradability, making it a great energy source for ruminants (Singh et al., 1989). *Grewia* is a genus with around 150 species worldwide, 42 of which are located on the Indian subcontinent (Bhagta et al., 2021).

Soil consists of definite chemical, physical, mineralogical and biological properties, which provides a medium for plant growth (Thakre et al., 2012). The knowledge of physiochemical properties viz; organic carbon, available Nitrogen (N), Phosphorus (P_2O_5), Potassium (K_2O), pH, electrical conductivity, soil texture and bulk density of soil is also important to determine the available nutrient status in soil and to develop specific fertilizer recommendations. (Sumithra et al., 2013). As pH is a good indicator of the balance of available nutrients in the soil whereas, Electrical Conductivity can almost be viewed as the quantity of available nutrients in the soil. (Smith and Doran, 1996) [use most recent reference 1996 is too old](#). The response of trees to increasing atmospheric CO_2 concentrations is often mediated by the availability of nutrients in the soil (Schleppi et al., 2019). Whether terrestrial ecosystems, forests, crop land trees are sources or sinks for CO_2 and their growth will ultimately depend on interactions of the C cycle with the cycles of nutrients, especially nitrogen (N) and phosphorus (P) (Ellsworth et al., 2017). An increased production of exoenzymes has been found in several studies with CO_2 enrichment, and this effect has depended on the availability of N in the soil (Drake et al., 2013, Meier et al., 2017, Ochoa-Hueso et al., 2017). Nitrogen (N) is one of the most important biological elements for plants, agricultural crops and forest trees, because it is a component of amino acids, proteins, genetic materials, pigments, and other key organic molecules (Chen et al., 2014, Ji et al., 2015, Liu et al., 2018). N has

an irreplaceable role in organ construction, material metabolism, fruit yield, and the quality formation of fruit trees (Bai et al., 2016). Soil nitrogen (N) deficiencies can affect the photosynthetic N-use efficiency (PNUE), mesophyll conductance (g_m), and leaf N allocation (Tang et al., 2019). Potassium (K) is used for flowering purpose, it is also required for building of protein, photosynthesis, fruit quality and reduction of diseases (Valente et al., 2012). Potassium is an activator of dozens of important enzymes, such as protein synthesis, sugar transport, N and C metabolism, and photosynthesis. It plays an important role in the formation of yield and quality improvement (Marschner, 2012, Oosterhuis et al., 2014). Potassium has strong mobility in plants and plays an important role in regulating cell osmotic pressure and balancing the cations and anions in the cytoplasm (Hu et al., 2016a). Phosphorus (P) is also an essential plant nutrient for various tree growth functions (Jonard et al. 2015). Plants take up Phosphorus in its inorganic form as phosphate (Becquer et al., 2014). Phosphorus limitation decreases the efficiency of plant respiration (Jiang et al., 2019) and night respiration may increase along the N/P ratio. Therefore, considering the importance of the physiochemical properties of the soil mentioned above, the present study was carried out to perform a correlation and nutritional analysis of the soil under the populations of *Grewia optiva* Drummond in Himachal Pradesh.

2. Materials and Methods.

Site Description

In this section describe where the research was done, the seasons include climatic conditions, type of soils

Two composite soil samples representative of the different population were drawn from the two depths *i.e.*, 0-15cm (Surface layer) and 15-30cm (subsurface layer). These samples were collected underneath the selected populations of *Grewia optiva* Drummond. Each sample was air dried, grounded with wooden pestle and mortar, sieved through 2 mm sieve and stored in plastic containers. The soil samples were then analysed for morphological (soil colour), physiochemical (bulk density, particle density, porosity, pH, EC), and available nutrient status using standard methods (Table 1).

Table 1 : Soil parameters under study with their methods of measurements

Sr.No.	Soil Parameters	Analysis Method Used
1	Soil colour	Munsell soil colour chart
2	pH	Digital pH meter (Jackson,1973).
3	Organic carbon (%)	Chromic acid titration method (Walkley and Black, 1934)
4	Available Nitrogen (kg ha ⁻¹)	Micro Kjeldhal Method (Subbaiah and Asija, 1956)
5	Available Phosphorus (kg ha ⁻¹)	0.5 M sodium bicarbonate (NaHCO ₃) at 8.5 pH (Olsen <i>et al.</i> , 1954)
6	Available K (kg ha ⁻¹)	Flame photometric method (1N NH ₄ OAC extractable) (Merwin and Peech, 1951)
7.	EC (dSm ⁻¹)	Digital EC meter (Jackson,1973)

3. 4 Experimental design

Describe the treatments , experimental design used, how the experiment was laid out, if possible just put the experiementa layout with treatments in this section

3.5 Data Correction

3.6 In this section include the data that was corrected nd procedures involved

3- 3.6 Statistical Analysis:

Karl pearson correlation coefficient ($p < 0.05$) used to find correlation between physiochemical properties of soil and morphological characteristics of selected population of *Grewia optiva* which had been recorded simultaneously.

Formatted: Normal, Indent: Left: 0.25", No bullets or numbering

Formatted: Font: (Default) Arial, 12 pt, Bold, Font color: Text 1

Formatted

Formatted: Font: (Default) Arial, 12 pt, Bold, Font color: Text 1

Formatted: Normal, Indent: Left: 0.25", No bullets or numbering

Formatted: Font: (Default) Arial, 12 pt, Bold, Font color: Text 1

4. Results and Discussion:

Karl Pearson's correlation coefficients (at 5% level of significance) for tree and leaf morphometric characteristics (of different populations of *G. optiva*) and physiochemical characteristics of soil were worked out. It was evident from Table 2 and Fig. 1 that leaf area showed a significant positive correlation with soil N (0.509), SOC (0.407), soil P (0.728), and soil K (0.577). Leaf length showed a highly significant correlation with SOC (0.401), soil N (0.509), soil P (0.710), and soil K (0.592). The tree height (0.385), tree diameter (0.602), crown spread (N-S) (0.629), crown spread (E-W) (0.334), branch nodal length (0.436) and leaf width (0.470) showed a significant positive correlation with soil P. Soil K showed a significant positive correlation with tree height (0.774), tree diameter (0.645), crown spread (N-S) (0.576), crown spread (E-W) (0.314), branch nodal length (0.737) and leaf width (0.592). Soil pH showed highly significant correlation with Leaf width (0.449). Nitrogen Similar results recorded in Douglas-fir by (Brix and Ebell, 1969), which revealed that fertilization with Nitrogen increased basal area increment, stem height, and branch length, leaf area, leaf length-width, the no. of leaves per shoot increased markedly. (Wang et al., 2012) reported increased specific leaf area (SLA) and leaf area index (LAI) with N fertilization. A study on *Fagus sylvatica* by (Meier and Leuschner, 2008) reported positive effects on leaf area and LAI in forests with increased in nitrogen availability. In a similar study by (Herbert and Fownes, 1995) in native *Metrosideros polymorpha* forest showed that increased available phosphorus promoted an increase in photosynthetic area which led to increased tree growth. A similar study in *Eucalyptus grandis* by (Battie-Laclau et al., 2013) reported that K and Na applications enhanced tree leaf area by increasing both leaf longevity and the mean area of individual leaves. The most important role of pH is the control of nutrients solubility in soil. Nutrient availability usually decreases with increasing pH (Kazem et al., 2012). EC values affects uptake of nutrients by the tree. Very high or low EC decreases plants leaf size, leaf water content, leaf net photosynthetic rate (P_n), stomatal conductance (G_s), transpiration rate (T_r), (Sonneveld and De Kreij, 1996). Soil organic carbon is a natural resource for the sustainable development of human society and a key foundation for sustainable forestry development (Pan et al., 2015). It plays an important role in the formation and conservation of soil structure, soil nutrient cycling and soil biodiversity.

		**																	
CS(N-S)	0.602	0.728	0.588 **	1															
NPB	0.362	0.445	0.396	0.512 **	1														
NSB	- 0.178	0.026	0.442	0.065	0.215	1													
BNL	0.851	0.816	0.460	0.628	0.452	- 0.01 6	1												
LL	0.512	0.623	0.298	0.605	0.522	0.23 0	0.572	1											
LW	0.383	0.531	0.090	0.294	0.517	0.23 6	0.513	0.562	1										
LA	0.581	0.732	0.358	0.610	0.541	0.33 2	0.655	0.944	0.665	1									
PL	0.362	0.456	0.281	0.495	0.534	0.40 0	0.503	0.867	0.534	0.852	1								
BD	0.361	0.273	0.348	0.257	0.079	- 0.15 5	0.291	0.136	0.232	0.138	0.084	1							
PD	0.508	0.328	0.119	0.123	0.023	- 0.35 7	0.476	0.237	0.024	0.246	0.221	0.66 9	1						
Porosity	-	-	-	-	0.080	-	-	-	0.307	-	0.013	-	-	1					

	0.179	0.165	0.380	0.268		0.003	0.103	0.043		0.038		0.903	0.286							
SoilpH	0.225	0.221	0.078	0.037	0.028	0.008	0.416	0.135	0.449	0.148	0.282	-0.067	0.229	0.214	1					
EC	0.038	-0.042	0.105	-0.009	0.294	-0.011	0.154	0.046	-0.069	0.003	0.359	0.201	0.229	-0.145	0.328	1				
OC	0.062	0.192	0.178	0.273	0.127	0.217	-0.017	0.401**	0.169	0.407**	0.260	-0.120	-0.310	-0.018	-0.312	-0.433	1			
Soil N	0.129	0.230	0.220	0.464**	0.226	0.291	0.147	0.525**	0.202	0.509**	0.434**	0.297	0.350**	0.182	0.018	0.244	0.770	1		
Soil P	0.385**	0.602**	0.334**	0.629**	0.429**	0.049	0.436**	0.710**	0.470**	0.728**	0.607**	-0.064	0.088	0.130	0.236	0.010	0.522	0.738	1	
Soil K	0.774**	0.645**	0.314**	0.576**	0.289	-0.043	0.737**	0.458**	0.592**	0.577**	0.287	0.277	0.356**	-0.149	0.284	-0.232	0.174	0.229	0.437	1

**5% level of Significance, where TH-tree height, TD-tree diameter, CS-crown spread, NPB-number of primary branches, NSB-number of secondary branches, BNL-branch nodal length, LL-leaf length, LW-leaf width, LA-leaf area, PL-petiole length, BD-bulk density, PD-particle density, EC-electrical conductivity, OC-organic carbon, N-nitrogen, P-phosphorus, K-potassium




Fig. 1: correlation between physiochemical characteristics of soil and tree and leaf morphometric characteristics of *Grewia optiva* Drummond.

here TH-tree height, TD-tree diameter, CS-crown spread, NPB-number of primary branches, NSB-number of secondary branches, BNL-branch nodal length, LL-leaf length, LW-leaf width, LA-leaf area, PL-petiole length, BD-bulk density, PD-particle density, EC-electrical conductivity, OC-organic carbon, N-nitrogen, P-phosphorus, K-potassium

Conclusion :

The correlation developed between tree morphological and soil characteristics will help in quantify the impact of different soil characteristics on tree and leaf morphometric characteristics and help in selection of superior populations, further improvement and fertilizers recommendation dose.

Authors' Contribution

In this section describe the role played by the authors involved

Competing interest

In this section the author should declare if there is any competing interest

References:

Comment [FKM2]: Include most recent references

Bai, L., Deng, H., Zhang, X., Yu, X., Li, Y., 2016. Gibberellin Is Involved in Inhibition of Cucumber Growth and Nitrogen Uptake at Suboptimal Root-Zone Temperatures. PLoS One 11(5), e0156188. <https://doi.org/10.1371/journal.pone.0156188>

Battie-Laclau, P., Laclau, J.P., Piccolo, M.C., Arenque Beri, C., Mietton, L., Muniz, M.R., Meille-Buckeridge, M.S., Nouvellon, Y., Ranger, J., Bouillet, J.P., 2013. Influence of Potassium and Sodium Nutrition on Leaf Area Components in *Eucalyptus grandis* Trees. Plant and Soil 371, 19–35.

Becquer, A., Trap, J., Irshad, U., Ali, M.A., Claude, P., 2014. From soil to plant, the journey of P through trophic relationships and ectomycorrhizal association. Frontiers in Plant Science 5, 548. doi: 10.3389/fpls.2014.00548

Bhagta, S., Thakur, P., Sharma D., 2021. Genetic divergence study in *Grewia optiva* through Quantitative and Molecular markers. International Journal of Economic Plants, 029-033. DOI: [HTTPS://DOI.ORG/10.23910/2/2020.0396](https://doi.org/10.23910/2/2020.0396)

Brix, H., Ebell, L.F., 1969. Effects of Nitrogen fertilization on Growth, leaf area, and photosynthesis rate in Douglas-Fir. Forest Sciences 15, 189-196.

Chen, L.H., Dong, T.F., Duan, B.L., 2014. Sex-specific carbon and nitrogen partitioning under N deposition in *Populus cathayana*. *Trees* 28, 793–806. <https://doi.org/10.1007/s00468-014-0992-3>.

Drake, J.E., Gallet-Budynek, A., Hofmockel, K.S., Bernhardt, E.S., Billings, S.A., Jackson, R.B., 2013. Increases in the flux of carbon belowground stimulate nitrogen uptake and sustain the long-term enhancement of forest productivity under elevated CO₂. *Ecology Letters* 14, 349–357. doi: 10.1111/j.1461-0248.2011.01593.x

Ellsworth, D.S., Anderson, I.C., Crous, K.Y., Cooke, J., Drake, J.E., Gherlenda, A.N., 2017. Elevated CO₂ does not increase Eucalypt forest productivity on a low-phosphorus soil. *Nature Climate Change* 7, 279–283. doi: 10.1038/nclimate3235

Herbert, D.A., Fownes, J.H., 1995. Phosphorus limitation of forest leaf area and net primary production on a highly weathered soil. *Biogeochemistry* 29, 223-235.

Hu, W., Zhao, W., Yang, J., Oosterhuis, D.M., Loka, D.A., Zhou, Z., 2016a. Relationship between potassium fertilization and nitrogen metabolism in the leaf subtending the cotton (*Gossypium hirsutum* L.) boll during the boll development stage. *Plant Physiology and Biochemistry* 101, 113–123. doi: 10.1016/j.plaphy.2016.01.019

Jackson, M.L., 1973. *Soil Chemical Analysis*. Prentice Hall of India (Pvt.) Ltd., New Delhi. 85, 251-252.

Ji, D.H., Mao, Q.Z., Watanabe, Y., Kitao, M., Kitaoka, S., 2015. Effect of nitrogen loading on the growth and photosynthetic responses of Japanese larch seedlings grown under different light regimes. *Journal of Agricultural Meteorology* 71, 232–238. <https://doi.org/10.2480/agrmet.D-14-00027>

Jiang, M., Caldararu, S., Zaehle, S., Ellsworth, D.S., Medlyn, B.E., 2019. Towards a more physiological representation of vegetation phosphorus processes in land surface models. *New Phytologist* 222, 1223–1229. doi: 10.1111/nph.15688

Jonard, M., Furst, A., Verstraeten, A., 2015. Tree mineral nutrition is deteriorating in Europe. *Global Change Biology* 21, 418-430.

Katoch, R., Singh, S.K., Tripath, A., Kumar, N., 2017. Effect of seasonal variation in biochemical composition of leaves of fodder trees prevalent in the mid-hill region of Himachal Pradesh. *Range Management and Agroforestry* 38, 234-40.

Kazem, H., Tayebbeh, M., Farani, S., Jamaati-e-Somarin, 2012. Effect of elemental sulphur and compost on pH, electrical conductivity and phosphorus availability of one clay soil. *African Journal of Biotechnology* 11, 1425-1432.

Liu, N., Wang, J., Guo, Q., Wu, S., Rao, X., Cai, X., Lin, Z., 2018. Alterations in leaf nitrogen metabolism indicated the structural changes of subtropical forest by canopy addition of nitrogen. *Ecotoxicology and Environmental Safety* 160, 134–143. <https://doi.org/10.1016/j.ecoenv.2018.05.037> (2018)

Marschner, H., 2012. *Marschner's Mineral Nutrition of Higher Plants*. Cambridge, MA: Academic press.

Meier, I.C., Finzi, A.C., Phillips, R.P., 2017. Root exudates increase N availability by stimulating microbial turnover of fast-cycling N pools. *Soil Biology and Biochemistry* 106, 119–128. doi: 10.1016/j.soilbio.2016.12.004

Meier, I.C., Leuschner, C., 2008. Leaf Size and Leaf Area Index in *Fagus sylvatica* Forests: Competing Effects of Precipitation, Temperature, and Nitrogen Availability. *Ecosystems* 11, 655–669

Merwin, H.D., Peech, M., 1951. Exchangeability of soil potassium in the sand, silt and clay fractions as influenced by the nature of the complementary exchangeable cation. *Soil Science Society of America Journal*. 15, 125-28.

Miller, C., Turk, L.M., 2002. *Fundamentals of soil science Biotech*. Books, 1123/74, Trinagar, Delhi, India pp. 157

Mukherjee, A., Modal, T., Bist, J.K., Pattanayak, A., 2018. Farmer's preference of fodder trees in mid hills of Uttarakhand: A comprehensive ranking using analytical hierarchy process. *Range Management and Agroforestry* 39, 115-20.

Ochoa-Hueso, R., Hughes, J., Delgado-Baquerizo, M., Drake, J.E., Tjoelker, M.G., Pineiro, J., 2017. Rhizosphere-driven increase in nitrogen and phosphorus availability under elevated atmospheric CO₂ in a mature *Eucalyptus* woodland. *Plant and Soil* 416, 283–295. doi: 10.1007/s11104-017-3212-2

Olsen, S.R., Cole, C.V., Watanable, F.S., Dean, L.A., 1954. Estimation of available phosphorus in soil by extraction with NaHCO_3 , USDA Circular, US Washington. 939, 19p.

Oosterhuis, D., Loka, D., Kawakami, E., Pettigrew, W., 2014. The physiology of potassium in crop production. *Advances in Agronomy* 126, 203–234. doi: 10.1016/B978-0-12-800132-5.00003-1

Pan, G., Lu, H., Li, L., Zhang, J., Zhang, X., 2015. Soil Carbon Sequestration with Bioactivity: A New Emerging Frontier for Sustainable Soil Management. *Advances in Earth Science* 30, 940–951.

Schleppi, P., Korner, C., Klein, T., 2019. Increased Nitrogen availability in the soil under mature *Picea abies* trees exposed to elevated CO_2 concentrations. *Frontiers in Forests and Global Change* 2, 59.

Singh, B., Makkar, P.S., Negi, S.S., 1989. Rate and extent of digestion and potentially digestible dry matter and cell wall of various tree leaves. *Journal of Dairy Science* 72, 3233-39.

Smith, J.L., Doran, J.W., 1996. Measurement and use of pH and electrical conductivity for soil quality analysis. *Methods for assessing soil quality*, 169-185.

Sonneveld, C., De Kreij, C., 1996. Response of cucumber (*Cucumis sativus* L.) to an unequal distribution of salts in the root environment. *Plant and Soil* 209, 45-56.

Subbaiah, B.V., Asija, G.L., 1956. A rapid method for the estimation of available nitrogen in soil. *Current Science* 25, 258-260.

Sumithra, S., Ankalaiah, C., Rao, D., Yamuna, R.T., 2013. A case study on physico-chemical characteristics of soil around industrial and agricultural area of yerraguntla, kadapa district, AP, India. *International Journal Geology Earth and Environmental Sciences* 3, 28-34.

Tang, J., Sun, B., Cheng, R., Shi, Z., Luo D., Liu, S., Centritto, 2019. Effects of soil nitrogen (N) deficiency on photosynthetic N-use efficiency in N-fixing and non-N-fixing tree seedlings in subtropical China. *Scientific Reports* 9, 4604 <https://doi.org/10.1038/s41598-019-41035-1>

Thakre, Y.G., Choudhary, M.D., Raut, R.D., 2012. Physicochemical Characterization of Red and Black Soils of Wardha Region. International Journal of Chemical and Physical Sciences 1, 60-66.

Valente, D.S.M., Queiroz, D.M., Pinto, F., Santos, N.T., Santos, F.L., 2012. Definition of Management Zones in Coffee Production Fields Based on Apparent Soil Electrical Conductivity. Scientia Agricola 69, 173-179.

Walkley, A.J., Black, I.A., 1934. Estimation of soil organic carbon by the chromic acid titration method. Soil Science 37, 29-38.

Wang, D., Mathew, W.M., Jindong, S., Xiaohui, F., Fernando, M., Dokyoung, L., Michael C., 2012. Impact of nitrogen allocation on growth and photosynthesis of *Miscanthus* (*Miscanthus X giganteus*) GCB Bioenergy 4, 688–697.

Yosuf, M., Li J., Lu J., Ren, T., Cong R., Fahad, S., Li, X., 2017. Effects of fertilization on crop production and nutrient-supplying capacity under rice-oilseed rape rotation system. Scientific reports 7, 1-9.

UNDER PEER REVIEW