

An analysis of the Correlation between physiochemical characteristics of soil and the morphological characteristics of *Grewia optiva* Drummond in The Northwestern Himalayan Rregion.

Abstract:

Continued research Study on the pPhysiochemical properties of soil is critical for the long-term maintenance of agricultural cropping systems, (including trees) and field trees, in order to harness their economic benefits, as well as their economic development. The current study was carried out in the Department of Tree Improvement and Genetic Resources, COF, Nauri, Solan (H.P.) during the period 2020-2022 to quantify the impact of soil nutrient variation on the morphological characteristics of *Grewia optiva* Drummond in different districts with variable climate and heterogeneous soils. The impacts on growth parameters (tree height, crown spread, leaf traits) of selected populations of *Grewia optiva* Drummond were statistically analyzed using Karl Pearson correlation coefficient. Two composite soil samples representative of the different population-populations were drawn from the two depths *i.e.*, 0-15cm (Surface layer) and 15-30cm (subsurface layer). These samples were collected underneath the selected populations of *Grewia optiva* Drummond. There was a highly positive correlation observed between leaf area and soil N (0.509), leaf area and SOC (0.407), leaf area and soil P (0.728), and leaf area and soil K (0.577). Leaf length showed a highly significant correlation with SOC (0.401), soil N (0.509), soil P (0.710), and soil K (0.592). The tree height (0.385), tree diameter (0.602), crown spread (N-S) (0.629), crown spread (E-W) (0.334), branch nodal length (0.436) and leaf width (0.470) showed a significant positive correlation with soil P. Soil K showed a significant positive correlation with tree height (0.774), tree diameter (0.645), crown spread (N-S) (0.576), crown spread (E-W) (0.314), branch nodal length (0.737) and leaf width (0.592). soil pH demonstrated showed a highly significant correlation with Leaf width (0.449).

1. Introduction:

Grewia optiva, often known as Biul/Bihul/Bhimal, is a Tiliaceae plant. This species is favoured by mountainous farmers in Uttarakhand, Himachal Pradesh, Nepal, and elsewhere for qualities like as palatability, rapid growth, ease of propagation, and fodder production (Mukherjee et al., 2018). It supplies fodder during the lean season when there is no alternative to green fodder. (2017; Katoch et al. It possesses more than 70 (%) potential DM digestibility and 56.7 ~~per-cent~~ effective degradability, making it a great energy source for ruminants (Singh et al., 1989). *Grewia* is a genus with around 150 species worldwide, 42 of which are located on the Indian subcontinent (Bhagta et al., 2021).

Soil consists of definite chemical, physical, mineralogical and biological properties, which ~~provides~~ provide a medium for plant growth (Thakre et al., 2012). The knowledge of physiochemical properties viz; organic carbon, available Nitrogen (N), Phosphorus (P_2O_5), Potassium (K_2O), pH, electrical conductivity, soil texture and bulk density of soil is also important to determine the available nutrient status in soil and to develop specific fertilizer recommendations. (Sumithra et al., 2013). As pH is a good indicator of the balance of available nutrients in the soil whereas, Electrical Conductivity can almost be viewed as the quantity of available nutrients in the soil. (Smith and Doran, 1996). The response of trees to increasing atmospheric CO_2 concentrations is often mediated by the availability of nutrients in the soil (Schleppi et al., 2019). Whether terrestrial ecosystems, forests, ~~crop-land~~ cropland trees are sources or sinks for CO_2 and their growth will ultimately depend on interactions of the C cycle with the cycles of nutrients, especially nitrogen (N) and phosphorus (P) (Ellsworth et al., 2017). An increased production of exoenzymes has been found in several studies with CO_2 enrichment, and this effect has depended on the availability of N in the soil (Drake et al., 2013, Meier et al., 2017, Ochoa-Hueso et al., 2017). Nitrogen (N) is one of the most important biological elements for plants, agricultural crops and forest trees, because it is a component of amino acids, proteins, genetic materials, pigments, and other key organic molecules (Chen et al., 2014, Ji et al., 2015, Liu et al., 2018). N has an

irreplaceable role in organ construction, material metabolism, fruit yield, and the quality formation of fruit trees (Bai et al., 2016). Soil nitrogen (N) deficiencies can affect the photosynthetic N-use efficiency (PNUE), mesophyll conductance (g_m), and leaf N allocation (Tang et al., 2019). Potassium (K) is used for flowering purpose, it is also required for building of protein, photosynthesis, fruit quality and reduction of diseases (Valente et al., 2012). Potassium is an activator of dozens of important enzymes, such as protein synthesis, sugar transport, N and C metabolism, and photosynthesis. It plays an important role in the formation of yield and quality improvement (Marschner, 2012, Oosterhuis et al., 2014). Potassium has strong mobility in plants and plays an important role in regulating cell osmotic pressure and balancing the cations and anions in the cytoplasm (Hu et al., 2016a). Phosphorus (P) is also an essential plant nutrient for various tree growth functions (Jonard et al. 2015). Plants take up Phosphorus in its inorganic form as phosphate (Becquer et al., 2014). Phosphorus limitation decreases the efficiency of plant respiration (Jiang et al., 2019) and night respiration may increase along the N/P ratio. Therefore, considering the importance of the physiochemical properties of the soil mentioned above, the present study was carried out to perform a correlation and nutritional analysis of the soil under the populations of *Grewia optiva* Drummond in Himachal Pradesh.

2. Materials and Methods.

Two composite soil samples representative of the different population were drawn from the two depths *i.e.*, 0-15cm (Surface layer) and 15-30cm (subsurface layer). These samples were collected underneath the selected populations of *Grewia optiva* Drummond. Each sample was air dried, grounded with wooden pestle and mortar, sieved through 2 mm sieve and stored in plastic containers. The soil samples were then analysed for morphological (soil colour), physiochemical (bulk density, particle density, porosity, pH, EC), and available nutrient status using standard methods (Table 1).

Table 1 :Soil parameters analyzed under study with their methods of measurements

Sr.No.	Soil Parameters	Analysis Method Used
1	Soil colour	Munsell soil colour chart
2	pH	Digital pH meter (Jackson,1973).
3	Organic carbon (%)	Chromic acid titration method (Walkley and Black, 1934)
4	Available Nitrogen (kg ha ⁻¹)	Micro Kjeldhal Method (Subbaiah and Asija, 1956)
5	Available Phosphorus (kg ha ⁻¹)	0.5 M sodium bicarbonate (NaHCO ₃) at 8.5 pH (Olsen <i>et al.</i> , 1954)
6	Available K (kg ha ⁻¹)	Flame photometric method (1N NH ₄ OAC extractable) (Merwin and Peech, 1951)
7.	EC (dSm ⁻¹)	Digital EC meter (Jackson,1973)

3. Statistical Analysis:

Karl Pearson correlation coefficient ($p < 0.05$) used to find correlation between physiochemical properties of soil and morphological characteristics of selected population of *Grewia optiva* which had been recorded simultaneously.

4. Results and Discussion:

Karl Pearson's correlation coefficients (at 5% level of significance) for tree and leaf morphometric characteristics (of different populations of *G. optiva*) and physiochemical characteristics of soil were worked out. It was evident from Table 2 and Fig.1 that leaf area showed a significant positive correlation with soil N (0.509), SOC (0.407), soil P (0.728), and soil K (0.577). Leaf length showed a highly significant correlation with SOC (0.401), soil N (0.509), soil P (0.710), and soil K (0.592). The tree height (0.385), tree diameter (0.602), crown spread (N-S) (0.629), crown spread (E-W) (0.334), branch nodal length (0.436) and leaf width (0.470) showed a significant positive correlation with soil P. Soil K showed a significant positive correlation with tree height (0.774), tree diameter (0.645), crown spread (N-S) (0.576), crown spread (E-W) (0.314), branch nodal length (0.737) and leaf width (0.592). soil

pH showed highly significant correlation with Leaf width (0.449). Nitrogen Similar results recorded in Douglas-fir by (Brix and Ebell, 1969), which revealed that fertilization with Nitrogen increased basal area increment, stem height, and branch length, leaf area, leaf length-width, the no. of leaves per shoot increased markedly. (Wang et al., 2012) reported increased specific leaf area (SLA) and leaf area index (LAI) with N fertilization. A study on *Fagus sylvatica* by (Meier and Leuschner, 2008) reported positive effects on leaf area and LAI in forests with increased in nitrogen availability. In a similar study by (Herbert and Fownes, 1995) in native *Metrosideros polymorpha* forest showed that increased available phosphorus promoted an increase in photosynthetic area which led to increased tree growth. A similar study in *Eucalyptus grandis* by (Battie-Laclauet et al., 2013) reported that K and Na applications enhanced tree leaf area by increasing both leaf longevity and the mean area of individual leaves. The most important role of pH is the control of nutrients solubility in soil. Nutrient availability usually decreases with increasing pH (Kazem et al., 2012). EC values affects uptake of nutrients by the tree. Very high or low EC decreases plants leaf size, leaf water content, leaf net photosynthetic rate (P_n), stomatal conductance (G_s), transpiration rate (T_r), (Sonneveld and De Kreij, 1996). Soil organic carbon is a natural resource for the sustainable development of human society and a key foundation for sustainable forestry development (Panet et al., 2015). It plays an important role in the formation and conservation of soil structure, soil nutrient cycling and soil biodiversity. Nitrogen is considered to be the most important nutrient, and plants absorb more nitrogen than any other element. Nitrogen is essential in the formation of protein, and protein makes up much of the tissues of most living things. Increase in available phosphorus affected positively leaf area, crown spread, no. of branches and fruit. Phosphorus, is linked to a plant's ability to use and store energy, including the process of photosynthesis. It's also needed to help plants grow and develop normally (Yosuf et al., 2017). Potassium is known to affect cell division, cell permeability formation of carbohydrates, translocation of sugars, various enzyme actions and resistance of some plants to certain diseases (Miller and Turk, 2002). It helps strengthen plants' abilities to resist disease and plays an important role in increasing crop yields and overall quality. Potassium also protects the plant when the weather is cold or dry, strengthening its root system and preventing wilt. Thus available soil

						7	0.017	**		**		0.120	0.310		0.312	0.433				
Soil N	0.129	0.230	0.220	0.464**	0.226	0.291	0.147	0.525**	0.202	0.509**	0.434**	0.297	0.350**	0.182	0.018	0.244	0.770	1		
Soil P	0.385**	0.602**	0.334**	0.629**	0.429**	0.049	0.436**	0.710**	0.470**	0.728**	0.607**	0.064	0.088	0.130	0.236	0.010	0.522	0.738	1	
Soil K	0.774**	0.645**	0.314**	0.576**	0.289	-0.043	0.737**	0.458**	0.592**	0.577**	0.287	0.277	0.356**	-0.149	0.284	-0.232	0.174	0.229	0.437	1

**5% level of Significance, where TH-tree height, TD-tree diameter, CS-crown spread, NPB-number of primary branches, NSB-number of secondary branches, BNL-branch nodal length, LL-leaf length, LW-leaf width, LA-leaf area, PL-petiole length, BD-bulk density, PD-particle density, EC-electrical conductivity, OC-organic carbon, N-nitrogen, P-phosphorus, K-potassium




Fig. 1: correlation between physiochemical characteristics of soil and tree and leaf morphometric characteristics of *Grewia optiva* Drummond.

here TH-tree height, TD-tree diameter, CS-crown spread, NPB-number of primary branches, NSB-number of secondary branches, BNL-branch nodal length, LL-leaf length, LW-leaf width, LA-leaf area, PL-petiole length, BD-bulk density, PD-particle density, EC-electrical conductivity, OC-organic carbon, N-nitrogen, P-phosphorus, K-potassium

Conclusion :~~The~~The correlation developed between tree morphological and soil characteristics will help in ~~quantify~~quantifying the impact of different soil characteristics on tree and leaf morphometric characteristics and help in the selection of superior populations, further improvement and fertilizers recommendation dose.

References:

Bai, L., Deng, H., Zhang, X., Yu, X., Li, Y., 2016. Gibberellin Is Involved in Inhibition of Cucumber Growth and Nitrogen Uptake at Suboptimal Root-Zone Temperatures. PLoS One 11(5), e0156188. <https://doi.org/10.1371/journal.pone.0156188>

Battie-Laclau, P., Laclau, J.P., Piccolo, M.C., Arenque Beri, C., Mietton, L., Muniz, M.R., Meille-Buckeridge, M.S., Nouvellon, Y., Ranger, J., Bouillet, J.P., 2013. Influence of Potassium and Sodium Nutrition on Leaf Area Components in *Eucalyptus grandis* Trees. Plant and Soil 371, 19–35.

Becquer, A., Trap, J., Irshad, U., Ali, M.A., Claude, P., 2014. From soil to plant, the journey of P through trophic relationships and ectomycorrhizal association. Frontiers in Plant Science 5, 548. doi: 10.3389/fpls.2014.00548

Bhagta, S., Thakur, P., Sharma D., 2021. Genetic divergence study in *Grewia optiva* through Quantitative and Molecular markers. International Journal of Economic Plants, 029-033. DOI: [HTTPS://DOI.ORG/10.23910/2/2020.0396](https://doi.org/10.23910/2/2020.0396)

Brix, H., Ebell, L.F., 1969. Effects of Nitrogen fertilization on Growth, leaf area, and photosynthesis rate in Douglas-Fir. Forest Sciences 15, 189-196.

Chen, L.H., Dong, T.F., Duan, B.L., 2014. Sex-specific carbon and nitrogen partitioning under N deposition in *Populus cathayana*. Trees 28, 793–806. <https://doi.org/10.1007/s00468-014-0992-3>.

Drake, J.E., Gallet-Budynek, A., Hofmockel, K.S., Bernhardt, E.S., Billings, S.A., Jackson, R.B., 2013. Increases in the flux of carbon belowground stimulate nitrogen

uptake and sustain the long-term enhancement of forest productivity under elevated CO₂. *Ecology Letters* 14, 349–357. doi: 10.1111/j.1461-0248.2011.01593.x

Ellsworth, D.S., Anderson, I.C., Crous, K.Y., Cooke, J., Drake, J.E., Gherlenda, A.N., 2017. Elevated CO₂ does not increase Eucalypt forest productivity on a low-phosphorus soil. *Nature Climate Change* 7, 279–283. doi: 10.1038/nclimate3235

Herbert, D.A., Fownes, J.H., 1995. Phosphorus limitation of forest leaf area and net primary production on a highly weathered soil. *Biogeochemistry* 29, 223-235.

Hu, W., Zhao, W., Yang, J., Oosterhuis, D.M., Loka, D.A., Zhou, Z., 2016a. Relationship between potassium fertilization and nitrogen metabolism in the leaf subtending the cotton (*Gossypium hirsutum* L.) boll during the boll development stage. *Plant Physiology and Biochemistry* 101, 113–123. doi: 10.1016/j.plaphy.2016.01.019

Jackson, M.L., 1973. *Soil Chemical Analysis*. Prentice Hall of India (Pvt.) Ltd., New Delhi. 85, 251-252.

Ji, D.H., Mao, Q.Z., Watanabe, Y., Kitao, M., Kitaoka, S., 2015. Effect of nitrogen loading on the growth and photosynthetic responses of Japanese larch seedlings grown under different light regimes. *Journal of Agricultural Meteorology* 71, 232–238. <https://doi.org/10.2480/agrmet.D-14-00027>

Jiang, M., Caldararu, S., Zaehle, S., Ellsworth, D.S., Medlyn, B.E., 2019. Towards a more physiological representation of vegetation phosphorus processes in land surface models. *New Phytologist* 222, 1223–1229. doi: 10.1111/nph.15688

Jonard, M., Furst, A., Verstraeten, A., 2015. Tree mineral nutrition is deteriorating in Europe. *Global Change Biology* 21, 418-430.

Katoch, R., Singh, S.K., Tripath, A., Kumar, N., 2017. Effect of seasonal variation in biochemical composition of leaves of fodder trees prevalent in the mid-hill region of Himachal Pradesh. *Range Management and Agroforestry* 38, 234-40.

Kazem, H., Tayebbeh, M., Farani, S., Jamaati-e-Somarin, 2012. Effect of elemental sulphur and compost on pH, electrical conductivity and phosphorus availability of one clay soil. *African Journal of Biotechnology* 11, 1425-1432.

Liu, N., Wang, J., Guo, Q., Wu, S., Rao, X., Cai, X., Lin, Z., 2018. Alterations in leaf nitrogen metabolism indicated the structural changes of subtropical forest by canopy addition of nitrogen. *Ecotoxicology and Environmental Safety* 160, 134–143. <https://doi.org/10.1016/j.ecoenv.2018.05.037> (2018)

Marschner, H., 2012. *Marschner's Mineral Nutrition of Higher Plants*. Cambridge, MA: Academic press.

Meier, I.C., Finzi, A.C., Phillips, R.P., 2017. Root exudates increase N availability by stimulating microbial turnover of fast-cycling N pools. *Soil Biology and Biochemistry* 106, 119–128. doi: 10.1016/j.soilbio.2016.12.004

Meier, I.C., Leuschner, C., 2008. Leaf Size and Leaf Area Index in *Fagus sylvatica* Forests: Competing Effects of Precipitation, Temperature, and Nitrogen Availability. *Ecosystems* 11, 655–669

Merwin, H.D., Peech, M., 1951. Exchangeability of soil potassium in the sand, silt and clay fractions as influenced by the nature of the complementary exchangeable cation. *Soil Science Society of America Journal*. 15, 125-28.

Miller, C., Turk, L.M., 2002. *Fundamentals of soil science* Biotech. Books, 1123/74, Trinagar, Delhi, India pp. 157

Mukherjee, A., Modal, T., Bist, J.K., Pattanayak, A., 2018. Farmer's preference of fodder trees in mid hills of Uttarakhand: A comprehensive ranking using analytical hierarchy process. *Range Management and Agroforestry* 39, 115-20.

Ochoa-Hueso, R., Hughes, J., Delgado-Baquerizo, M., Drake, J.E., Tjoelker, M.G., Pineiro, J., 2017. Rhizosphere-driven increase in nitrogen and phosphorus availability under elevated atmospheric CO₂ in a mature *Eucalyptus* woodland. *Plant and Soil* 416, 283–295. doi: 10.1007/s11104-017-3212-2

Olsen, S.R., Cole, C.V., Watanable, F.S., Dean, L.A., 1954. Estimation of available phosphorus in soil by extraction with NaHCO₃, USDA Circular, US Washington. 939, 19p.

Oosterhuis, D., Loka, D., Kawakami, E., Pettigrew, W., 2014. The physiology of potassium in crop production. *Advances in Agronomy* 126, 203–234. doi: 10.1016/B978-0-12-800132-5.00003-1

Pan, G., Lu, H., Li, L., Zhang, J., Zhang, X., 2015. Soil Carbon Sequestration with Bioactivity: A New Emerging Frontier for Sustainable Soil Management. *Advances in Earth Science* 30, 940–951.

Schleppi, P., Korner, C., Klein, T., 2019. Increased Nitrogen availability in the soil under mature *Picea abies* trees exposed to elevated CO_2 concentrations. *Frontiers in Forests and Global Change* 2, 59.

Singh, B., Makkar, P.S., Negi, S.S., 1989. Rate and extent of digestion and potentially digestible dry matter and cell wall of various tree leaves. *Journal of Dairy Science* 72, 3233-39.

Smith, J.L., Doran, J.W., 1996. Measurement and use of pH and electrical conductivity for soil quality analysis. *Methods for assessing soil quality*, 169-185.

Sonneveld, C., De Kreij, C., 1996. Response of cucumber (*Cucumis sativus* L.) to an unequal distribution of salts in the root environment. *Plant and Soil* 209, 45-56.

Subbaiah, B.V., Asija, G.L., 1956. A rapid method for the estimation of available nitrogen in soil. *Current Science* 25, 258-260.

Sumithra, S., Ankalaiah, C., Rao, D., Yamuna, R.T., 2013. A case study on physico-chemical characteristics of soil around industrial and agricultural area of yerraguntla, kadapa district, AP, India. *International Journal Geology Earth and Environmental Sciences* 3, 28-34.

Tang, J., Sun, B., Cheng, R., Shi, Z., Luo D., Liu, S., Centritto, 2019. Effects of soil nitrogen (N) deficiency on photosynthetic N-use efficiency in N-fixing and non-N-fixing tree seedlings in subtropical China. *Scientific Reports* 9, 4604 <https://doi.org/10.1038/s41598-019-41035-1>

Thakre, Y.G., Choudhary, M.D., Raut, R.D., 2012. Physicochemical Characterization of Red and Black Soils of Wardha Region. *International Journal of Chemical and Physical Sciences* 1, 60-66.

Valente, D.S.M., Queiroz, D.M., Pinto, F., Santos, N.T., Santos, F.L., 2012. Definition of Management Zones in Coffee Production Fields Based on Apparent Soil Electrical Conductivity. *Scientia Agricola* 69, 173-179.

Walkley, A.J., Black, I.A., 1934. Estimation of soil organic carbon by the chromic acid titration method. *Soil Science* 37, 29-38.

Wang, D., Mathew, W.M., Jindong, S., Xiaohui, F., Fernando, M., Dokyoung, L., Michael C., 2012. Impact of nitrogen allocation on growth and photosynthesis of *Miscanthus* (*Miscanthus X giganteus*) *GCB Bioenergy* 4, 688–697.

Yosuf, M., Li J., Lu J., Ren, T., Cong R., Fahad, S., Li, X., 2017. Effects of fertilization on crop production and nutrient-supplying capacity under rice-oilseed rape rotation system. *Scientific reports* 7, 1-9.

UNDER PEER REVIEW