

# Standardization of media for propagation of Bougainvillea (*Bougainvillea* comm. Ex Juss.) : Root parameters

## ABSTRACT :

The present experiment entitled “Standardization of media for propagation of Bougainvillea (*Bougainvillea* comm. Ex Juss.)” was carried out at Horticultural nursery, College of Agriculture, IGKV, Raipur, (C.G) during the year 2019-20 and the experiment was conducted in Completely Randomized Design (CRD) with 16 treatments and 5 replications comprising of soilless media and biofertilizer Azotobacter. All the media treatments studied recorded superior vegetative and rooting attributes over control. Among the media treatments studied soil + sand + cocopeat recorded maximum sprouting percentage and survival percentage. All other Root attributes viz. The minimum days to rooting (144.80), The longest root length (6.08), highest number of roots per cutting (63.54), The maximum number of cutting attained rooting (24.00) discovered idealistic in media containing Soil + Sand + Cocopeat. The findings revealed that all increasing media treatments had a significant effect on all characters relevant to shoot and root growth and also the development of Bougainvillea. The findings revealed that all the characters linked root growth were greatly affected by all the growing media treatments and also the development of Bougainvillea. The maximum overall production cost has been seen in T6 (Soil + Sand + Cocopeat) and the least production cost in T13 (Soil + Azotobacter). Combination of rooting media has been beneficial to significantly increase the vegetative growth parameters and economics as compared to control in Bougainvillea.

Keywords : *Azotobacter*, cocopeat, Bougainvillea.

## INTRODUCTION

Bougainvillea is a genus of thorny ornamental vines, bushes, and trees belonging to the four o' clock family, Nyctaginaceae. This evergreen shrub has a variety of flower colours like Red, Orange, Pink, Purple, White etc. It is mainly grown in open field condition. They flower most heavily in winter and early spring, but some plants put forth scattered clusters all year. Depending on the variety, it can be grown as a bush, hedge, in pots, as a bonsai, as standard or semi standard shrub or can be trained in pergola or arches.

Likewise, other ornamental plants, the multiplication of Bougainvillea is done by using vegetative part *i.e.* Stem cutting, Layering, Budding. Stem cutting is regarded as the cheapest and easiest method of Bougainvillea; however the success rate of propagation is very limited. Rooting percentage or else we can say rooting responses varies from variety to variety. As we know, in most of ornamental plants, the success of rooting in cuttings is mainly depends upon variety, time of planting and use of growth regulators (Rowezak, 2001)<sup>[6]</sup>. Rainy season is the best time for planting Bougainvillea.

Use of different types of rooting media having different characteristics is important for quality rooting in cuttings. Though there is no universal rooting mix, different medias are used according to cutting types, different seasons etc. Highest rooting percentage (> 97%) was seen with basal stem cutting compared with terminal and middle cutting. Highest number of roots/cuttings ( 27.66) was seen with basal stem cuttings after 60 days of planting the cuttings. Highest plant height(67.66 cm) and number of leaves/cutting (46.00) was seen with soil+sand (1:1) after 90 days of planting the cuttings (Eed *et al.* 2015)<sup>[3]</sup>.

We are aware of the fact that, success rate of bougainvillea is very limited. Hence, to improve rooting in cuttings, several rooting hormones are widely used during propagation. The .The variety ‘Mahara’ is easily rooted with the help of auxin treatment. Growth hormones which are used are indole -3-butyric acid (IBA), 1-naphthalene acetic acid (NAA), indole -3-acetic acid (IAA). These hormones with different concentrations enhance the growth and development of roots.

## MATERIAL AND METHODS

The field experiment was performed during the year 2019-2020 at the Horticulture research cum Instructional Farm, College of Agriculture, IGKV, Raipur (C.G).

**Table 1 : Treatment Details:**

S.no	Treatments	Treatment details
1.	T <sub>1</sub>	Sand
2.	T <sub>2</sub>	Cocopeat
3.	T <sub>3</sub>	Rice husk
4.	T <sub>4</sub>	Sawdust
5.	T <sub>5</sub>	Soil + FYM + Sawdust
6.	T <sub>6</sub>	Soil + Sand + Cocopeat
7.	T <sub>7</sub>	Sand + FYM
8.	T <sub>8</sub>	Sand + Azotobacter
9.	T <sub>9</sub>	Soil + Cocopeat
10.	T <sub>10</sub>	Soil + Sand + Ricehusk
11.	T <sub>11</sub>	Soil + Sand
12.	T <sub>12</sub>	Cocopeat + Azotobacter
13.	T <sub>13</sub>	Soil + Azotobacter
14.	T <sub>14</sub>	Ricehusk + Azotobacter
15.	T <sub>15</sub>	Sawdust + Azotobacter
16.	T <sub>16</sub>	Soil (control)

After cutting from the mother plants, the hardwood cuttings (10-15 cm) of pencil thickness were prepared. The leaves were carefully trimmed from the shoots to prevent damage to the axillary buds. The cuttings were prepared to differentiate both ends during planting by giving a slanting cut on the upper end and a straight cut on the lower end. The cuttings were planted immediately in media mixtures after quick dip treatment of IBA @ 4000 ppm for (<5 sec.) directly in the polythene bags. 1g of IBA was first dissolved in a few drops of 0.1N NaOH for 4000 ppm of IBA solution and then up to 250 ml of distilled water was made up in order to achieve the right concentration of IBA. In the polybags with pre-punched bottom holes for drainage, various treatments containing different media were filled in. Various intercultural operations were carried out when needed. Various shoot and root attributes were noted and Statistically evaluated using Anova for Completely Randomized Design (CRD).

## RESULT AND DISCUSSION

The minimum days to rooting (144.80) was recorded with T6 (Soil + Sand + Cocopeat). Likewise, the maximum days to rooting (152.40) was recorded with control T16 (Soil). this can be because of cocopeat, which has low nitrogen, calcium and magnesium but it is high in phosphorus and potassium (Gohil *et al.* 2018)<sup>[4]</sup> in ornamental pot plants. Benefits of phosphorus in plants is that, it is good for the root growth and development.

The longest root length (6.08) was observed in T6 (Soil + Sand + Cocopeat). The least root length (3.60) was observed in T15 (Sawdust + *Azotobacter*) at 180 DAP.

The increase in the length of the cutting roots treated with growth regulators may be due to the accumulation of metabolism at the site of auxin application, cell enlargement, increased carbohydrate hydrolysis, new protein synthesis and auxin-induced cell division. The standard concentration of IBA combined with media increases the duration of the roots of bougainvillea, an effect recorded by Eed *et al.* (2015)<sup>[3]</sup>.

Data recorded at 180 DAP, treatment T6 (Soil + Sand + Cocopeat) showed highest number of roots per cutting (63.54). The lowest number of roots (39.78) was recorded with control T16 (Soil). These findings are in agreement with the reports of Nazari *et al.* (2011)<sup>[5]</sup> in hyacinth.

The maximum number of cutting attained rooting (24.00) was recorded in T6 (Soil + Sand + Cocopeat). The minimum number of cutting attained rooting (17.00) was recorded in Control T16 (Soil). Highest survival percentage was also found in T<sub>6</sub>. This may be attributed to metabolite aggregation at the site of auxin application, accelerated carbohydrate hydrolysis, new protein synthesis, auxin-induced cell elongation and cell division. Increases in the rooting percentage may be attributable to proper IBA concentration, high carbohydrates in turn, and low nitrogen levels contribute to further root forming (Carlson, 1929)<sup>[2]</sup>.




Fig 1. Graphical representation of effect of growing media on Bougainvillea (Root parameters)

Table 2. Effect of growing media on Bougainvillea

Tr. No.	Treatments	Days to rooting	Length of root 180 DAP	Number of roots 180 DAP
T1	Sand	145.80	4.14	42.24
T2	Cocopeat	149.80	4.18	51.04
T3	Ricehusk	145.40	4.42	57.73
T4	Sawdust	149.00	4.32	51.92
T5	Soil + FYM + Sawdust	147.60	5.24	63.36
T6	Soil + Sand + Cocopeat	144.80	6.08	63.54
T7	Sand + FYM	146.00	4.24	58.08
T8	Sand + <i>Azotobacter</i>	145.80	4.58	49.98
T9	Soil + Cocopeat	148.60	4.58	51.22
T10	Soil + Sand + Ricehusk	145.00	4.02	52.27
T11	Soil + Sand	146.00	4.26	52.10
T12	Cocopeat + <i>Azotobacter</i>	147.40	4.30	49.98
T13	Soil + <i>Azotobacter</i>	149.80	4.42	52.80
T14	Ricehusk + <i>Azotobacter</i>	146.00	2.40	55.62
T15	Sawdust + <i>Azotobacter</i>	147.00	3.60	46.29
T16	Soil (control)	152.40	4.54	39.78
	SEm ±	1.55	0.30	3.48
	C.D @ 5%	4.38	0.87	9.82

Table 3. Effect of growing media on Bougainvillea

Tr. No.	Treatments	Number of cutting attained rooting	Survival %
T1	Sand	19	64

T2	Cocopeat	20	68
T3	Ricehusk	19	64
T4	Sawdust	20	64
T5	Soil + FYM + Sawdust	22	80
T6	Soil + Sand + Cocopeat	24	88
T7	Sand + FYM	23	84
T8	Sand + <i>Azotobacter</i>	22	84
T9	Soil + Cocopeat	19	68
T10	Soil + Sand + Ricehusk	23	84
T11	Soil + Sand	19	68
T12	Cocopeat + <i>Azotobacter</i>	22	80
T13	Soil + <i>Azotobacter</i>	19	68
T14	Ricehusk + <i>Azotobacter</i>	23	80
T15	Sawdust + <i>Azotobacter</i>	22	80
T16	Soil (control)	17	56
	SEm ±	0.28	
	C.D @ 5%	0.78	

## CONCLUSION

As cocopeat contains 20 percent to 30 percent fibre of varying length, it is considered as a good growing media. Cocopeat is high in phosphorus and potassium (Gohil *et al.* 2018)<sup>[4]</sup>. Benefits of phosphorus in plants is that, it is good for the root growth and development. Whereas, potassium helps in strong build up of stems. It holds upto 8-9 times its weight in water, hence it has high water holding capacity which may cause poor aeration within the media, thus affecting the oxygen diffusion to the roots. The solution to this problem is to incorporate coarser material into cocopeat could improve the aeration status of the media (Awang *et al.* 2009)<sup>[1]</sup>.

## FUTURE SCOPE

As there is scarcity of land, hence the supply of quality topsoil is declining rapidly. Thus, Promoting the use of soil-free media in ornamental crop production.

## REFERENCE

1. Awang, Y., Shaharom, A.S., Mohamad, R.B. and Selamat, A. 2009. Chemical and physical characteristics of cocopeat-based media mixtures and their effects on the growth and development of *Celosia cristata*. AM. J. Agri. & Biol. Sci., 4(1): 63-71.
2. Carlson, M.C. 1929. Microchemical studies of rooting and non-rooting rose cuttings. Bot. Gaz., 87(1): 64.
3. Eed, A.M., Albana'a, B. and Almaqtari, S. 2015. The effect of growing media and stem cutting type on rooting and growth of *Bougainvillea spectabilis* plants. Univ. Aden J. Nat. and Appl. Sc., 19(1): 141-147.
4. Gohil, P., Gohil, M., Rajatiya, J., Halepotara, F., Solanki, M., Malam, V.R. and Barad, R. 2018. Role of growing media for ornamental pot plants. Int. J. Pure App. Biosci., 6(1): 1219-1224.
5. Nazari, F., Farahmand, H., Khos-Khui, M. and Salehi, H. 2011. Effects of coir as a component of potting media on growth, flowering and physiological characteristics of hyacinth. Int. J. of Agric. and Food Sci., 1(2): 34-38.
6. Rowezak, M.M.A. 2001. Response of some Ornamental plants to Treatment with Growth substances. Sc. Thesis, Fac. Agric., Cairo Univ.