

Original Research Article

Nutritional composition, phytochemistry, and in vivo potentials of *Thaumatococcus daniellii* (Benn.) rhizome extracts

ABSTRACT

The evidence on the advantages of plant rhizomes remains sparse. Proximate analysis, phytochemical screening (qualitative and quantitative), and lethal dose investigation of the aqueous and ethanolic extracts of the unpeeled and peeled *Thaumatococcus daniellii* (Benn.) rhizome (TdR) and their effects in albino rats are presented in this paper. In order to investigate the extract's effect on albino rats, fifteen animals were divided into five groups (n = 3). Group 1 is the control, and groups 2–5 were orally administered 300 mg/kg body weight of aqueous and ethanolic extracts of unpeeled and peeled TdR for 7 days, respectively. After the experimental period, blood glucose, serum total protein, albumin, globulin, cholesterol, and triglyceride concentrations were determined spectrophotometrically. The qualitative analysis carried out revealed the presence of flavonoids, reducing sugar, free anthraquinones, cardiac glycosides, and glycosides in both aqueous and ethanolic extracts of the unpeeled and peeled rhizomes. The quantitative analysis shows that total phenol has the highest percentage of constituents compared to niacin, flavonoids, and tannins. Both qualitative and quantitative analyses revealed the absence of alkaloids and phlobatannins. The blood glucose concentration was significantly ($P \leq .05$) decreased in animals administered with all the crude rhizome extracts, while the total serum albumin, globulin, and protein concentrations were significantly ($P \leq .05$) increased. Likewise, the extracts of the peeled and unpeeled rhizomes caused a significant ($P \leq .05$) increase in serum cholesterol concentration as well as triglycerides compared to the control. The results revealed that the nutritional composition and phytochemistry of both the aqueous and ethanolic extracts of the unpeeled and peeled TdR rhizome and the administration of 300 mg/kg body weight had significant effects on the biochemical parameters of the rat.

Keywords: Glucose, flavonoids, cholesterol, protein, peeled, unpeeled

1. INTRODUCTION

Thaumatococcus daniellii (Td) (Benn.) is an underutilized rhizomatous plant found in the tropical rain forests and coastal areas of West Africa. In Nigeria, it is called “Ewe-Eran” or “Adundunmitan”, and its stalk, leaves, fruits, and rhizomes contribute to the economy of the rural people. It is commonly referred to as miracle fruit or berry, serendipity berry, sweet prayer plant, soft cane, and katempfe. Td, a perennial monocotyledonous herb, propagates itself by rhizomes and forms an undergrowth of forest trees in its natural habitat [1-9]. The aril of its fruit is a natural source of a sweet protein called thaumatin, which is about 3000 times sweeter than sucrose solution (8-10%) [1,7,8,10]. The removal of thaumatin from the arils of Td fruits leads to the generation of waste (seed, pericarp, and pulp), which has been shown to constitute over 93% of fruit weight and found to have no significant difference in the biological value, net protein utilization, and protein efficiency ratio value between the casein base diet and the Td base diet; therefore, they can be used as livestock feed [7].

In the traditional and folkloric management of some diseases and ailments, the different parts of Td are employed. Several researchers worked on the leaves, fruits, fruit waste (seed, pericarp, and pulp), and stalks of the Td plant. It has been found that these parts have medicinal properties such as preservative, antioxidant, antimicrobial, and anti-diabetic, to mention a few [2,7,11-14]. There is a paucity of scientific data on the medicinal properties of the Td rhizome, which is a part of the plant. Therefore, this study is to conduct research on the aqueous and ethanolic extracts of unpeeled and peeled Td rhizomes by investigating their nutritional composition, phytochemistry, lethal dose, and in vivo effects.

2. MATERIAL AND METHODS

2.1 Plant material and sample preparation .

The rhizomes of the *Thaumatococcus daniellii* (Td) plant were collected from a farm at Lusada village in Ogun State, Nigeria, and validated by the Department of Botany, Lagos State University, Ojo, Lagos. The rhizomes were rinsed carefully with tap water to remove the dirt and divided into two equal parts. One part was peeled and the other unpeeled (Fig. 1); both parts were cut into pieces for aqueous and ethanolic extraction. Briefly, 100 g of unpeeled and peeled rhizomes of Td were extracted with 70% ethanol for 4 hours using the Soxhlet extractor apparatus. Another 100 g of unpeeled and peeled samples were soaked in 250 cm³ of distilled water for 12 hours and filtered using Whatman filter paper No. 42 (125 mm) to get the aqueous extract. The crude extracts were collected into sterile bottles and stored until further use.



Figure 1. The unpeeled (A) and peeled (B) rhizomes of *Thaumatococcus daniellii*

2.2 Proximate analysis

The Association of Official Analytical Chemists' recommended method was used to determine the TdR's proximate composition (moisture, crude protein, crude fat, and ash) [15]. The difference was used to compute carbohydrate content.

2.3 Phytochemical screening

2.3.1 Qualitative analysis

The determination of tannins, phlobatanins, reducing sugar, glycosides, alkaloids, cardiac glycosides, saponins, flavonoids, bound anthraquinone, anthracyanides, terpenoids, and free anthraquinone in the crude aqueous and ethanolic extracts of unpeeled and peeled TdR was performed according to the methods of Ogunrinola *et al.* [16] and Kalaichelvi and Dhivya [17].

2.3.2 Quantitative analysis

The estimation of total phenolic content (TPC), total flavonoid content (TFC), niacin content (NC), tannin content (TC), and alkaloid content (AC) was performed in the aqueous and

ethanolic extracts of peeled and unpeeled TdR as described by the modified methods of Ogunrinola *et al.* [16], Ghasemi *et al.* [18], Iqbal *et al.* [19], and Chlopicka *et al.* [20].

2.4 Acute toxicity studies (Lethal dose (LD₅₀))

The modified method of Lorke [21], as modified by the method of Adu *et al.* [2], was used to estimate the acute toxicity (LD₅₀) of the aqueous and ethanolic extracts of unpeeled and peeled TdR when administered orally. Briefly, fifty-six rats were allotted equally into twenty-eight (n = 2) well-ventilated plastic cages. The first 7 cages were administered with an aqueous extract of unpeeled TdR; the second 7 cages were administered with an aqueous extract of peeled TdR; the third set of cages were administered with an ethanolic extract of unpeeled TdR; and the last 7 cages were given an ethanolic extract of peeled TdR orally at dosage levels of 50 mg/kg, 100 mg/kg, 150 mg/kg, 200 mg/kg, 250 mg/kg, 300 mg/kg, and 350 mg/kg body weight, respectively, for 72 hours.

2.5 Animals and Experimental Procedure

A total of fifteen (15) Wistar male albino rats weighing between 120 - 150 g were used for the experiments. The animals were housed in well ventilated cages to acclimatize for a week and allowed to drink and feed freely. The animals were randomly and equally distributed into five groups (n = 3) as follows: aqueous and ethanolic extracts of TdR.

Group I	Control rats, given distilled water
Group II	Oral administration of 300 mg/kg body weight of unpeeled aqueous extract of TdR
Group III	Oral administration of 300 mg/kg body weight of peeled aqueous extract of TdR
Group IV	Oral administration of 300 mg/kg body weight of unpeeled ethanolic extract of TdR
Group V	Oral administration of 300 mg/kg body weight of peeled ethanolic extract of TdR

After seven days of administration, the animals were fasted overnight and sacrificed under light ketamine anaesthesia. Blood was collected via cardiac puncture and separated into serum, which was stored for analysis.

2.6 Biochemical analysis

Blood glucose level, serum total protein, albumin, and globulin determination

The blood glucose level was measured before and after treatment by taking a blood sample from the tail. Blood was dripped on the end of the strip, and blood glucose was read on the glucometer after 10 seconds. The serum total protein and albumin were analyzed using the commercial Randox kits, products of Randox Laboratories, U.K. And serum globulin was determined as the difference between serum total protein and albumin.

Lipid profile determination concentrations of total cholesterol and triglycerides in the serum were determined with commercial kits (Spin React S.A., Santa Colona, Sant Esteve de Bas, Spain).

2.7 Statistical analysis

The results are presented as mean \pm standard error of mean (SEM) and were analyzed for statistical significance by one-way analysis of variance (ANOVA). The values with $P \leq .05$ were considered statistically significant.

3. RESULTS AND DISCUSSION

3.1 Proximate analysis

The proximate analysis is used for the estimation of the quantitative properties of food substances, including moisture, crude protein, total fat, ash, total carbohydrate, and dietary fibre [15,23]. Table 1 represents the proximate composition of the unpeeled and peeled TdR. The results revealed that while crude protein content has the lowest percentage of 0.13 and 0.94 in both unpeeled and peeled rhizome samples, the percentage of carbohydrate content is higher (76.75% and 88.53%), respectively. The moisture content of a sample is the amount of water lost during drying and serves as one of the main factors in storage, which may be due to the proliferation of microorganisms [24].

Table 1. Proximate analysis of unpeeled and peeled *Thaumatococcus daniellii* (Benn.) rhizomes (TdR) (g per 100g)

Rhizome samples	Moisture Content (%)	Ash Content (%)	Crude Protein Content (%)	Crude Fat Content (%)	Carbohydrate Content (%)
unpeeled	1.94	4.78	0.13	16.40	76.75
peeled	1.05	4.88	0.94	4.60	88.53

In this study, the percentage moisture content of unpeeled TdR was higher than that of peeled TdR samples, which might be due to the removal of the rhizome bark. The reduced moisture content shows that the peeled TdR samples can withstand microorganism growth and extend the shelf-life of TdR, which means that it has environmental benefits. The ash content is the amount of total mineral (inorganic) residue leftover after the combustion of organic matter in a food sample until it reaches a constant weight [25]. The result revealed a higher percentage of ash content in the peeled TdR than in the unpeeled TdR samples. This enhancement of the ash content might be due to the presence of more fibres and veins in the peeled TdR samples [26].

The percentage crude protein is the amount of total nitrogen (protein nitrogen and a few non-protein nitrogens) multiplied by protein factors [24]. In this study, the TdR samples used a 6.25 protein factor to convert nitrogen to protein and then found that it was higher in the peeled samples. This is similar to the report of Pazhanichamy *et al.* [27]. Although the observed value is lower than the value, which makes the rhizomes less advantageous as a rich source of plant protein for humans, they can contribute to the formation of hormones that control growth, repair, and maintenance of the plant. Total fat is the amount of fatty acids, fat-soluble vitamins, and steroids in a food sample [24]. Total fat in this study was found to be higher in unpeeled TdR samples, indicating that it can help with the transportation and absorption of fat-soluble vitamins such as vitamins A, D, E, and K and provide energy to the plant. Carbohydrates are the main components of the structural materials (cell walls, cell sap, and protoplasm) in plants.

The carbohydrates produced by the leaves were translocated to the rhizomes, which serve as storage organs [24,28]. In this study, carbohydrate was calculated based on the difference method, and the total carbohydrate of TdR samples was found to be higher in peeled samples. This is in accordance with the study carried out by Pazhanichamy *et al.* [27]. This shows that the natural structure (cell walls and protoplasm) of the rhizome is still maintained [28]. These results differ from the report of Oforibika *et al.* [29] and may be due to the geographical location where the samples were collected and the research protocols.

3.2 Phytochemical screening

The detection of active principles in medicinal plants plays an important role in regards to their potential pharmacological effects [28]. Appropriate solvents, including organic and/or aqueous solutions, have been reported for extracting active compounds. Because of the

polarity differences between water and ethanol, they are often recommended for extract preparation [30-32]. The results of phytochemical constituents in the aqueous and ethanolic extracts of unpeeled and peeled TdR are depicted in Table 2. Qualitative analysis depicts the presence of saponins, flavonoids, reducing sugar, free anthraquinone, cardiac glycosides, and glycosides in the aqueous extract of unpeeled and peeled TdR. And the presence of tannins, flavonoids, reducing sugar, free anthraquinone, bound anthraquinones, anthacyanides, terpenoids, cardiac glycosides, and glycosides in the ethanolic extract of unpeeled and peeled TdR, respectively. This may be due to the high polarity of the ethanol used for the extraction.

In both the unpeeled and peeled aqueous and ethanolic extracts, the quantitative analysis (percentage of crude extract) showed the presence of tannins, niacin, total phenol, and flavonoids but the absence of alkaloids. It was observed that total phenol has the highest percentage, whereas tannins have the lowest percentage, respectively, in all the samples. The results that were observed are comparable to those that Majaw and Moirangthem [28] and Taoheed *et al.* [32] reported. It has been reported that these bioactive constituents are suggested to be associated with antibacterial, antidiarrheal, antioxidant, and antiviral activity [28,34,35]. The high amount of reducing sugar will help in the central metabolic pathways and in the production of secondary metabolites that enhance the medicinal properties of plants [36,37].

Table 2. Phytochemical constituents of aqueous and ethanolic of unpeeled and peeled *Thaumatococcus daniellii* (Benn.) rhizomes (TdR)

Phytochemicals	Groups			
	Unpeeled aqueous extract of TdR	Peeled aqueous extract of TdR	Unpeeled ethanolic extract of TdR	Peeled ethanolic extract of TdR
Qualitative analysis				
Saponins	++	+	nd	nd
Tannins	nd	nd	+	+
Flavonoids	++	+	+	+
Reducing sugar	+	++	+++	++++
Free anthraquinone	++	+	+	+
Bound anthraquinone	nd	nd	+	+
Phlobatannins	nd	nd	nd	nd
Anthacyanides	nd	nd	+	+
Terpenoids	nd	nd	+	+
Cardiac glycosides	+	+	+	+
Glycosides	+	+	++	++
Alkaloids	nd	nd	nd	nd
Quantitative analysis: Percentage of crude extracts				
Tannins	0.08	0.06	0.15	0.28
Niacin	1.13	0.36	0.39	0.69
Total Phenol	5.68	3.65	1.49	3.58
Flavonoid	1.86	0.22	0.67	0.31
Alkaloids	nd	nd	nd	nd

TdR = *Thaumatococcus daniellii* (Benn.) rhizomes; ++++ = Present in very highly concentration; +++ = Present in very high concentration; ++ = Present in moderately high concentration; + = Present in trace concentration; -nd = Not detected

3.3 Acute toxicity studies (Lethal dose (LD₅₀))

The results of acute toxicity studies show that there was no mortality within 72 hours after the oral administration of 50 mg/kg, 100 mg/kg, 150 mg/kg, 200 mg/kg, 250 mg/kg, 300 mg/kg, and 350 mg/kg body weight of aqueous and ethanolic extracts of unpeeled and peeled TdR, respectively. Therefore, our research team used 300 mg/kg body weight for the animal study.

3.4 Determination of blood glucose, serum total protein, albumin, globulin, cholesterol, and triglyceride concentrations

There was a significant ($P \leq .05$) reduction in blood glucose concentrations after the administration of aqueous and ethanolic extracts of unpeeled and peeled TdR compared to the control. The total serum albumin, globulin, and protein concentrations were found to increase significantly ($P \leq .05$) in all groups compared to the control (Table 3). The administration of all the aqueous and ethanolic extracts of unpeeled and peeled TdR caused a significant ($P \leq .05$) increase in serum cholesterol concentration as well as triglycerides compared to control (Table 4). The highest levels of cholesterol and triglycerides increase more with the administration of the peeled ethanolic extract of TdR.

This study observed up- and down-regulation of total protein, albumin, globulin, blood glucose, cholesterol, and triglyceride concentrations in the aqueous and ethanolic extracts of unpeeled and peeled TdR, respectively (Table 4). These alterations are due to the phytochemical constituents present in TdR. Their effects on total protein and globulin may be due to an alternation in the intracellular protein synthesis mechanism and immunoglobulin production. The increased concentration of albumin may be due to high protein formation from the alimentary tract or protein in the liver [34,38,39]. The reduction in the glucose concentration revealed that TdR possesses a glucose-lowering effect, which may be partly due to the action of the flavonoid, thereby modulating some cell signalling like glucose uptake in the intestine [34,40-42].

Table 3. Effects of *Thaumatococcus daniellii* (Benn.) rhizomes (TdR) extracts on blood glucose, serum total protein, albumin and globulin concentrations in the animal

Groups	Blood glucose (mg/dl)		Total protein (g/dl)	Albumin (g/dl)	Globulin (g/dl)
	Before	After			
I	29.35 ± 2.11 ^a	35.35 ± 2.11 ^a	7.10 ± 0.10 ^a	1.94 ± 0.05 ^a	5.16 ± 0.04 ^a
II	35.20 ± 1.62 ^b	31.25 ± 1.72 ^b	11.40 ± 1.20 ^b	5.01 ± 0.26 ^b	6.39 ± 0.98 ^b
III	74.75 ± 1.70 ^c	73.75 ± 1.72 ^c	14.1 ± 0.30 ^c	4.42 ± 0.46 ^c	9.68 ± 0.66 ^c
IV	38.61 ± 1.80 ^d	28.63 ± 2.86 ^d	10.20 ± 0.20 ^d	4.45 ± 0.30 ^d	5.75 ± 0.28 ^d
V	99.77 ± 2.52 ^e	88.75 ± 2.30 ^e	13.5 ± 0.30 ^e	5.67 ± 0.13 ^e	7.83 ± 0.40 ^e

I = Control; II = 300 mg/kg body weight of unpeeled aqueous extract; III = 300 mg/kg body weight of peeled aqueous extract; IV = 300 mg/kg body weight of peeled ethanolic extract; V = 300 mg/kg body weight of peeled ethanolic extract. Values are represented as the mean ± S.E.M for 3 rats in each group. Values having different superscripts within a column differ significantly from each other ($P \leq .05$).

Likewise, tannins inhibited the activation of α -amylase and α -glucosidase activities to reduce the blood glucose concentration by phosphorylation of the insulin receptor and translocation of glucose transporter 4, and also inhibited the important genes for adipogenesis [43,44]. Phenols and terpenoids also inhibit the transcription factor associated with the activation of genes involved in the biosynthesis of cholesterol, fatty acids, and triglycerides, thereby increasing glycogenesis and decreasing glycogenolysis, as well as inhibiting aldose reductase [45-48]. The glycosides and cardiac glycosides mediate hypoglycaemic activity by

increasing insulin secretion through adenosine monophosphate-activated protein kinase [49,50].

Table 4. Effects of *Thaumatococcus daniellii* (Benn.) rhizomes (TdR) extracts on serum cholesterol and triglyceride concentrations in the animal

Groups	Cholesterol Concentration (mg/dl)	Triglyceride Concentration (mg/dl)
I	69.70 ± 6.71 ^a	105.54 ± 4.20 ^a
II	86.36 ± 14.05 ^b	89.54 ± 9.86 ^b
III	77.82 ± 13.64 ^c	106.77 ± 6.70 ^c
IV	102.12 ± 15.97 ^d	117.54 ± 5.73 ^d
V	75.76 ± 19.19 ^e	99.85 ± 5.20 ^e

I = Control; II = 300 mg/kg body weight of unpeeled aqueous extract; III = 300 mg/kg body weight of peeled aqueous extract; IV = 300 mg/kg body weight of peeled ethanolic extract; V = 300 mg/kg body weight of peeled ethanolic extract. Values are represented as the mean ± S.E.M for 3 rats in each group. Values having different superscripts within a column differ significantly from each other ($P \leq .05$).

4. CONCLUSION

The present study investigated the phytochemical constituents of TdR, determined its acute toxicity and effects on blood glucose, total protein, albumin, globulin, cholesterol, and triglyceride concentrations. This is the first scientific report on the phytochemical constituents and health benefits of aqueous and ethanolic extracts of unpeeled and peeled TdR. The presence of phytochemicals acts through a number of diverse mechanisms to bring about some potential health benefits of TdR for humans and animals. Further research is needed to investigate the effectiveness of the aqueous and ethanolic extracts of the unpeeled and peeled TdR against degenerative diseases. And there should be awareness of the potential of TdR for animal nutrition and medical purposes.

Ethical Approval

All experiments were performed in compliance with the principles for an ethical guide for the care and use of laboratory animals [22], which were approved by the Animal Ethical Committee of the Department and the University.

REFERENCES

1. Fadahunsi O, Adegbola P, Olorunnisola S, Akinloye O. Phytochemistry, nutritional composition, and pharmacological activities of *Thaumatococcus daniellii* (Benth): A review. *BioTechnologia*. 2021;102(1):101-17. <https://doi.org/10.5114/bta.2021.103766>
2. Adu OB, Adeyemo GA, Falua OB, Fajana OO, Ogunrinola OO, Saibu GM, *et al*. The Effect of *Thaumatococcus daniellii* Leaf Extracts on Immunological and Oxidative Stress Markers in Rat. *Asian J. Biochem. Genet. Mol. Biol.* 2021; 7(4): 6-14. <https://doi.org/10.9734/AJBGMB/2021/v7i430179>
3. Chinedu SN, Iheagwam FN, Anichebem CJ, Ogunnaike GB, Emiloju OC. Antioxidant and biochemical evaluation of *Thaumatococcus daniellii* seeds in rat. *J. Bio. Sci.* 2017; 17(8): 381-387. <https://doi.org/10.3923/jbs.2017.381.387>

4. Chinedu SN, Oluwadamisi AY, Popoola ST, David BJ, Epelle T. Analyses of the leaf, fruit and Seed of *Thaumatococcus daniellii* (Benth.): Exploring potential uses. Pak. J. Biol. Sci. 2014; 17(6): 849-854. <https://doi.org/10.3923/pjbs.2014.849.854>
5. Ogoloma UJ, Wegu M, Abbey BN. Phytochemical Analysis and Effect of Methanolic Root/Leaf Extracts of *Thaumatococcus danielli* on some Biochemical Parameters in Wistar Rat. Academ. Arena. 2017; 9(6): 64-74. <https://doi.org/10.7537/marsaaj090617.10>.
6. Ogoloma UJ, Wegu M, Abbey BN. Haematological Effects of Methanolic Root and Leaf Extracts of *Thaumatococcus danielli* in Wistar Rat. Biomed. Nursing. 2017; 3(3): 42-55. <https://doi.org/10.7537/marsbnj030317.05>
7. Elemo BO, Adu OB, Ogunrinola OO, Efuwape TO, Olaleye KO, Kareem AA. Biological evaluation of *Thaumatococcus danielli* waste protein. Pak. J. Nutr. 2011; 10: 1048-1052.
8. Elemo BO, Elemo GN, Agboola OO, Oyedun AB. Studies on some anti-nutritive factors and in-vitro protein digestibility of *Thaumatococcus danielli* (Benth) wastes. Niger. J. Biochem. Mol. Biol. 2001; 16: 43-46.
9. Raimi OG, Elemo BO, Fatai AA, Bankole HA, Kazeem MI, Banjoko AO. Isolation and partial characterization of a protease enzyme from *Thaumatococcus daniellii* waste. African J. Biotech. 2011; 10(16): 3186-3190. <http://doi.org/10.5897/AJB10.2065>
10. Ubani CD, Uko OE, Wariso CA, Kalu AA. Evaluation of the nutrient composition and hepatotoxic potential of *Thaumatococcus daniellii*. GSC Biol. Pharm. Sci. 2022; 18(3): 011-015. <https://doi.org/10.30574/gscbps.2022.18.3.0051>
11. Shalom NC, Franklyn NI, Makinde BT, Thorpe BO, Emiloju OC. Data on in vivo antioxidant, hypolipidemic and hepatoprotective potential of *Thaumatococcus daniellii* (Benn.) Benth leaves. Data in Brief. 2018; 20: 364-370. <https://doi.org/10.1016/j.dib.2018.08.016>
12. Adeogun O, Adekunle A, Ashafa A. Chemical composition, lethality and antifungal activities of the extracts of leaf of *Thaumatococcus daniellii* against foodborne fungi. Beni-Suef Univ. J. Basic Appl. Sci. 2016; 5: 356-368. <https://doi.org/10.1016/j.bjbas.2016.11.006>
13. Olorunnisola OS, Adetutu A, Owoade AO, Okoh OO, Oyewo EB, Adegbola P. Ethnopharmacological and in-vitro anti-diabetic study of some medicinal plants commonly used in Ogbomoso, South Western Nigeria. J. Appl. Biosci. 2016; 105: 10064-10084. <http://doi.org/10.4314/jab.v105i1.3>
14. Shalom NC, Adetayo YO, Samuel TP, Bolaji JD, Tamuno E. Analyses of the leaf, fruit and seed of *Thaumatococcus daniellii* (benth). Exploring potential uses. Pak. J. Biol. Sci. 2014; 17(6): 849-854. <https://doi.org/10.3923/pjbs.2014.849.854>
15. AOAC (Association of Official Analytical Chemists) Official Methods of Analysis of Association of Analytical Chemists international, 17th ed. Horwitz, W. (ed). Vol I and II. AOAC International Publs, Maryland USA. 2000; Ch. 45: 12-20.
16. Ogunrinola OO, Odulate JE, Elemo BO. Proximate analysis and pharmacognostical investigation of some medicinal plants: *Naudea latifolia*; *Morinda lucida*; *Alstonea congestis* and *Anchornea cordifolia*. J. Res. Rev. Sci. 2004; 3: 190-193.
17. Kalaichelvi K, Dhivya SM. Phytochemical screening and antibacterial activity of leaf extract of *Martynia annua*, L. and *Premna latifolia*, Roxb. J. Med. Plants Studies. 2016; 4(4): 84-87. ISSN 2320-3862
18. Ghasemi Pirbalouti A, Siahpoosh A, Setayesh M, Craker L. Antioxidant activity, total phenolic and flavonoid contents of some medicinal and aromatic plants used as herbal teas and condiments in Iran. J. Med. food. 2014; 17(10): 1151-1157. <https://doi.org/10.1089/jmf.2013.0057>

19. Iqbal E, Salim KA, Lim LB. Phytochemical screening, total phenolics and antioxidant activities of bark and leaf extracts of *Goniothalamus velutinus* (Airy Shaw) from Brunei Darussalam. J. King Saud University-Science. 2015; 27(3): 224-232. <https://doi.org/10.1016/j.jksus.2015.02.003>
20. Chlopicka J, Pasko P, Gorinstein S, Jedryas A, Zagrodzki P. Total phenolic and total flavonoid content, antioxidant activity and sensory evaluation of pseudocereal breads. LWT-Food Sci. Technol. 2012; 46(2): 548-555. <https://doi.org/10.1016/j.lwt.2011.11.009>
21. Lorke D. A new approach to practical acute toxicity testing. Arch. Toxicol. 1983; 54: 275-287.
22. National Institutes of Health, NIH publication (1985) Guide for the care and use of laboratory animals. National Academies Press.
23. Puwastien P, Siong TE, Kantasubrata J, Craven G, Feliciano RR, Judprasong K. Asean Manual of Food Analysis. (1st ed.) Thailand: Institute of Nutrition, Mahidol University. 2011; 196.
24. Ganogpichayagrai A, Suksaard C. Proximate composition, vitamin and mineral composition, antioxidant capacity, and anticancer activity of *Acanthopanax trifoliatum*. J. Adv. Pharm. Technol. Res. 2020; 11(4): 179-183. https://doi.org/10.4103/japtr.JAPTR_61_20
25. Harris GK, Marshall MR. Ash analysis. Food analysis. 2017; 287-297. https://doi.org/10.1007/978-3-319-45776-5_16
26. Oduntan AO, Olaleye O, Akinwande BA. Effect of plant maturity on the proximate composition of *Sesamum radiatum Schum* leaves. J. Food studies.(2012; 1(1): 69-76. <https://doi.org/10.5296/jfs.v1i1.1806>.
27. Pazhanichamy K, Pavithra S, Rubini S, Lavanya B, Ramya I, Eevera T. Morphological, anatomical and proximate analysis of leaf, root, rhizome of *Costus igneus*. J. Pharm. Res. 2010; 3(4): 747-752. ISSN: 0974-6943
28. Majaw S, Moirangthem J. Qualitative and quantitative analysis of *Clerodendron colebrookianum Walp.* leaves and *Zingiber cassumunar Roxb.* rhizomes. Ethnobotanical Leaflets. 2009; (5),13: 578-589. <https://opensiuc.lib.siu.edu/eb/vol2009/iss5/3>.
29. Oforibika GA, Ogoloma JU, Tamunodiepriye E. Potential of *Thaumatococcus daniellii* in animal nutrition. Nature Sci. 2017; 15(10): 97-100. <https://doi.org/10.7537/marsnsj151017.13>
30. Abarca-Vargas R, Peña Malacara CF, Petricevich VL. Characterization of chemical compounds with antioxidant and cytotoxic activities in *bougainvillea x buttiana holttum* and *standl*, (Var. rose) extracts. Antioxidants. 2016; 5(4): 45. <https://doi.org/10.3390/antiox5040045>
31. Humaira F, Khan K, Zia M, Ur-Rehman T, Mirza B, Haq IU. Extraction optimization of medicinally important metabolites from *Datura innoxia* Mill: An in vitro biological and phytochemical investigation. BMC Complement. Altern. Med. 2015; 15: 376. <https://doi.org/10.1186/s12906-015-0891-1>
32. Anwar F, Przybylski R. Effect of solvents extraction on total phenolics and antioxidant activity of extracts from flaxseed (*Linum usitatissimum* L.). Acta Sci. Pol. Technol. Aliment. 2012; 11(3): 293-301.
33. Taoheed A, Tolulope A, Saidu A, Odewumi O, Sunday R, Usman M. Phytochemical properties, proximate and mineral composition of *Curcuma longa* Linn. and *Zingiber officinale* Rosc.: A comparative study. J. Scientific Res. Reports. 2017; 13(4): 1-7. <https://doi.org/10.9734/JSRR/2017/32623>

34. Chukwuma ER, Obioma N, Christopher OI. The phytochemical composition and some biochemical effects of Nigerian tigernut (*Cyperus esculentus* L.) tuber. *Pak. J. Nutr.* 2010; 9(7): 709-715.
35. Ayodeji OI, Adeleye O, Dada O, Adeyemi O, Anyasor GN. Phytochemical constituent and antioxidant activity of *Thaumatococcus daniellii* (Benth.) leaves (food wrapper). *Int. J. Pharmacol. Phytochem. Ethnomed.* 2016; 2: 55-61.
<https://doi.org/10.18052/www.scipress.com/IJPPE.2.55>
36. Khatri D, Chhetri SBB. Reducing sugar, total phenolic content, and antioxidant potential of nepalese plants. *BioMed. Res. Inter.* 2020. <https://doi.org/10.1155/2020/7296859>
37. Arsenault PR, Vail DR, Wobbe KK, Weathers PJ. Effect of sugars on artemisinin production in *Artemisia annua* L.: transcription and metabolite measurements. *Molecules.* 2010; 15(4): 2302-2318. <https://doi.org/10.3390/molecules15042302>
38. Selvakumar K, Bavithra S, Suganya S, Ahmad Bhat F, Krishnamoorthy G, Arunakaran J. Effect of quercetin on haematobiochemical and histological changes in the liver of polychlorinated biphenyls-induced adult male wistar rats. *J. biomarkers.* 2013. <https://doi.org/10.1155/2013/960125>
39. Olorunnisola OS, Bradley G, Afolayan AJ. Protective effect of *T. violacea* rhizome extract against hypercholesterolemia-induced oxidative stress in Wistar rats. *Molecules.* 2012; 17(5): 6033-6045. <https://doi.org/10.3390/molecules17056033>
40. Aba PE, Asuzu IU. Mechanisms of actions of some bioactive anti-diabetic principles from phytochemicals of medicinal plants: A review. *Indian Journal of Natural Products and Resources.* 2018; 9(2): 85-96. <http://nopr.niscpr.res.in/handle/123456789/44904>
41. Sangeetha KS, Umamaheswari S, Reddy CUM, Kalkura SN. Flavonoids: Therapeutic potential of natural pharmacological agents. *Inter. J. Pharm. Sci. Res.* (2016; 7(10): 3924-3930. [http://doi.org/10.13040/IJPSR.0975-8232.7\(10\).3924-30](http://doi.org/10.13040/IJPSR.0975-8232.7(10).3924-30)
42. Ramachandran V, Baojun X. Antidiabetic properties of dietary flavonoids: A cellular mechanism review. *Nutr. Metabol.* 2015; 12: 1-20. <https://doi.org/10.1186/s12986-015-0057-7>
43. Kunyanga CN, Imungi JK, Okoth M, Momanyi C, Biesalski HK, Vadivel V. Antioxidant and anti-diabetic properties of condensed tannins in acetonitrile extract of selected raw and processed indigenous food ingredients from Kenya. *J. Food Sci.* 2011; 76(4): 560-567. <https://doi.org/10.1111/j.1750-3841.2011.02116.x>
44. Liu X, Kim J, Li Y, Liu F, Chen X. Tannic acid stimulates glucose transport and inhibits adipocyte differentiation in 3T3-L1 cells. *J. Nutr.* 2005; 135(2): 165-171. <https://doi.org/10.1093/jn/135.2.165>
45. Panigrahy SK, Bhatt R, Kumar A. Targeting type II diabetes with plant terpenes: The new and promising antidiabetic therapeutics. *Biologia.* 2021; 76(1): 241-254.
<http://doi.org/10.2478/s11756-020-00575-y>
46. Zhao C, Yang C, Wai STC, Zhang YP, Portillo M, Paoli P, *et al.* Regulation of glucose metabolism by bioactive phytochemicals for the management of type 2 diabetes mellitus. *Crit. Rev. Food Sci. Nutr.* 2019; 59(6): 830-847.
<https://doi.org/10.1080/10408398.2018.1501658>
47. Holubkova A, Penesova A, Sturdík E, Mosovska S, Mikusova L. Phytochemicals with potential effects in metabolic syndrome prevention and therapy. *Acta Chimica Slovaca.* 2012; 5(2): 186-199. <https://doi.org/10.2478/v10188-012-0029-8>

48. Tang J, Li J, Qi W, Qiu W, Li P, Li B, *et al.* Inhibition of SREBP by a small molecule, Betulin, improves hyperlipidemia and insulin resistance and reduces atherosclerotic plaques. *Metabol.* 2011; 13:44–56. <https://doi.org/10.1016/j.cmet.2010.12.004>
49. Tofighi Z, Moradi-Afrapoli F, Ebrahimi SN, Goodarzi S, Hadjiakhoondi A, Neuburger M, *et al.* Securigenin glycosides as hypoglycemic principles of *Securigera securidaca* seeds. *J. Nat. Med.* 2016; 71(1): 272-280. <https://doi.org/10.1007/s11418-016-1060-7>
50. Eid HM, Martineau LC, Saleem A, Muhammad A, Vallerand D, Benhaddou-Andalousi A, *et al.* Stimulation of AMP-activated protein kinase and enhancement of basal glucose uptake in muscle cells by quercetin glycosides, active principles of the antidiabetic medicinal plant, *Vaccinium vitis-idea*. *Mol. Nutr. Food. Res.* 2010; 54(7): 991-1003. <https://doi.org/10.1002/mnfr.200900218>

UNDER PEER REVIEW