

Effect of Growth Regulators on Growth and Yield Attributes of Black Gram (*Vigna mungo* L.)

ABSTRACT

The field experiment was conducted during Zaid season of 2022 at the Crop Research Farm, Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P.) India. To study the Influence of Growth Regulators on the growth and yield of Black gram. The treatments consist of levels of Gibberellic Acid (GA₃ – 15 ppm, GA₃ – 30 ppm, GA₃ – 45 ppm) and Salicylic acid (SA – 50 ppm, SA – 75 ppm, SA – 100 ppm). The soil of experimental plot was sandy loamy in texture, nearly neutral in soil reaction (pH 7.8), low in organic carbon (0.35%). Results revealed that application of GA₃ 45 ppm + 50 ppm Salicylic acid (Treatment – 10). recorded with plant height (40.08 cm), plant dry weight (7.13 g/plant), number of pods/plant (36.27), number of seeds/pod (7.12), test weight (35.29 gm), 29.3 seed yield (957.86 kg/ha) were also recorded in treatment - 10 GA₃ 45 ppm + 100 ppm Salicylic acid at 15 & 40 DAS.

Keywords: *Black Gram*, Gibberellic Acid, *Growth Parameters*, Salicylic Acid, *Yield Attributes* and *Economics*.

INTRODUCTION

Leguminosae is the family that includes black gramme. The significant pulse crop grown throughout India's various agro-climatic zones is black gramme (*Vigna mungo* L.). Although this crop is farmed during the Kharif and Rabi seasons, the majority of its land is planted during the former, when intercropping with crops like sorghum, pearl millet, maize, cotton, castor, and pigeon pea is highly common. In terms of India's main pulse crops, black gramme holds a significant position. A complete array of nutrients, including 20 to 25% protein, 40 to 47% starch, ash lipids, carbs, and important vitamins, may be found in black gramme. In 2012–2013, the total area, production, and productivity of black gramme in India were 3.11 million hectares, 1.90 million tonnes, and 611 kg/ha, respectively (**Anonymous et al. 2014**).

A vast class of tetracyclic diterpenoid plant growth agents is known as the gibberellins. In order to control plant growth, GA₃ acts as a hormone, and this knowledge dates back to the 1950s (Solanki Mittal et al., 2018). Salicylic acid, also known as ortho-hydroxybenzoic acid, is a secondary metabolite that functions similarly to chemicals that control development. important in increasing the crop's yield (**Reham Dawar et al., 2020**).

Salicylic acid, or salicylate, is frequently referred to as "Aspirin" in acetylated form. Signalling is aided by salicylic acid. Salicylic acid is a member of an incredibly diverse class of plant phenols, which are substances with an aromatic ring bearing a hydroxyl group. In order to ascertain the impact of external application stress on the antioxidative enzymatic of salicylic acid (SA) under salt activities in black gramme, the green house experiment was carried out. Black gramme crop grain output has mostly increased as a result of the cumulative effect of yield-attributing traits, improved photosynthetic efficiency, and improvement in the reproductive and sinks' ability to utilise incoming assimilates as a result of the foliar application of GA₃.

MATERIALS AND METHODS

A field experiment was conducted during *zaid* 2022 at the Crop Research Farm, Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P.) India. The soil of experimental plot was sandy loamy in texture, nearly neutral in soil reaction (pH 7.2), low in organic carbon (0.72%), The treatments consist of Control, GA₃ 15 ppm + 50 ppm salicylic acid at 15 & 40 DAS, GA₃ 15 ppm + 75 ppm salicylic acid at 15 & 40 DAS, GA₃ 15 ppm + 100 ppm salicylic acid at 15 & 40 DAS, GA₃ 30 ppm + 50 ppm salicylic acid at 15 & 40 DAS, GA₃ 30 ppm + 75 ppm salicylic acid at 15 & 40 DAS, GA₃ 30 ppm + 100 ppm salicylic acid at 15 & 40 DAS, GA₃ 45 ppm + 50 ppm salicylic acid at 15 & 40 DAS, GA₃ 45 ppm + 75 ppm salicylic acid at 15 & 40 DAS, GA₃ 45 ppm + 100 ppm salicylic acid at 15 & 40 DAS. Ten treatments were replicated three times, and the experiment was set up using a randomised block design. By using the analysis of variance approach, statistical analysis of the acquired data was performed (**Gomez 1976**).

RESULT AND DISCUSSION

PRE AND POST-HARVEST PARAMETERS

Plant height - At 60 DAS, the significantly higher plant height of (40.08 cm) [Table.1] was recorded with treatment-10 (GA₃ 45 ppm + 100 ppm salicylic acid at 15 & 40 DAS). However, treatment-9 GA₃ 45 ppm + 75 ppm salicylic acid at 15 & 40 DAS (39.34cm) was found to be statistically at par with treatment-9 (GA₃ 45 ppm + 100 ppm salicylic acid at 15 & 40 DAS). The plant height of Black Gram increased significantly due to with At Harvest, Foliar application of GA₃ significantly influenced the black gram plant height (cm), number of pods/cluster, length of pod (cm), number of seedspod, number of pods/plant and grain yield/ plant (g). Two applications of 30 ppm GA₃ at flower and pod initiation stages (T10) recorded significantly higher number of pods/cluster, length of pod (cm), and grain yield/ plant. Similar results were reported by **Rehem Dawar *et al.* (2020)** and **Kodhati Vishnu *et al.* (2020)**.

Number of pods/plant – The significant and higher number of Pods/plant (36.27) were observed in treatment-10 with (GA₃ 45 ppm + 100 ppm Salicylic acid), which was significantly superior over rest of the treatments. However, treatment-8 (GA₃ 45 ppm + 50 ppm Salicylic acid), was found to be statistically at par with treatment-9(GA₃ 45 ppm + 75 ppm Salicylic acid). According to Singh and Totawat (2002), an improvement in the soil's nutrient status and the creation of a conducive environment for better root growth through the secretion of growth-promoting substances like Gibberellin, cytokinin, and auxin are responsible for the significant increase in the number of pods per plant following seed inoculation with growth regulators.

Number of seeds/pod - The significant and higher number of Seeds/pod (7.12) were observed in treatment-10 with (GA₃ 45 ppm + 100 ppm Salicylic acid), which was significantly superior over rest of the treatments. However, treatment-8 (GA₃ 45 ppm + 50 ppm Salicylic acid), was found to be statistically at par with treatment-9 (GA₃ 45 ppm + 100 ppm Salicylic acid). In this field experiment, the combined application of growth regulators hardly boosted the amount of seeds per pod. As seeds serve as a direct indicator of pollen viability, the presence of magnesium in multi-nutrient solutions may be to blame for the

increase in seeds per pod. According to Solanki mittal et al. (2018), magnesium has been shown to increase fruit set, pollen viability, and has a significant impact on pollen formation.

Seed yield - The significant and higher Seeds yield (957.86 kg/ha) were observed in treatment-10 with (GA₃ 45 ppm + 100 ppm Salicylic acid), which was significantly superior over rest of the treatments. However, treatment-8 (GA₃ 45 ppm + 100 ppm Salicylic acid), was found to be statistically at par with treatment-9 (GA₃ 45 ppm + 100 ppm Salicylic acid). Maximum seed yield (q) per hectare was recorded in T4 (SA 150 ppm). While, minimum seed yield (q) per hectare was recorded with treatment T7 (GA₃ 150 ppm) over the control. In this area, the combined use of growth regulators has only slightly boosted the amount of seeds per pod. The fact that the action of salicylic acid in mung bean reported greater seed output provided significant support for these findings. reported that salicylic acid's impact on chickpeas boosted seed production. reported that salicylic acid treatment improved the physiological efficiency of the crop and resulted in greater growth and yield of black gramme cv. All yield and yield components were increased in pea plant. NDU-1. The results of this study suggest that PGRs may be used to increase bean productivity by optimising yield-related parameters. (2017) Manjari et al.

CONCLUSION - It was concluded that with the application of GA₃ 45 ppm along with 50 ppm Salicylic acid at 15 & 40 DAS [Treatment-10, has performs positively and improves growth and yield parameters. Maximum seed yield, gross returns, net return and benefit cost ratio were also recorded with the application of GA₃ 45 ppm along with 100 Salicylic acid at 15 & 40 DAS [Treatment-10]. These results are based on a single season, therefore additional testing may be necessary for further conformation.

REFERENCES

- Aninda Chakraborty and Sanjoy Kumar Bordolui. (2021). Impact of Seed Priming with Ag Nanoparticle and GA₃ on Germination and Vigour in Green gram. *International Journal Of Current Microbiology and Applied Sciences*. **10**(03): 941-950.
- Bibek Laishram¹, T. Basanta Singh, Athokpam Kalpana, Merinda Wangkheirakpam, Sunil Kumar Chongtham and W. Jiten Singh. (2020). Effect of Salicylic Acid and Potassium Nitrate on Growth and Yield of Lentil (*Lens culinaris* L.) under Rainfed Condition. *International Journal of Current Microbiology and Applied Sciences*. **9**(11): 2779- 2791.
- Ergin, N. – Kyan, N. (2021). The effects of different gibberellic acid doses and application times on chickpea plants. *Journal of Bioscience and Agriculture Research*. **12**(05): 1812- 1814.
- Gomez, K.A., Gomez, A. A., (1976) Three or more factor experiment. (In:) *Statistical Procedure for Agricultural Research 2nd ed.*, 1976, pp.139 -141.
- Fatima Anaya, Rachid Fghire, Said Wahbi, Kenza Loutfi. (2015). Influence of salicylic acid on seed germination of *Vicia faba* L. under salt stress. *Journal of the Saudi Society of Agricultural Sciences*. **17**(1); 1-8.
- Ferdowsi noor, Feroza Hossain and Umme ara. (2017). Effects of gibberellic acid (GA₃) on growth and yield paramaters of French bean (*Phaseolus vulgaris* L.). *Journal of Asian. Soc. Bangladesh, Sciences*. **43**(1): 49-60.
- G.B. Vekaria, K.D. Rakholiya, V.D. Vora, J.T. Patel, G.S. Sutaria and P.D. Vekaria. (2017). Response of Sesame (*Sesamum indicum* L.) to Growth Regulator under Dry Farming Condition. *International Journal of Current Microbiology and Applied Sciences*. **6**(3): 1113-1120.
- Hafeez Ur Rahim, Sajjad Ahmad, Laiq Zada¹, Zaid Khan, Muhammad Ayoub Khan, Muhammad Haris, Amin Ullah, Usman. (2018). Yield and Growth Response of Maize Crop to Urea and Gibberellic Acid Potash Salt (Ga-K Salt) in Calcarious Soil. *JOJ Horticulture & Arboriculture*. **1**(2); 2641- 8215.
- Hasan Muhammad Zubair, Md. Mothaharul Islam, Md. Shahidur Rahman, Mohammad Mahbub Islam, Md. Shariful Islam and Mohammad Mustafizur Rahman. (2010). Role of GABA on Growth, Yield Contributing Characters and Yield of Sesame. *Plant Resource Management*. 184-188.

- Javed akhtar, rashid ahmad, M. yasin ashraf, Asif tanveer, ejaz ahmad waraich and hesham oraby. (2013). Influence of exogenous of salicylic acid on salt stressed mung bean (*vigna radiata* L.) growth and nitrogen metabolism. *Pakistan Journal of Botanical Sciences*. **45**(1): 119-125.
- Manjiri, Akanksha singh, Sarita devi gupta, Raj Bahadur and A.K.Singh. (2018). Response of Black gram (*Vigna mungo*) to foliar applied plant growth regulators. *International Journal of Current Microbiology and Applied Sciences* 2018; **7**:4058-4064.
- Manikandan Appu, Sathiyabama Muthukrishnan. (2014). Foliar Application of Salicylic Acid Stimulates Flowering and Induce Defense Related Proteins in Finger Millet Plants. *Universal Journal of Plant Science* **2**(1): 14-18.
- Neelambari1, Chetanaben Mandavia and S. Sree Ganesh. (2018). Curative Effect of Ascorbic Acid and Gibberellic Acid on Wheat (*Triticum astivum* L.) Metabolism under Salinity Stress. *International Journal of Current Microbiology and Applied Sciences*. **7**(1): 522-533.
- Nikita Nehal, Ah Khan, Nitish Sharma, Sanjay Kumar Tripathi and Mayanker Sing. (2018). Role of GA₃, SA and ABA on growth and yield of Indian mustard [*Brassica juncea* (L.) Czern. & Coss.] under rainfed condition. *Journal of Pharmacognosy and Phytochemistry*. **2**: 199-203.
- Pradip Kumar Saini, R.K. Yadav and Mayank Pratap. (2017). Effect of Foliar Application of GA₃, on Yield and Quality of Indian Mustard [*Brassica juncea* (L.) Czern. & Coss.] Under Sodic Soil. *International Journal of Current Microbiology and Applied Sciences*. **6**(12): 4156-4159.
- Reham Dawar, MD Giri, Ajit Kumar Meena, Gopal Patidar and Sandeep Rathod. (2020). Effect of foliar application of gibberellic acid on growth, yield and economic of black gram. *Journal of Pharmacognosy and Phytochemistry*. **9**(4): 3128 – 3190.
- Solanki Mital V, Trivedi Sandhya K, Kandoliya UK and Golakiya BA. (2018). Effect of exogenous application of salicylic acid on antioxidative enzymes in black gram (*Vigna mungo* (L.) Hepper) irrigated with saline water. *International Journal of Chemical Studies*. **6**(4): 2107- 2116.

Table 1. Influence of Phosphorus and Foliar Application of Zinc on Growth and Yield Attributes of Black Gram.

Treatments	Plant Height (cm)	Plant Dry Weight (g/plant)	Pods/Plant	Seeds/Pod	Test Weight (g/)	Seed Yield (kg/ha)
Control	35.05	5.43	26.28	3.72	29.87	627.50
T ₂	35.28	5.57	32.15	4.46	31.11	631.65
T ₃	36.28	5.73	33.19	4.91	32.18	670.77
T ₄	36.75	6.11	33.63	5.03	32.68	743.68
T ₅	37.12	6.13	34.06	5.27	32.97	754.60
T ₆	36.62	6.35	34.29	5.89	33.21	758.11
T ₇	37.83	6.38	34.54	6.19	33.54	784.53
T ₈	37.87	6.41	35.12	6.39	33.84	791.74
T ₉	39.34	6.53	35.73	6.83	34.47	932.33
T ₁₀	40.08	7.13	36.27	7.12	35.29	957.86
F test	S	S	S	S	S	S
S Em. (±)	0.49	0.23	1.06	0.21	0.38	12.04
CD (P=0.05)	1.47	0.69	3.15	0.62	0.96	35.78

UNDER PEER REVIEW