

Original Research Article
**SOIL PHYSICO-CHEMICAL PROPERTIES IN
DIFFERENT FORESTES OF TEHRI & PAURI-
GARHWAL HIMALAYS, UTTARAKHAND, INDIA.**

ABSTRACT

Garhwal Himalaya is the part of the Himalayan biodiversity hotspot and represents the western Himalayan landscape of the Indian Himalayas. The aim of present investigation was to analyse soil properties in different forest areas across different altitudes of Garhwal Himalayas i.e., *Quercus leucotrichophora* Forest (>1900m), *Anogeissus latifolia* Forest (1300m), Mixed Forest (900m) and Shrub Dominated Forest (600m). Physical properties and chemical properties of the soil were estimated using all standard procedures in department laboratory. According to findings of this study high-altitude temperate *Quercus leucotrichophora* Forest has dominated in all soil properties and greater potential to store soil organic carbon stock and Nitrogen ($29.64 \pm 3.48 \text{ t ha}^{-1}$) (Nitrogen $254.22 \pm 27.16 \text{ kg ha}^{-1}$) than Sub – tropical *Anogeissus latifolia* Dominated Forest, Mixed Forest, Shrub Dominated Forest with values of 18.34 ± 2.90 , 14.52 ± 2.27 and $10.74 \pm 1.57 \text{ t ha}^{-1}$.

Keywords: Garhwal Himalaya, *Quercus leucotrichophora*, *Anogeissus latifolia*, Forests, Carbon sink, Climate change, Soil Physico-Chemical properties.

1. INTRODUCTION

The Himalayas are the richest diverse and fragile forest ecosystem with tropical forests to alpine forests. Garhwal Himalayas are situated in the western part of Central Himalayas with wide altitudinal gradient range and rich biodiversity made it interesting for the research studies (Donald *et al.*, 1997), and have great potential to sequester carbon and mitigate climate change.

Vegetation plays an important role in soil formation. The vegetation enhances the aeration, hydraulic conductivity, water holding capacity, soil structure, and infiltration rate (Liorkar and Toley, 2001).

Forests play an important role in soil formation and also influence physico-chemical characteristics. The composition of the forest stand, ground cover, rate of tree growth, and other variables are influenced by the forest soil. Forest soils physiochemical properties vary through time and place because of changes in terrain, climate, physical weathering processes, plant cover, microbiological activity, and a number of other biotic and abiotic factors. Soil gets colour by processes called lithochromatic and pseudoachromatic processes. Soil colour is influenced by its mineral composition as well as organic matter, water content, and other factors. The soils with high calcium content are in white to grey colour and soils in high iron content are in red brown to black coloured.

Plant tissues (aboveground and belowground detritus) are the main source of soil organic matter (SOM), which influences physico-chemical characteristics. Soil organic carbon depends on organic matter availability in the soil, and it is an important factor for indicating soil quality, productivity. The physical and chemical characteristics of the soil are greatly influenced by the plants (Shameem *et al.*, 2011). Soil organic carbon (SOC) sequestration is the process of transfer and storage of atmospheric carbon into the soil through the decomposition. Estimates of SOC stocks are required to assess the role of soil in world carbon cycle, (Yang *et al.*, 2007). Soil pH is the concentration of H⁺ ions, used in measurement, the soil pH of 5.5 – 7.0 is the optimal suitable range for the most of the plant growth (Mc Clauety *et al.*, 2017). The water holding capacity (WHC) of soil determines how much amount of water retained by soil particles it is affected by soil structure and aggregate stability of soil. Bulk density is defined as the soil mass per unit volume, it indicates the compactness of the soil and its structure. It effects root growth and proliferation, aeration, soil water regime, biomass productivity (John *et al.*, 2001). The other soil physical properties

like moisture content, texture, soil porosity is influence the microbial activity, infiltration rate and gaseous exchange, nutrient uptake, overall plant growth, and Nutrient retention.

Soil nitrogen is available in soil as amonical, nitrate, nitrite forms is taken up and utilized by the plants, which influence plant growth, promotes productivity, species diversity, sustainability of forest ecosystems. It is estimated that more than 90% of the nitrogen reserves are in soil organic pool through N mineralization this nitrogen act as limiting factor for the productivity of forest ecosystems (Cheng *et al.*, 2016). The C:N ratio of soil refers to the proportion of carbon to nitrogen. In organic matter, carbon constantly predominates over nitrogen, the abbreviated C:N ratio is typically expressed as a single value (Flavel and Murphy, 2006). The increase in greenhouse gases emissions and climate change on earth all eyes shifting towards mitigation activities, while forest terrestrial ecosystems are the major source of soil carbon pool and nitrogen pools.

2. MATERIAL AND METHODS

2.1 Description of study area

This study was carried out in the Silkakal (30°14'59.9064" N 78° 46' 43.4676" E), Badiyalgarh (30° 14' 20.1552" N, 78° 48' 27.846" E) (Tehri Garhwal) and Adwani (30° 4' 20.4924" N, 78° 42' 58.626" E) (Pauri Garhwal) regions of Garhwal Himalaya forest areas in districts of Tehri and Pauri of Uttarakhand, India. (**fig 1**). The present investigation includes four sites of different forest lands areas i.e., *Anogeissus latifolia* Dominated Forest (ALDF: towards Silkakal), Mixed Forest Area (MF: towards Silkakal), Shrub Dominated Forest (SDF: towards Badiyalgarh) and *Quercus luecotrichophora* Forest (QLDF: Pauri, Adwani) across different altitudes ranges from **600 to 2000m asl**. The average annual rainfall is around 800 - 1500 mm and average temperature rise up to 35°C in summers and fall up to 15°C in winters.




Fig.1: View of Indian state of Uttarakhand (Left) and Study area (Right)

2.2 Data Collection

The 9 Soil samples were collected from 3 sites randomly from each forest area which has 3 depths i.e., 0- 20cm, 20-40cm and 40-60cm. Thus, A total of 36 samples were collected from all four forest areas. The physico-chemical properties of soil were analyzed by using all standard procedures (Table 1) in department laboratory and mean values reported among all samples (Table 3, Table 4 & Table 5).

Table 1: Methodology used for the analysis of different soil parameters.

S No	Soil Parameter	Formula/ Method	Reference
1.	Moisture Content (%)	$\frac{\text{Fresh weight of soil (g)} - \text{dry weight of soil (g)}}{\text{Dry weight of soil (g)}} \times 100$	Upreti, 2019
2.	Soil Texture and class	$\frac{\text{Weight of sieved soil proportion}}{\text{Total soil sample weight}} \times 100$ <p>Class: Based on texture percentage values and assessed by using texture triangle method.</p>	USDA, Groenendyk <i>et al.</i> , 2015
3.	Soil bulk density (g/cm^3)	$\frac{\text{Dry soil weight (g)}}{\text{Soil volume (cm}^3\text{)}}$ $\text{Soil volume (cm}^3\text{)} = 3.14 \times \text{radius}^2 \times \text{ring height (h)}$	ISO, 2017
4.	Water holding capacity (%)	$\frac{W2 - W3 - W4}{W3 - W1} \times 100$	Upreti, 2019
5.	Soil pH	Determined using dynamic digital pH meter	Jackson, 1958

6.	Soil organic carbon (%)	$\frac{10 (B-T)}{B} \times \frac{0.003 \times 100}{\text{weight of soil (g)}}$	Walkley & Black, 1934
7.	SOC Stock (t ha ⁻¹)	Soil bulk density x Soil depth x SOC (%)	Pearson, 2007
8.	Soil Nitrogen (kg/ha ⁻¹)	$\frac{14 \times T_v \times 0.02N \times 2.24 \times 10^6}{\text{Soil sample weight (gm)} \times 1000}$ Kg/ha x 0.4 = Kg/acre	Sáez-Plaza <i>et al.</i> , 2013
9	Soil colour	Munsell colour chart	Munsell, 2010
10	Soil organic matter (SOM)	SOM% = Soil organic carbon (%) x 1.724.	Budiman <i>et al.</i> , 2020.
11	C:N ratio	The proportion of carbon to nitrogen	Flavel and Murphy, 2006

3. RESULTS AND DISCUSSION

3.1 Soil Physical Properties

3.1.1 Moisture Content (MC%)

In present study, soil MC% ranges from 1.52 ± 0.62 to $7.48 \pm 2.15\%$. The highest MC was in *Q. leucotrichophora* dominated forest ($7.48 \pm 2.15\%$) because of good WHC of soil and higher proportion of clay particles as well as dense forest cover and followed by forest areas of MF, ALDF and SDF with the values of 5.22 ± 0.77 , 3.05 ± 0.42 and $1.52 \pm 0.62\%$ respectively. Mahato *et al.*, 2013 reported MC of 14.39% in forest lands of the Garhwal Himalaya. Chauhan *et al.*, 2020 also reported the MC ranged from 9.01 to 16.09% in sub-tropical forest belt of the Garhwal Himalayas. These values of MC are higher than the present study

3.1.2 Water holding capacity (WHC%)

The WHC of present study ranged from 15.29 to 63.68% in different forest areas. Amongst the study sites, the highest WHC (48.63%) was found in QLDF due to good amount of clay particles and hold maximum water content in the soil which followed by ALDF, SDF and MF with the reported values of 33.71, 31.37 and 29.52% respectively, all forests types have shown moderate level of WHC. Chauhan *et al.*, 2020 also reported the values of WHC ranged from 24.84 to 55.23% in the similar *Anogeissus latifolia* forests of Garhwal

Himalayas, the present study estimates also falls within the same range. Mahato *et al.*, 2016, reported WHC value of 24.17% for the forest of Garhwal Himalaya and this value is lower than the present study.

3.1.3 Bulk Density (BD g cm⁻³)

Bulk density of the present study ranges from 0.73 ± 0.07 to 0.84 ± 0.24 (Among over all individual sample values range 0.41 to 1.06 g cm³) in different forest areas. The highest BD was found in *Q. leucotrichophora* dominated forest 0.84 g cm³ followed by SDF, ALDF and MF with the values of 0.79, 0.73 and 0.70 g cm³. Chauhan *et al.*, (2020) reported BD values from 0.90 to 1.06 g cm³. Bhat *et al.*, (2012) in temperate Himalayan forests reported BD ranged from 1.41 to 1.59 g cm⁻³. Bam and Surendra (2013) study BD of forest lands (1.0 g cm⁻³) and shrub lands (1.4 g cm⁻³). Mahato *et al.*, (2016) in the study of community managed forest reported BD of 1.29 g cm⁻³. All other works reported BD values were higher than the present study. It might be because of higher proportion of gravel particles which contributed more than 50% of the tested samples, where compactness of soil reduces due to presence of gravel.

3.1.4 Soil texture (%)

The particle of sand ranges from 51.67 (qldf) to 66.67% (mf). silt particle also ranges from 7.35 (sf) to 9.99% (aldf) and clay particle ranges from 23.58 (mf) to 39.07% (qldf). the highest proportion of soil particles was of sand followed by silt and clay in all the depths. the texture class was sandy clay loam and sandy clay. chauhan *et al.*, (2020) reported that the highest proportion reported of sand particle followed by silt and clay and texture class was sandy clay. mahato *et al.*, (2016) stated the highest proportion of sand particle followed by silt and clay. munesh *et al.* (2012) study on *a. latifoila* stands reported similar proportion of soil particles.

3.1.5 Texture class

A. latifolia dominated forest and mixed forest having texture classes of Sandy Clay Loam and Q. leucotrichophora dominated forest and shrub forest have Sandy Clay. The variation in texture class was not reported with the soil depths.

3.1.6 Soil Colour

Amongst the forest areas, various soil colour has been reported such as Pale Brown colour (A. latifolia dominated forest), Light Grey Colour (Mixed Forest), Yellowish Brown, Reddish Yellow & Pale Brown colour (Shrub Forest) and Brown to Yellowish Brown coloured (Q. leucotrichophora dominated forest) reported. Soil colour also used to predict the presence mineral properties of soil (**Table 2**).

Table.2 Study area forest soil colour and its mineral interpretation

Forest area	Depth (cm)	Soil colour	Mineral pigment (source USDA)
ALDF	0-20	pale brown	gypsum
	20-40	greyish brown	-
	40-60	brown	geiothite, brown
MF	0-20	light grey	quartz
	20-40	greyish brown	-
	40-60	light grey	quartz
SDF	0-20	Reddish yellow	hematite, lepidocrocite
	20-40	light yellowish brown	gypusm
	40-60	pale brown	gypsum
QLDF	0-20	brown	geothite, humus
	20-40	light yellowish brown	gypsum

3.2 Soil Physical Properties

3.2.1 Soil pH

Soil pH value ranges from 5.25 to 7.67 in the present study area. The highest (6.91) pH was found in ALDF followed by QLDF (6.69), MF (6.26) and SDF (6.02). The study revealed that the pH of all sites was Slightly Acidic. In depth wise, pH shown increasing trend with increasing depth. Chauhan et al. (2020) reported pH values ranged from 6.44 – 6.79. Mutanol *et al.*, (2016) estimated pH value of A. latifolia stands as 6.5. These values are similar to the present study. Mahato *et al.*, (2016) study on Community Forest of Garhwal

Himalaya reported pH value of 5.9. Rawat *et al.*, (2022) reported pH of 5.6. Shukla *et al.*, (2021) reported pH values from 4.86-5.16 for Mixed Forest stands & Single dominant forest (Sal, Teak), these values are quite lower of the present study.

3.2.1.1 Effect of soil pH on nutrient availability

Jackson *et al.*, (2018) suggested the function of pH and the availability of vital plant nutrients (**Table 3**) are closely connected. The plants' ability to get these nutrients is correlated with varied pH conditions. With the exception of phosphorus, the macronutrients such as nitrogen, calcium, potassium, magnesium, and Sulphur are more easily obtainable in the pH range of 6.5-8. However, the micronutrients are available at a pH range of 5-7, which is slightly acidic. These are the ranges where nutrients are most readily supplied to plants in a sufficient amount. In present study, soil pH ranges from 5.25 to 7.67 i.e., slightly acidic soils which states that better availability of nutrients in soil for good plant growth in all present forest areas.

3.2.2 Soil Organic Matter (SOM%)

It was analyzed that SOM ranges from 4.79 to 0.27%. On mean the highest SOM was found in *Q. leucotrichophora* dominated forest was 2.99% followed by *A. latifolia* dominated forest, mixed forest and shrub dominated forest with the values of 2.14, 1.70 and 1.15% respectively.

3.2.3 Soil organic carbon (SOC %)

SOC % ranged from 0.67 ± 0.32 to $1.73 \pm 0.16\%$ (Among over all individual samples the values ranged from 0.16 to 4.08%) with the highest SOC found in QLDF (1.73 ± 0.16) followed by ALDF ($1.24 \pm 0.16\%$), MF ($0.99 \pm 0.08\%$) and SDF ($0.67 \pm 0.32\%$). Chauhan *et al.*, (2020) reported SOC from 1.09–1.36%. Mahato *et al.*, (2016) study on Community Forest of Garhwal Himalaya reported SOC % of 1.28. Gupta and Rout (1989) study on Mixed-forest (*A. latifolia* & *Lannea coromandelica*) reported SOC of 2.99%..

3.2.4 SOC Stock (t ha-1)

SOC stock ranges from 10.74 ± 1.57 to 29.64 ± 3.48 t ha⁻¹ (Among over all individual sample values ranged from 2.81 to 46.70 t ha⁻¹). The highest SOC stock was found in high altitude QLDF (29.64 ± 3.48 t ha⁻¹) followed by ALDF, MF and SDF with values of 18.34 ± 2.90 , 14.52 ± 2.27 and 10.74 ± 1.57 t ha⁻¹. Munesh *et al.*, (2012) studied in *A. latifolia* forests and reported SOC stock of species *Rhus parviflora* (168.00 t ha⁻¹) and *Lantana camara* (164.16 t ha⁻¹). These values were higher compared to *A. latifolia* forest (161.28 t ha⁻¹). Shahid and Joshi *et al.*, (2018) studied that Carbon Stock Variation in Different Forest Types of Western Himalaya where carbon stock density varied between 129.81 and 136.00 Mg C ha⁻¹. Mahato *et al.*, (2016) reported SOC stock 218.57 t ha⁻¹ in Community Forest of Garhwal Himalaya. Sheikh *et al.*, (2011) reported 60.8-185.6 t ha⁻¹ in *Quercus* of Garhwal Himalaya. The SOC stock values of all the studies were higher than the present study. Singh *et al.*, (2011) reported SOC stock of 64-72 t ha⁻¹ in oak forest of Garhwal Himalaya. Shukla *et al.*, (2021) for mixed forest stands & Single dominant forest (Sal, Teak) estimated SOC stock of 75.9 -107.7 t ha⁻¹. Chauhan *et al.* (2020) study on *A. latifolia* forest stands of Garhwal Himalayas reported stock of 14.94 -23.78 t ha⁻¹ were similar to the present study.

3.2.5 Soil Nitrogen (kg ha⁻¹)

Nitrogen ranges from 193.79 ± 8.15 to 254.22 ± 27.16 (Among over all individual samples values ranged from 170.61 to 306.07 kg ha⁻¹). The highest (254.22 ± 27.16 kg ha⁻¹) nitrogen was found in QLDF followed by ALDF (205.47 ± 7.10 kg ha⁻¹), MF (197.07 ± 1.50 kg ha⁻¹) and SDF (193.79 ± 8.15 kg ha⁻¹). Chauhan *et al.*, (2020) study on *A. latifolia* forest reported soil nitrogen values from 291.64 - 323.01 kg ha⁻¹. Rawat *et al.*, (2022) reported nitrogen of 217.38 kg ha⁻¹ in Oak Forest. Mutanal *et al.*, (2016) study on *A. latifolia* forest reported nitrogen of 235.01 kg ha⁻¹. All this studies, reported similar and close values to the present study. Shukla *et al.* 2021 study on Mixed Forest stands & Single dominant forest (Sal, Teak) estimated stored soil nitrogen values ranged from 210 -260 kg ha⁻¹.

3.2.6 C:N ratio

The C:N ratio ranged from 5.54 to 11.65. The highest mean C:N ratio was found in Q. leucotrichophora dominated forest (11.65 ± 1.01) followed by A. latifolia dominated forest (8.93 ± 1.29), mixed forest (7.37 ± 0.48) and lowest shrub dominated forest (5.54 ± 0.83) respectively. Chauhan *et al.* (2020) study reported C/N ratio of < 10 within range of present study. Donald *et al.* (2011) estimated C/N ratio range of 11.6 – 45.3 in different tree species, the both N concentration and the C/N ratio was strongly related to tree species and C content. The Low C/N ratio was found in present study could be because of high soil nitrogen compared SOC. When the C:N ratio is between 1 and 15, N is quickly mineralized and released, making it available for plant uptake. The faster nitrogen is released into the soil of usage of crops with the lower the C:N ratios (Watson *et al.* 2002).

The Depth wise distribution of Soil Physico-chemical properties in the study area is demonstrated in the **Table.5**.

Table. 3 Mean values of soil physico- chemical properties

Physico-chemical properties of soil in different forest areas							
Parameter	Forest area				Depth wise		
	ALDF (1300m)	MFA (900m)	SDF (600m)	QLDF (1900m)	0-20	20-40	40-60
Moisture (%)	3.05 ± 0.42	5.22 ± 0.77	1.52 ± 0.62	7.48 ± 2.15	4.36±1.14	4.9±1.51	3.67±0.78
WHC (%)	33.71 ± 4.20	29.52 ± 3.72	31.37 ± 6.58	48.63 ± 14.15	43.52 ± 3.60	33.51±4.83	30.27±4.46
Soil pH	6.91 ± 0.17	6.26 ± 0.07	6.02 ± 2.06	6.69 ± 2.29	6.23±0.16	6.48±0.18	6.73±0.21
SOC (%)	1.25 ± 0.16	0.99 ± 0.08	0.67 ± 0.32	1.73 ± 0.16	1.45±0.19	1.23±0.11	0.81±0.6
BD (g cm ³)	0.73 ± 0.07	0.704 ± 0.06	0.79 ± 0.18	0.84 ± 0.24	0.82±0.03	0.84±0.04	0.63±0.04
SOC Stock (t ha ⁻¹)	18.34 ± 2.90	14.52 ± 2.27	10.74 ± 1.57	29.64 ± 3.48	28.73±5.20	49.29 ± 4.79	59.14 ± 1.94
Nitrogen (kg ha ⁻¹)	205.47 ± 7.10	197.07 ± 1.50	193.79 ± 8.15	254.22 ± 27.16	221.76 ±14	220.75±25.40	199.26 ± 5.83
SOM (%)	2.14 ± 0.28	1.70 ± 0.14	1.15 ± 0.55	2.99 ± 0.42	2.31±0.30	1.85±0.23	1.31±0.21

C/N Ratio	8.93 ± 1.29	7.37 ± 0.48	5.54 ± 0.83	11.65 ± 1.01	12.95 ± 2.38	22.32 ± 2.60	29.67 ± 2.92
------------------	----------------	----------------	----------------	-----------------	-----------------	-----------------	-----------------

UNDER PEER REVIEW

Table.4 Mean values of soil texture and texture classes

Particle	Forest areas			
	ALDF	MF	SDF	QLDF
Sand (%)	61.21 ± 1.27	66.67 ± 1.29	57.63 ± 2.37	51.67 ± 1.22
Silt (%)	9.99 ± 0.49	9.27 ± 1.00	7.35 ± 0.43	9.27 ± 1.03
Clay (%)	28.91 ± 1.15	23.58 ± 1.46	34.05 ± 2.52	39.07 ± 0.80
Texture class	Sandy Clay Loam	Sandy Clay Loam	Sandy Clay	Sandy Clay
Gravel (%)	67.19 ± 0.99 (Sand, silt, clay=32.81%)	69.46 ± 2.40 (Sand, silt, clay=30.54%)	66.92 ± 3.36 (Sand, silt, clay=33.08%)	52.13 ± 5.84 (Sand, silt, clay=47.87%)

Table.5 Depth wise distribution of soil Physico-chemical properties in the study area.

Forest area	Depth (cm)	SOC (%)	SOC stock (t ha ⁻¹)	Nitrogen (kg ha ⁻¹)	SOM (%)	BD (g cm ⁻³)	Soil pH (1:2.5)	WHC (%)	Moisture (%)
ALDF (1300 m)	0-20	3.06	26.67	225.79	3.06	0.74	6.68	45.87	3.29
	20-40	2.17	19.81	210.73	2.17	0.84	6.92	31.04	3.09
	40-60	1.22	8.53	195.63	1.22	0.61	7.31	24.23	2.77
MF (900m)	0-20	2.16	39.56	199.66	2.16	0.88	6.08	43.6	6.11
	20-40	1.62	13.45	190.66	1.62	0.71	6.32	21.23	5.67
	40-60	1.34	8.05	200.7	1.34	0.52	6.38	23.23	3.88
SDF (650m)	0-20	1.41	12.68	195.68	1.41	0.75	5.8	32.32	1.14
	20-40	1.22	12.55	175.61	1.22	0.90	5.97	33.57	1.47
	40-60	0.84	6.37	210.07	0.84	0.75	6.3	28.22	1.94
QLDF (1900m)	0-20	2.62	36.44	265.92	2.62	0.94	6.37	52.29	6.88
	20-40	2.42	36.04	306.07	2.42	0.93	6.74	48.21	9.46
	40-60	2.03	16.44	190.66	2.03	0.65	6.93	45.38	6.12

UNDER PEER REVIEW

Table.6 Comparative studies of earlier works

Vegetation type	Moisture (%)	WHC (%)	BD(g cm ³)	Soil pH	SOC (%)	SOC Stock (t ha ⁻¹)	SOM (%)	Nitrogen (kg ha ⁻¹)	References
<i>A. latifolia</i> dominated forest (ALDF)	3.05	33.71	0.73	6.91	1.24	18.34	2.14	205.47	Present study
Mixed Forest (MF)	5.22	29.52	0.70	6.26	0.99	14.52	1.70	197.07	
Shrub dominated forest (SDF)	1.52	31.37	0.79	6.02	0.67	10.74	1.15	193.79	
<i>Q. leucotrichophora</i> Dominated forest (QLDF)	7.48	48.63	0.84	6.69	1.73	29.64	2.99	254.22	
<i>A. latifolia</i>	-	30.90	-	-	-	161.28	-	-	Kumar et al. 2012
Shrubs forest (<i>Lantana camara</i> & <i>Rhus parviflora</i>)	-	-	-	-	-	168.00	-	-	Kumar et al. 2012
Mixed forest (<i>A. latifolia</i> & <i>Lannea coromendalica</i>)	-	49.80	1.31	7.60	2.99	-	-	-	Gupta and Rout (1989)
<i>A. latifolia</i> forest	-	-	1.30	6.5	-	-	-	235.01	Mutanal et al. (2016)
Community Forest of Garhwal Himalaya	14.39	24.17	1.29	5.90	1.28	218.57	-	-	Mahato et al. (2016)
<i>Quercus leucotrichophora</i> forest	-	-	-	-	-	64-72	-	-	Singh et al. (2011)
<i>Quercus leucotrichophora</i> forest	-	-	-	-	-	60.8-185.6	-	-	Sheik et al. (2011)
Shrub land	-	-	-	6.38	2.06	-	-	-	Worku et al. (2014)
<i>Quercus. L</i>	-	-	-	5.6	1.95	-	-	217.38	Rawat et al.2022
Mixed forest stands & Single dominant forest (Sal, Teak)	27.01-1.03	-	-	4.86-5.16	-	75.9-107.7	-	0.21-0.26	Shukla et al. 2021

<i>A. latifolia</i> Forest	9.01-16.09	24.84-55.23	0.90-1.06	6.35-6.79	1.36	14.94 - 23.78	-	291.64 - 323.01	Chauhan et al. 2022
Forest land	-	-	1.00	-	-	98.01	-	-	Bam & Surendra et al. 2013
Barren land	-	-	1.2	-	-	83.6	-	-	
Agriculture land	-	-	1.4	-	-	36.6	-	-	
Shrub land	-	-	-	-	-	10.8%	-	55.9%	Wang & Kang et al. 2013
Mixed forest (<i>Quercus</i> & <i>Rhododendron</i>)	25.83	-	1.47	-	-	23.2 mg/ha	-	-	Bhat et al. 2012
Grass land	33.21	-	1.53	-	-	25.0	-	-	
Tropical forest	-	-	-	-	-	64.3	-	-	Gachhadar et al. 2022
Sub-tropical forest	--	-	-	-	-	84.4	-	-	
Mountain forest	-	-	-	-	-	95.3	-	-	
Chauras Campus (Garhwal University)	3.30 ±0.28	31.53±0.72%	1.38±0.4 g/cm ³	6.84±0.11	0.86±0.06%	302.62 t ha ⁻¹		133.12±5.79	Prashanth <i>et al.</i> , In Press

4. CONCLUSION

This study revealed that soil physico-chemical properties changes with different forest areas, altitude and also with soil depth. The Highest SOC and Nitrogen reported in high altitude QLDF forest area, followed by ALDF, MF, SDF. Same trend followed in almost all soil properties like MC, WHC, SOC%, SOM, Nitrogen, SOC stock), highest pH found in ALDF. Highest Bulk density was found in SDF followed by QLDF, ALDF, MF.

C/N ratio, Soil pH, Nitrogen and SOC stock was increased with increase in depth, While MC, WHC, SOM, SOC (%), BD, were found decreasing with soil depth.

This study concluded that among four different forest areas, the high-altitude temperate *Quercus leucotrichophora* Forest (QLDF) has greater potential to store soil organic carbon stock and Nitrogen than Sub – tropical *Anogeissus latifolia* Dominated Forest (ALDF), Mixed Forest (MF), Shrub Dominated Forest (SDF). In almost all other soil properties are dominated by *Quercus leucotrichophora* Forest followed by *Anogeissus latifolia* Dominated Forest, Mixed Forest, Shrub Dominated Forest. This study revealed the carbon sequestering potential of four major Himalayan region forest types and their key role to combat the climate change.

CONSENT

I on behalf of all authors confirms that all authors are read the copy of this manuscript and all authors are approved for submission of this manuscript.

REFERENCES

- Bam, Surendra Bhandari, Suresh (2013). Comparatives Study of Soil Organic Carbon (SOC) under Forest, Cultivated and Barren Land: A Case of Chovar Village, Kathmandu. *Nepal Journal of Science and Technology*. 14. 103-108. 10.3126/njst.v14i2.10422.
- Bhat Jahangeer, Kumar Munesh, Negi Ajeet, Pala Nazir, Todaria Np (2012). Soil Organic Carbon Stock and Sink Potential in High Mountain Temperate Himalayan Forests Of India. *International journal of current research*. vol.4, pp 206 – 209.
- Budiman Minasny, Alex B, McBratney, Alexandre M.J.-C. Wadoux Erwin Nyak Akoeb, Tengku Sabrina, Precocious 19th century soil carbon science, *Geoderma Regional*, Volume 22, 2020, e00306, ISSN 2352-0094, <https://doi.org/10.1016/j.geodrs.2020.e00306>.
- Chauhan Monika, Kumar Munesh Dr (2020). Impact of Carbon Stocks of *Anogeissus latifolia* on Climate Change and Socioeconomic Development: a Case Study of Garhwal Himalaya, India. *Water Air and Soil Pollution*. pp 231. 10.1007/s11270-020-04803-8.
- Donald B, Zobel, Surendra P Singh (1997). Himalayan Forests and Ecological Generalizations *BioScience*, Oxford University Press. Vol. 47, No. 11 (Dec., 1997), pp. 735-745 (11 pages).
- Donald S, Ross, Scott W Bailey, Gregory B Lawrence, James B Shanley, Guinevere Fredriksen, Austin E Jamison, Patricia A, Brousseau (2011). Near-Surface Soil Carbon, Carbon/Nitrogen Ratio, and Tree Species Are Tightly Linked across Northeastern United States Watersheds. *Forest Science*. pp 57(6)
- Flavel TC, Murphy DV (2006). Carbon and nitrogen mineralization rates after application of organic amendments to soil. *J Environ Qual*, vol 35: pp183–193.

- Groenendyk DG, Ferre TP, Thorp KR, Rice AK (2015). Hydrologic-process-based soil texture classifications for improved visualization of landscape function. *PloS one*, 10(6), e0131299.
- Gupta S.R, Rout K.S (1989). Analysis of Forest Vegetation of Morni Hills in Northeast Haryana. *Plant Science*, 99(2): 117-126.
- Ilorkar VM, Totey NG (2001). Floristic diversity and soil studies in Navegaon National Park (Maharashtra). *Indian J For* 24: pp 442-447.
- International Organization for Standardization (ISO). (2017). ISO 11272-2017. Soil Quality Determination of Dry Bulk Density. International Organization for Standardization, Geneva.
- Jackson Khaidem, Thomas Thounaojam, Meetei, Thounaojam Thomas (2018). Influence of soil pH on nutrient availability: A Review. *Journal On Emerging Technology and Innovative Research*, 707; 44
- Jackson M.L (1958). Soil Chemical Analysis. *Prentice Hall Inc.*, New Jersey, USA.
- John M Kimble, Rattan Lal, Richard Birdsey, Linda S Health, Characterization of soil organic carbon pools, in methods for soil organic carbon. Eds., CRC Press, Boca Raton, FL, 2001.
- Mahato S, Dasgupta S, Todaria N.P (2016). Tree and soil carbon stock by community managed forests in Garhwal Himalaya. *Plant Archives*. Vol.16 (2), pp 805-811.
- McCauley K.B, Hawkins F, Serra M, Thomas D.C, Jacob A, Kotton D.N 2017. Efficient Derivation of Functional Human Airway Epithelium from Pluripotent Stem Cells via Temporal Regulation of Wnt Signaling. *Cell Stem Cell*. 20(6):844-857.
- Munesh Kumar*, Manish Kumar, S. S. Khan, M. Joshi, S. Kumar, J. A. Bhat (2012). Soil organic carbon stock potential of shrubs, occupying natural *Anogeissus latifolia* forest in sub-tropical belt of Garhwal Himalaya. *Journal of Ecology and the Natural Environment*. Vol. 4(8), pp. 219-222.

- Munsell Color (Firm). (2010). Munsell soil color charts: with genuine Munsell color chips. Grand Rapids, MI: Munsell Color.
- Mutanal S.M, Patil HY, Mokashi M.V (2016). Economic evaluation multipurpose tree species degraded lands Karnataka, *International Journal Forestry and Crop Improvement* 7(1):35-40 of Morni hills in northeast Haryana. *Plant Science* 99(2): 117-126.
- Pearson TR (2007). Measurement guidelines for the sequestration of forest carbon (Vol. 18). US Department of Agriculture, Forest Service, Northern Research Station.
- Prashanth Vempally, Murari Chiluveri, Munesh Kumar, Shakith MA, Sushanth Thalari, Manoj Chandra. *Environment and Ecology*, 2023 EE41: (In Press).
- Sáez-Plaza P, Navas MJ, Wybraniec S, Michałowski T, Asuero AG (2013). An overview of the Kjeldahl method of nitrogen determination. Part II. Sample preparation, working scale, instrumental finish, and quality control. *Critical Reviews in Analytical Chemistry*, 43(4), 224-272
- Shameem SA, Kangroo NI, Bhat GA (2011). Comparative assessment of edaphic features and herbaceous diversity in lower Dachigam national park, Kashmir, Himalaya. *J Ecol Nat Environ* 3: 196-204
- Sheikh Mehraj, Kumar Munesh, Bussmann Rainer (2009). Altitudinal variation in soil organic carbon stock in coniferous subtropical and broadleaf temperate forests in Garhwal Himalaya. *Carbon balance and management*. 4. 6. 10.1186/1750-0680-4-6.
- Singh J S. (2005). Sustainable development of the Indian Himalayan region: Linking ecological and economic concerns. *Current science*. 90.
- United states Department of Agriculture (USDA), NRCS Colour of the soil. www.nrcs.usda.com
- Upreti Brij (2019) Analysis of soil physical properties of different land forms in and around Nagal Hatnala region, Dehradun. 3. 34 - 38.

Walkley and I.A. Black. 1934. An examination of Degtjareff method for determining soil organic matter and a proposed modification of the Chromic Acid titration method. *Soil Science*. 37: 29-37

Yang Y, Mohammat A, Feng J, Zhou R, Fang J (2007). Storage, patterns and environmental controls of soil organic carbon in China. *Biogeochemistry*, 84: 131–141.

DEFINITIONS, ACRONYMS, ABBREVIATIONS

Here is the Definitions section. This is an optional section.

Term: Definition for the term

UNDER PEER REVIEW