

Response of Vermicompost and Neem cake on Soil Health and Yield attributes of Cluster bean (*Cyamopsis tetragonoloba* L.) var. Pusa Naubahar

Abstract

The present investigation carried out at research farm of department of Soil Science and Agricultural Chemistry, Naini Agricultural Institute, SHUATS, Prayagraj, U.P. during the *zaid* season of 2021-22 with the objective to response of different levels of vermicompost and neem cake on soil health, growth and yield attributes of Cluster bean [*Cyamopsis tetragonoloba* L.] Var. Pusa Naubahar. The experiment was laid out in a randomized block design with nine treatment combinations, consisting of three vermicompost levels (0, 50 and 100%) and neem cake (0, 50 and 100%). In soil parameters bulk density (Mg m^{-3}) of soil was recorded 1.18 Mg m^{-3} in treatment T₉ (100% NPK + 100% VC + 100% NC). Similar results were also reported in the particle density (Mg m^{-3}) of soil was recorded 2.14 Mg m^{-3} in treatment T₉ (100% NPK + 100% VC + 100% NC). Soil pore space was recorded 48.22% in treatment T₉ (100% NPK + 100% VC + 100% NC). It was observed that Soil pH after harvesting 7.56 which was recorded in T₉ (100% NPK + 100% VC + 100% NC). Electrical conductivity (dS m^{-1}) after harvesting was 0.270 recorded with T₉ (100% NPK + 100% VC + 100% NC). Organic carbon (%) of soil after harvesting was 54.5 in T₉ (100% NPK + 100% VC + 100% NC). Available nitrogen in soil was 276.49 kg ha^{-1} after harvesting in T₉ (100% NPK + 100% VC + 100% NC). Available phosphorus in soil was 30.74 kg ha^{-1} after harvesting and highest was in T₉ (100% NPK + 100% VC + 100% NC). Available potassium in soil was 262.65 kg ha^{-1} after harvesting and highest was in T₉ (100% NPK + 100% VC + 100% NC). It was observed that for post-harvest treatment T₉ (100% NPK + 100% VC + 100% NC) was best in terms of growth, yield and economic parameters with maximum plant height 82.25 cm, number of leaves plant^{-1} 26.19, number of branch plant^{-1} 6.90, pods plant^{-1} 79.50, pod yield 56.13 and maximum cost benefit ratio (C: B) of (1:2.52).

Keywords: Soil properties, Inorganic fertilizers, Vermicompost, Neem cake, Cluster bean.

Comment [Ma1]: This abstract should include method of analysis, summary of key results/discussion and conclusion.

Comment [Ma2]: Something is missing here, there is no connection. Separate objective from the location of the farm.

Introduction

Soil is a medium for plant growth. Crop production is based largely on soils. Some of the soil properties affecting plant growth include: soil texture (coarse fine), aggregate size, porosity, aeration (permeability), and water holding capacity, pH, bulk density, particle density. The rate of water movement into the soil (infiltration) is influenced by its texture, physical condition (soil structure and tilth), and the amount of vegetative cover on the soil surface. Organic matter tends to increase the ability of all soils to retain water, and also increases infiltration rates of fine textured soils. Bulk density reflects the soil's ability to function for structural support, water and solute movement, and soil aeration. Soil pH directly affects the solubility of many of the nutrients in the soil needed for proper plant growth and development (**Ghaffari et al., 2011**). Soil pH measurement is useful because it is a predictor of various chemical activities within the soil. As such, it is also a useful tool in making management decisions concerning the type of plants suitable for location, the possible need to modify soil pH (either up or down), and a rough indicator of the plant availability of nutrients in the soil. Three elements, carbon, oxygen and hydrogen, are essential to plant growth and are supplied by air and water. The other essential elements are referred to as plant nutrients, and are provided by the soil, or are added as fertilizers, and enter plants almost exclusively through the roots (**Singh et al., 2008**). Cluster bean popularly known as 'guar' is an important self-pollinated, multipurpose, relatively drought resistant and restorative leguminous vegetable crop mainly grown under rainfed condition in arid and semi-arid regions of India during *kharif* season. It is produced for the manufacturing of gum as well as for use as feed, fodder, vegetables, and green manure. Due to the fact that legumes improve soil fertility, they play a significant part in the nitrogen cycle for succeeding crops. It is very hardy and drought tolerant crop. **Chaudhari, J.A. (2018)** Its deep penetrating roots enable the plant to utilize available moisture more efficiently and thus offer better scope for rainfed cropping. The crop also survives even at moderate salinity and alkalinity conditions. There is no other legume crop so hardy and drought tolerant as cluster bean, which is especially suited for soil and climate of Rajasthan. India is the source of about 80 per cent (3 million hectares) of the world production (**Anonymous, 2011**). Seed of cluster bean contain 28-33 % gum India leads the list of the major guar producing countries of the world contributing to around 75 to 80% in the world's total production of around 7.5 to 10 lakh tonnes. In India Rajasthan is leading

Comment [Ma3]: Not in references.

producer of the guar seed and guar gum. It around 70% of the total production in India. Haryana and Gujarat have second and third position respectively. Rajasthan has an area of 30 lakh hectare, production of 15.46 lakh tones with a productivity of 515 kg ha⁻¹. Haryana and Gujarat state themselves at the second and third positions regarding the production in India (**Kherawat *et al.*, (2013)**). The pods of cluster bean are as rich in food value as that of French bean. According to **Aykroyd (1963)** the composition of cluster bean is 81.0 (g) moisture, 10.8 (g) carbohydrate, 3.2 (g) protein, 1.4 (g) of fat, 1.4 (g) of minerals, 0.09 (mg) thiamine, 0.03 (mg) riboflavin, vitamin C, vitamin A (100 g⁻¹) of edible portion.

Vermicompost

Vermicompost is the product of the composting process using various species of worms, usually red wigglers, white worms and other earthworms, to create a mixture of decomposing vegetable or food waste, bedding materials, and vermicast. Vermicast (also called worm castings, worm humus, worm manure, or worm feces) is the end-product of the breakdown of organic matter by earthworms. These castings have been shown to contain reduced levels of contaminants and a higher saturation of nutrients than the organic materials before vermicomposting. Vermicompost is rich in NPK (nitrogen 2- 3%, phosphorus 1.55-2.25% and potassium 1.85-2.25%), micronutrients, beneficial soil microbes and also contain plantgrowth hormones & enzymes. (**Gandhi, 2010**)

Neem cake

Neem cake has an adequate quantity of NPK in organic form for plant growth. Being a totally botanical product it contains 100% natural NPK content and other essential micro nutrients as N (Nitrogen 2.0 to 5.0%), P (Phosphorus 0.5 to 1.0%), K (Potassium 1.0 to 2.0%),Ca (Calcium 0.5 to 3.0%), Mg (Magnesium 0.3 to 1.0%), S(Sulphur 0.2 to 3.0%), Zn (Zinc 15 to 60 ppm), Cu (Copper 4to 20 ppm), Fe (Iron 500 to 1200 ppm), Mn (Manganese 20 to 60 ppm). It is rich in both Sulphur compounds and bitter limonoids. Neem cake improves the organic matter content of the soil, helping improve soil texture, water holding capacity, and soil aeration for better root development (**Lokanadhan *et al.*, 2012**)

Comment [Ma4]: The last part of the introduction should indicate objective(s) and addition to knowledge.

Comment [Ma5]: Find a common name for the two (Vermicompost and Neem cake) as a subsection not under Introduction

Materials and Method

The experiment was conducted at research farm of department of Soil Science and Agricultural Chemistry which is situated 6 km away from Prayagraj city on the right bank

of Yamuna River, the experimental site is located in the sub-tropical region with 25°24'23" N latitude, 81°50'38" E longitude and at an altitude of 98 m above mean sea level. The area of Prayagraj district comes under subtropical belt in the South east of Uttar Pradesh, which experience extremely hot summer and fairly cold winter. The maximum temperature of the location reaches up to 46°C-48°C and seldom falls as low as 4°C – 5°C. The relative humidity ranges between 20 to 92 %. The average rainfall in this area is around 1013.4 mm annually. The soil of experimental area falls in order of *Inceptisol*. The soil samples were randomly collected from three different sites in the experiment plot prior to tillage operation from a depth of 0-15 cm. The size of the soil sample was reduced by coning and quartering the composites soil sample and was air dried passed through a 2 mm sieve for preparing the sample for physical and chemical analysis.

Comment [Ma6]: Complete Randomized Block Design (RCBD) model is recommended for this study to identify factors (treatments) that are significance or not significance.

Table 1: Treatment combinations of cluster bean

Treatment	Treatment combination
T ₁	[Farmers practice]
T ₂	[NPK @ 0% + VC @ 50% + NC @ 50%]
T ₃	[NPK @ 0% + VC @ 100% + NC @ 100%]
T ₄	[NPK @ 50% + VC @ 0% + NC @ 0%]
T ₅	[NPK @ 50% + VC @ 50% + NC @ 50%]
T ₆	[NPK @ 50% + VC @ 100% + NC @ 100%]
T ₇	[NPK @ 100% + VC @ 0% + NC @ 0%]
T ₈	[NPK @ 100% + VC @ 50% + NC @ 50%]
T ₉	[NPK @ 100% + VC @ 100% + NC @ 100%]

Table 2: Chemical Analysis of pre-sowing soil samples

Parameters	Method employed	Results
Soil pH (1:2)	(Jackson, 1958)	7.56

Soil EC (dS m ⁻¹)	(Wilcox, 1950)	0.236
Organic Carbon (%)	(Walkley and Black's, 1947)	0.481
Available Nitrogen (Kg ha ⁻¹)	(Subbaih and Asija, 1956)	217.30
Available Phosphorus (Kg ha ⁻¹)	(Olsen <i>et al.</i> , 1954)	18.63
Available Potassium (Kg ha ⁻¹)	(Toth and Prince, 1949)	209.57

Results and Discussion

After harvesting the maximum Bulk density (Mg m⁻³) of soil was recorded 1.38 Mg m⁻³ in treatment T₁ (control) and minimum Bulk density (Mg m⁻³) of soil was recorded 1.18 Mg m⁻³ in treatment T₉ (100% NPK + 100% VC + 100% NC). The maximum particle density (Mg m⁻³) of soil was recorded 2.38 Mg m⁻³ in treatment T₁ (100% NPK+ 100% VC + 100% NC) and minimum particle density (Mg m⁻³) of soil was recorded 2.14 Mg m⁻³ in treatment T₉ (control). The maximum soil pore space was recorded 48.22 % in treatment T₉ (100% NPK+ 100% VC + 100% NC) and minimum soil pore space was recorded 44.82% in treatment T₁ (Control). The maximum soil pH was recorded 7.56 in treatment T₁ (control) and minimum soil pH was recorded 7.14 in treatment T₉(100% NPK + 100% VC + 100% NC). The maximum EC (dS m⁻¹) of soil was recorded 0.270 dS m⁻¹ in treatment T₉ (100% NPK +100% VC + 100% NC) and minimum EC (dS m⁻¹) of soil was recorded 0.236 dS m⁻¹ in treatment T₁ (control). The maximum % organic carbon in soil was recorded 0.545% in treatment T₉ (100% NPK +100% VC + 100% NC) which was significantly higher than any other treatment combination and the minimum % Organic carbon in soil was recorded 0.481 % in treatment T₁ (control). Legumes have potential to improve soil nutrients status through biological nitrogen fixation and incorporation of biomass into the soil as green manure. The maximum available Nitrogen in soil was recorded 276.49 kg ha⁻¹ in treatment T₉ (100% NPK +100% VC + 100% NC) which was significantly higher than any other treatment combination and the minimum available Nitrogen in soil was recorded 217.30 kg ha⁻¹ in treatment T₁ (control). The increase in available Nitrogen in soil after crop harvest by VC and NC seed inoculation might be due to increased efficiency of Nitrogen fixing capacity and nodule formation. The maximum available Phosphorus in soil was recorded 30.74 kg ha⁻¹ in treatment T₉ (100% NPK +100% VC + 100% NC) which was significantly higher than any other treatment combination and the minimum available Phosphorus in soil was recorded 18.63 kg ha⁻¹ in treatment T₁ (control). The maximum available potassium in soil was recorded 262.65 kg ha⁻¹ in treatment T₉ (100% NPK +100% VC+ 100% NC)

which was significantly higher than any other treatment combination and the minimum available potassium in soil was recorded 209.57 kg ha⁻¹ in treatment T₁ (control).

Table 3: Physical properties of soil sample after harvesting of Cluster bean

Treatment	Bulk Density (Mg m ⁻³)	Particle Density (Mg m ⁻³)	Pore space (%)
T1	1.38	2.38	44.82
T2	1.27	2.20	47.15
T3	1.26	2.16	48.96
T4	1.32	2.34	44.18
T5	1.26	2.26	46.97
T6	1.16	2.12	47.19
T7	1.31	2.42	45.03
T8	1.27	2.20	46.19
T9	1.18	2.14	48.22




Fig 1: Physical properties of soil sample after harvesting of cluster bean

Comment [Ma7]: Use either table or chart since the two represent the same data.

Table 4: Chemical properties of soil sample after harvesting of cluster bean

Treatments	pH	EC (dSm ⁻¹)	Organic Carbon (%)	Available Nitrogen (Kg ha ⁻¹)	Available Phosphorus (Kg ha ⁻¹)	Available Potassium (Kg ha ⁻¹)
T ₁	7.56	0.236	0.481	217.30	18.63	209.57

T ₂	7.43	0.257	0.488	222.43	22.67	213.28
T ₃	7.36	0.259	0.511	227.31	21.95	215.51
T ₄	7.41	0.240	0.512	233.54	26.28	219.77
T ₅	7.40	0.252	0.532	237.31	26.06	230.55
T ₆	7.30	0.265	0.573	243.50	28.79	232.60
T ₇	7.19	0.247	0.521	250.48	27.19	251.00
T ₈	7.17	0.248	0.522	265.66	30.09	254.45
T ₉	7.14	0.270	0.545	276.49	30.74	262.65




Fig. 2: Chemical properties of post-harvest soil

Conclusion

On the basis of above finding, it is concluded that application of NPK @ 100 % + NC @ 100 % + VC @ 100 % provided the significantly highest vegetative growth as well as yield attributes and yield and showed positive effect on net return up to ₹ 82992.08 ha⁻¹ with C:B ratio of 1:2.52. The available nitrogen, phosphorus and potassium content of soil after harvest of crop was increased significantly with application of NPK @ 100 % + NC @ 100 % + VC @ 100 % . The bulk density, particle density, pH, electrical conductivity of soil after harvest stage of crop were found to be non-significant while, the organic carbon,

Comment [Ma8]: It is only inferential statistics such as RCBD model that this claim can be established with certain probability.

nitrogen, phosphorus, nitrogen, porosity and water holding capacity of soil after harvest stage of crop were significant with different treatments of Inorganic fertilizer Vermicompost and Neem Cake application. Therefore, it is suggested that application of NPK @ 100 % + NC @ 100 % + VC @ 100 % found most suitable dose for cluster bean to obtain higher yield.

Comment [Ma9]: Suggestion is not conclusion.

References

Anonymous. Ministry of Agricultural, Government of India, 2011.

Aykroyd WR. The nutritive value of Indian foods and planning of satisfactory diet. Indian council of medical research, special report, series, 1963, 42.

Bouyoucos, G. J. (1927). The hydrometer as a new method for the mechanical analysis of soils. *Soil Sci*, **23**: 393-395.

Comment [Ma10]: Cite at least four more current articles of not more than three years from the date of publication. Make sure all articles cited in the text are cited in the references and vice versa.

Brown, S. and Cotton, M. (2011). Changes in soil properties and carbon content following compost application. *Compost Science and Utilization*. **19** (1): 88-97..

Chaudhari, J.A. (2018). Effect of organic sources of nutrients and biofertilizers on growth, yield and quality of clusterbean (*Cyamopsis tetragonoloba* L.) cv. Pusa Navbahar. Thesis, M.Sc. (Unpublished). S.D. Agricultural University, Sardarkrushinagar, Gujarat.

Comment [Ma11]: Not cited in the text.

Chavan, B. L., Vedpathak, M. M. and Pirgonde, B. R. (2015) Effects of organic and chemical fertilizers on cluster bean (*Cyamopsis tetragonoloba*). *European Journal of Experimental Biology*, **5**(1), 34-38.

Gandhi A, Sivakumar K. Impact of Vermicompost carrierbased bioinoculants on the growth, yield and quality of rice (*Oryza sativa* L.) cv. nlr 145. *The Ecoscan*. 2010; **4**(1):83-88.

Ghaffari, A., Klumperink, E. A., Soer, M. C., & Nauta, B. (2011). Tunable high-Q N-path band-pass filters: Modeling and verification. *IEEE Journal of Solid-State Circuits*, **46**(5), 998-1010.

Jackson, ML (1958) The pH was determined in 1:2 soil water suspensions using digital pH meter

Lokanadhan S, Muthukrishnan P, Jeyaraman S. Neem products and their agricultural operations. *Journal of Biopesticide*. 2012; 5:72-73.

Olsen, S.R., Watanale, F. S., (1954) A method to determine a phosphorus adsorption measured by colorimetric method *soil sci.soc.Am.J.21,144-149*

Singh, R.P., Bisen. J.S., Yadav, P.K., Singh, S.N. Singh, R. K. and Singh, J. (2008) Integrated use of sulphur and molybdenum on growth, yield and quality of black gram. *Legume Research* 31:214-217

Subbiah. B.V. and Asija (1956) A rapid procedure for estimation of available nitrogen in soil *curr. Sci.* 25:259.

Toth, S.J. A. L. Prince (1949) Estimation of cation exchange capacity and exchangeable calcium, potassium and sodium contents of soils by flame photometer techniques. *Soil sci.* 67. 439-445.

Wilcox LV. Electrical conductivity Am. water works Assoc. J 1950; 42:775-776. Yadav SL, Tarence T, Yadav KC, Serawat R, Seervi KS. Response of different levels of NPK and neem cake on soil health, growth and yield attributes of field pea (*Pisum sativum* L.) cv. T-19. *Journal of Pharmacognosy and Phytochemistry*. 2018; 7(3):2773-2776.

UNDER PEER REVIEW