

Original Research Article

Growth and yield of urdbean [*Vigna mungo* (L.) Hepper] influenced by vermicompost and vermiwash

ABSTRACT

The field experiment was conducted at Agronomy farm, S.K.N. College of Agriculture, Jobner (Rajasthan) to evaluate the Growth and yield of urdbean [*Vigna mungo* (L.) Hepper] influenced by vermicompost and vermiwash. The field experiment was carried out during *Kharif*, 2020 on loamy sand soil. The experiment comprising 20 treatment combinations with different doses of vermicompost and vermiwash was carried out in a 3 times replicated factorial randomised block design. Results revealed that application of vermicompost 4.5 t/ha significantly enhanced the growth attributes of urdbean viz., plant height, number of branches/plant, dry matter accumulation at 50 DAS and at harvest, CGR (Crop Growth Rate), LAI (Leaf Area Index), chlorophyll content, number of total and effective root nodules, fresh and dry weight of nodules. It also recorded significantly higher seed, stover and biological yields (940, 1920 and 2860 kg/ha) over lower levels. Results further showed that application of vermiwash 10% foliar spray at the time of flower initiation stage also resulted in significant increase in the above all mentioned parameters.

Key words: *Foliar spray, LAI, Seed yield, Stover yield, Urdbean, Vermicompost, Vermiwash*

1. INTRODUCTION

India is the largest pulse growing country contributing 9-10 % in the national food basket. In India, pulses are produced with a minimum use of resources. Pulses are a 'Smart Food' as these are critical for food basket. In addition, pulses are highly water efficient crops which can grow in drought prone areas and help in improving soil fertility by fixing atmospheric nitrogen in soil. As a result of stagnant pulse production, the per capita availability of pulses has been decreased from 60 gm in the year 1951 to 43 gm/day in the year 2016 [4]. Urdbean is a pulse grown in Indian subcontinent. Like its relative, the mung bean, it has been reclassified from the Phaseolus to the Vigna genus. Black gram originated from India, where it has been in cultivation from ancient time and is one of most highly prized pulses of India. Urdbean is an erect, dense, annual plant having tap root system with smooth, rounded nodules. The pods are narrow, cylindrical and upto six cm long. The plant grows 30-80 cm long with large hairy leaves and 4-6 cm seed pods. Urdbean contains 24% protein, 60% carbohydrate, 1.3% fat, 3.2% minerals, 0.9% fiber. It also contains calcium 54 mg, phosphorus 385 mg, iron 9.1 mg/100 gm and the calorific value of 347 Kcal/100 gm. It is consumed in the form of 'dal' (whole or split, husked and unhusked) and used as nutritive fodder for animals. It is also a green manuring crop. Urdbean can be grown in all seasons i.e. *Kharif, Rabi* and *Summer* varying upon different geographical area and climatic condition. Among different sources of organic manure,

Comment [A1]: This term is little known in the international scientific community. Therefore, I recommend that it be better explained in the text, citing for example, what season it is at the location of the experiment.

Comment [A2]: What were the doses? This information should be reported.

Comment [A3]: Write the abbreviation for gram correctly: "g" and not "gm".

Comment [A4]: Are "urdbean" and "black gram" the same bean varieties? If true, choose only one term to write the manuscript.

Comment [A5]: Write in brackets the station for "Kharif" and for "Rabi".

vermicompost is the most important manure for nutrient supplementation or management for increasing crop production. The vermicomposting technology has a benefit over composting, which is generally due to 'humus' content in vermicompost ejected by earth-worms and in conventional composting system it takes a very long time to form humus, because of slow rotting process of organic matter. It is the best method among all other composting techniques as it is cost effective, eco-friendly and very effective for sanitization of solid waste more particularly against the waste of biological origin. It is considered as a bio-oxidative process in which earthworms and microbes along with other degradable community interact and accelerate the decomposition process of organic waste [16]. Vermicompost has been reported for several beneficial effects in soil such as improvement in chemical, physical and biological conditions and can be used effectively to achieve sustainable agricultural growth [5]. Vermiwash is coelomic fluid extraction which contains enzymes, plant growth hormones, vitamins, macro and micro nutrients. Vermiwash is an organic liquid fertilizer, which have excellent growth promoting effects besides serving as biopesticides for plants [7]. Vermiwash has potential application in sustainable development in agriculture with respect to its origin, cost effectiveness, availability, reproductibility, reliability as well as biopesticide and ecofriendly soil conditioner which makes it more useful [20]. Vermiwash keeps providing the biological available nutrients to plants over a period of time and it also has biochemical factors which is not present in the conventional compost and also not found in the chemical fertilizers. Vermiwash can also reduce the use of chemical fertilizers, pesticides [2].

2. MATERIAL AND METHODS

The field trial was conducted in 2020 at the S.K.N. College of Agriculture's Agronomy Farm in Jobner (26°05' N, 75°28' E), Rajasthan, India. The study site is located in Rajasthan's agroclimatic zone IIIa (Semi-Arid Eastern Plains Zone) at an altitude of 427 metres above mean sea level. This region often has a semi-arid environment with temperature extremes in both the summer and the winter. The temperature can get as high as 48°C in the summer and as low as -2.5°C in the winter. Frost is frequent throughout the winter. This tract experiences 350 to 450 mm of rainfall on average annually, the majority of which is brought on by the south-west monsoon in the months of July and August. In the winter and summer, there is hardly any rain. At the college meteorological observatory, the average weekly meteorological observations for the entire crop season were noted. The soil had an electrical conductivity of 1.42 dS/m of saturation extract at 25°C and had an alkaline loamy-sand composition (coarse sand 25.2, fine sand 57.1, silt 9.3, and clay 7.8%). The soil had low available nitrogen (124.8 kg/ha) and organic content (0.18%) while, medium available phosphorus (16.53 kg/ha) and potassium (153.8 kg/ha). The field experiment, which included 20 treatment combinations, was replicated three times and used a factorial randomised block design. It included five treatments of vermicompost (control, 1.5 t/ha, 3.0 t/ha, 4.5 t/ha, and 6.0 t/ha) and four treatments of vermiwash foliar spray (control, 5%, 10%, and 15%). In accordance with the treatments per plot, vermicompost was applied before to the final ploughing. Vermiwash was foliar sprayed on at the flower initiation stage using a knapsack sprayer. The experiment has 60 plots of 4.0 m x 2.4 m size each. The variety 'Uttara' of urdbean was sown after two ploughings (disc ploughing than cross ploughing) followed by planking on 10th July 2020 at the depth of 4-5 cm with the help of 'Kera' method using a seed rate of

20 kg/ha. The experiment was conducted on organic land and any kind of chemicals or fertilizers were not used in the entire period. Thinning, hoeing and weeding were done after 15 and 20 DAS, respectively to maintain row to row and intra row distance of 30 X 10 cm and keep the field weed free and maintain proper aeration in soil. The experiment was basically planned for semi-arid eastern zone but under dry spell, two supplemental irrigations were given.

Comment [A6]: It remains to be explained what the procedure was for the statistical analyses.

3. RESULT AND DISCUSSION

3.1 Effect on Growth parameters

Vermicompost : The increase in levels of vermicompost a significant differences were observed in plant height, number of branches/plant, dry matter accumulation and CGR at 50 DAS and at harvest, leaf area index, chlorophyll content, number of total and effective root nodules and fresh and dry weight of root nodules. Significantly highest plant height (30.2 cm) & (36.0 cm) and no. of branches/plant (11.9) & (13.0) were found with the application of vermicompost 6.0 t/ha whereas, vermicompost 4.5 t/ha remain at par in both the parameters as compare with other treatments at 50 DAS and At harvest. Along with that dry matter g/m row length (79.9) & (116.3) and CGR g/m²/day (2.47) & (1.40) also remained significantly highest with the application of vermicompost 6.0 t/ha at 50 DAS and At harvest however it remained at par with vermicompost 4.5t/ha. The significantly highest leaf area index (3.89), chlorophyll content (3.89 mg/g) effective nodules (30.5), fresh weight of nodule (102.9 mg) and dry weight of nodule (59.1 mg) were found with the application of vermicompost 6.0 t/ha. Whereas, it was also found to be at par with the application of vermicompost 4.5 t/ha as compare with other level of vermicompost. The application of vermicompost to urdbean had a positive impact on growth and biomass production. Vermicompost improves the physical and biological qualities of soil, including the availability of nearly all of the needed plant nutrients for plant growth and development. As a result, balanced nutrition in a favourable environment may have contributed in the formation of new tissues and development, resulting in an increase in plant height, dry matter accumulation, CGR, leaf area index and chlorophyll content. Findings of the present investigation were in close agreement with those of [1], [3], [21]. Higher vegetative growth was observed in the current study with the application of vermicompost at 4.5 t/ha, indicating that it increased the number of beneficial microbes in the root zone during the early stages of infection and stimulated nodule growth by releasing bacterial cells in the cortex [6] which resulted growth in number of effective nodules and their weight. Similar results were also obtained by [12],[18].

Comment [A7]: The manuscript has a serious error in the methodology that seriously compromises the interpretation of the data obtained, as well as the conclusions: the authors did not use adequate statistical procedures to analyze and interpret the results. The dose factor of vermicompost is quantitative. Therefore, regression analysis should have been performed. The authors opted for the comparison of means. And this hides the information that there is a mathematical relationship between the increase in doses in each of the characteristics evaluated.

Comment [A8]: Considering that the experiment used a factorial design, the authors did not report whether or not there was an interaction effect.

Comment [A9]: The interpretation of the results and discussion should be based on regression analysis. And it should be reported if there was interaction between the two factors.

Vermiwash : With the increase in the per cent of vermiwash a significant effect was observed on plant height, number of branches/plant, dry matter and CGR. Application of vermiwash 15% had given significantly highest plant height (29.7 cm) and (35.5 cm), number of branches/plant (11.9) and (13.0) however, the dry matter g/m row length (77.1) & (112.6) and CGR g/m²/day (2.37) & (1.36) also, remained significantly highest with Application of vermiwash 15% at 50 DAS and At harvest. The per cent of vermiwash had also effected the leaf area index, chlorophyll content, number of total and effective root nodules and fresh and dry weight of root nodules. The significantly highest leaf area

Comment [A10]: The discussions presented here were given as if there was no effect of interaction. If true, this should be explained.

index (3.26), chlorophyll content (3.82 mg/g), effective nodules (30), fresh weight of nodule (102.8 mg) and dry weight of nodule (58.7 mg) were also found with the application of vermiwash 15%. Whereas, it was found to be at par with the application of vermiwash 10% as compared with other level of vermiwash. It is well known that foliar spray of vermiwash in urdbean crops provides nearly all of the critical plant nutrients required for plant growth and development. The higher amount of organic carbon, nitrogen, phosphorus, potassium, copper, zinc, calcium and magnesium nutrients, vitamins, amino acids and plant growth hormones found in vermiwash could be attributed to the fact that vermiwash 10 % spray was responsible for rapid growth and development of plants. It also has a lot of microorganisms that fix nitrogen and dissolve phosphate [15], [19] reported comparable results in blackgram. Higher chlorophyll content and leaf area due to vermiwash treatment resulted in higher photosynthesis and a faster growth rate, which may had influenced growth parameters such as plant height, branch number, dry matter accumulation, CGR and number and weight of effective nodules. The results of current study were also consistent with previous findings of [13], [14].

Table 1. Effect of varying levels of vermicompost and vermiwash on plant height, number of branches/plant, dry matter accumulation and CGR of urdbean at different stages

Treatments	plant height (cm)		number of branches/plant		dry matter accumulation (g/m row length)		CGR (g/m ² /day)	
	50 DAS	At Harvest	50 DAS	At Harvest	50 DAS	At Harvest	25-50 DAS	50 DAS-at harvest
Vermicompost								
Control	25.5	29.6	8.1	8.4	61.4	88.5	1.87	1.04
1.5 t/ha	27.0	31.8	9.4	10.0	66.8	96.4	2.05	1.14
3.0 t/ha	28.6	33.7	10.6	11.6	71.8	104.7	2.21	1.26
4.5 t/ha	29.9	35.6	11.5	12.7	76.9	112.4	2.37	1.36
6.0 t/ha	30.2	36.0	11.9	13.0	79.9	116.3	2.47	1.40
SEm±	0.43	0.57	0.24	0.29	1.55	2.38	0.05	0.03
CD (P = 0.05)	1.24	1.64	0.69	0.83	4.43	6.81	0.13	0.08
Vermiwash								
Control	26.4	31.1	9.1	9.7	63.6	93.2	1.95	1.13
5%	27.8	32.7	10.1	10.9	69.8	100.5	2.15	1.18
10%	29.0	34.2	10.9	11.8	75.0	108.3	2.31	1.28
15%	29.7	35.5	11.2	12.2	77.1	112.6	2.37	1.36
SEm±	0.39	0.51	0.21	0.26	1.38	2.13	0.04	0.03
CD (P = 0.05)	1.11	1.46	0.61	0.74	3.96	6.09	0.12	0.07

Table 2. Effect of varying levels of vermicompost and vermiwash on leaf area index, chlorophyll content, number of total and effective nodules, fresh and dry weight of nodules of urdbean

Treatments	leaf area index	Chlorophyll content (mg/g)	Total nodules	Effective nodules	Fresh weight of nodules (mg)	Dry weight of nodules (mg)
Vermicompost						
Control	2.83	2.83	26.8	25.2	95.3	47.5

1.5 t/ha	3.19	3.19	28.8	26.8	98.3	52.1
3.0 t/ha	3.52	3.52	30.5	28.4	100.4	56.6
4.5 t/ha	3.80	3.80	32.2	30.1	102.3	58.9
6.0 t/ha	3.89	3.89	32.8	30.5	102.9	59.1
SEm±	0.09	0.09	0.47	0.47	1.65	0.96
CD (P = 0.05)	0.25	0.25	1.36	1.36	4.72	2.74
Vermiwash						
Control	2.74	2.94	27.5	26.0	96.2	50.3
5%	3.01	3.34	29.0	27.4	98.7	53.2
10%	3.18	3.69	31.7	29.3	101.5	57.3
15%	3.26	3.82	32.6	30.0	102.8	58.7
SEm±	0.07	0.08	0.42	0.42	1.47	0.86
CD (P = 0.05)	0.19	0.23	1.21	1.21	4.22	2.45

3.2 Effect on Yield

Vermicompost : The application of 4.5 t/ha vermicompost improved seed yield (940 kg/ha), stover yield (1920 kg/ha) and biological yield (2860 kg/ha) which was significantly higher over control, vermicompost 1.5 t/ha and vermicompost 3.0 t/ha, while remained at par with vermicompost 6.0 t/ha (Table 3). The significant rise in seed yield under the effect of vermicompost was mostly due to enhanced growth and as a result an improvement in many yield parameters. Humic acid in vermicompost increases the availability of both native and added micronutrients in soil, resulting in increased plant growth, yield characteristics and yield. These results were in line with those of [10], [11], [17].

Vermiwash : The foliar spray of vermiwash resulted an increase in seed yield, stover yield and biological yield of urdbean over control (Table 3). In urdbean, vermiwash 10% spray was found to be comparable to vermiwash 15% spray in terms of yield. The increased effectiveness of foliar spraying of nutrients and growth stimulants to plants due to vermiwash spraying could be the reason for the improved yield qualities. When foliar spray of vermiwash was done, smaller amounts of IAA and GA present in vermiwash could have created stimuli in the plant system, which increased the production of growth regulator in the cell system and the action of growth regulators in the plant system stimulated the necessary growth and development, resulting in better yield. Crop yield is the result of a complex interaction of physiological and biochemical processes that alter the anatomy and morphology of growing plants. Similar set of findings were also reported by [9],[8].

Table 3. Effect of varying levels of vermicompost and vermiwash on seed, stover and biological yields of urdbean

Treatments	Yield (kg / ha)		
	seed	Stover	biological
Vermicompost			
Control	660	1456	2116
1.5 t/ha	770	1673	2443

3.0 t/ha	864	1822	2686
4.5 t/ha	940	1920	2860
6.0 t/ha	985	1951	2936
SEm±	22	28	48
CD (P = 0.05)	62	81	138
Vermiwash			
Control	759	1625	2384
5%	820	1734	2554
10%	885	1832	2717
15%	910	1866	2776
SEm±	20	25	43
CD (P = 0.05)	56	72	123

4. CONCLUSION

Based on the results of one year experimentation, it may be inferred that application of vermicompost @ 4.5 t/ha was found better for obtaining higher plant height, number of branches/plant, dry matter accumulation, number of effective nodules and their weight with higher seed yield (940 kg/ha) in urdbean. Similarly, foliar spray of vermiwash demonstrated significant positive effect on the growth characters and yield of urdbean. Vermiwash 10 % spray had been the most effective vermiwash spray for enhancing productivity (885 kg/ha). Through this study, it is evident that the utilization of these organic amendments can enhance various important parameters associated with urdbean. However, further research is warranted to optimize the application rates and explore the long-term effects of vermicompost and vermiwash on urdbean.

REFERENCE

- [1] Alkobaisy, J.S. and Mutlag, N. Effect of the use of vermicompost and rhizobial inoculation on some soil characteristics, growth and yield of mungbean (*Vigna radiata* L.). Iraqi Journal of Agricultural Sciences. 2021; **52**(1): 163-169.
- [2] Bansod, V.M. Effect of eisenia fetida vermiwash on the growth of dianthus plant. International Journal of Multidisciplinary Educational Research. 2020; **9**(1): 450-454.
- [3] Choudhary, V.K., Kumar, P.S. and Bhagawati, R. Influence of organic nutrient sources on growth, seed yield and economics of cowpea under mid hills of Arunachal Pradesh. Journal of Food Legumes. 2013; **26**(3): 51-54.
- [4] DAC, 2018. Pulses in India: Retrospect and Prospects 2018, Directorate of Pulses Development GOI, Ministry of Agriculture and Farmers Welfare, Department of Agriculture, Cooperation and Farmers Welfare. 2018.
- [5] Das, S., Teron, R., Duary, B., Bhattacharya, S.S. and Kim Ki-Hyun. Assessing C–N balance and soil rejuvenation capacity of vermicompost application in a degraded landscape: A study in an alluvial river basin with *Cajanus cajan*. Environmental Research. 2019; **177**: 108591.
- [6] Ghanshyam, K.R. and Jat, R.K. Productivity and soil fertility as affected by organic manure and inorganic fertilizer in green gram (*Vigna radiata*) wheat (*Triticum aestivum*) system. Indian Journal of Agronomy. 2010; **55**: 16-21.

Comment [A11]: The conclusion was compromised due to the statistical procedure used by the authors. Clearly, there was a mathematical relationship between the increase in vermicompost doses and each of the evaluated characteristics. But, this was not shown by the authors. Could it be that the effect of the vermiwash doses was the same in each of the vermicompost doses?? This will only be possible to know if something is informed about the interaction.

- [7] Jandaik, S., Kumar, V. and Thakur, P. Vermiwash: Plant growth enhancer and antifungal agent. *International Journal of Extensive Research*. 2015; **2**: 38-41.
- [8] Kapase, P. V., Deotale, R. D., Sawant, P. P., Sahane, A.N. and Banginwar, A.D. Effect of foliar sprays of humic acid through vermicompost wash and NAA on morpho-physiological parameters, yield and yield contributing parameters of chickpea. *Journal of Soils and Crops*. 2013; **24**(1): 107-114.
- [9] Khairnar, A.V. and Gunjal, B.S. Effect of potash fertilization and foliar spray of vermiwash on growth and yield of green gram (*Vigna radiata* L.). *International Journal of Agriculture Sciences*. 2012; **8**(1): 307-308.
- [10] Kumawat, N., Singh, R.P., Kumar, R., Yadav, T.P. and Hariom. Effect of integrated nutrient management on productivity, nutrient uptake and economics of rainfed pigeonpea (*Cajanus cajan*) and blackgram (*Vigna mungo*) intercropping system. *Indian Journal of Agricultural Sciences*. 2010; **85**(2): 171-6.
- [11] Meena, J.S., Verma, H.P. and Pancholi, P. Effect of fertility levels and biofertilizers on yield, quality and economics of cowpea. *Agriculture for Sustainable Development*. 2014; **2**: 162-164.
- [12] Mehta, H.K., Reddy, R.S., Jayamma, P. and Kumar, R.N. Influence of biofertilizers, vermicompost and chemical fertilizers on growth, nodulation, nutrient uptake, seed yield and economics of black gram. *International Journal of Applied Biology and Pharmaceutical Technology*. 2012; **6**: 249-251.
- [13] Meyyappan, M. And Sivakumar, G. Effect of foliar application of liquid organic and inorganic fertilizers along with NAA on cowpea (*Vigna unguiculata*). *Annals of Plant and Soil Research*. 2020; **22**(4): 454-456.
- [14] Nath, G. and Singh, K. Effect of foliar spray of biopesticides and vermiwash of animal, agronomy and kitchen wastes on soybean. *Botany Research International*. 2011; **4**(3): 52-57.
- [15] Nath, G. and Singh, K. Combined effect of vermiwash with biopesticides against infestation of pod borer (*Helicoverpa armigera* hub.). *International Journal of Zoological Investigations*. 2015; **1**(1): 40-51.
- [16] Patwa, A., Parde, D., Dohare, D., Vijay, R. and Kumar, R. Solid waste characterization and treatment technologies in rural areas: An Indian and international review. *Environmental Technology & Innovation*. 2020; **20**: 101066.
- [17] Raghawendra, S. and Kedar, P. Effect of vermicompost, rhizobium and DAP on growth, yield and nutrient uptake by chickpea. *Journal of Food Legumes*. 2008; **21**(2): 112- 114.
- [18] Rekha, G.S., Valivittan, K. and Kaleena, P.K. Studies on the influence of vermicompost and vermiwash on the growth and productivity of black gram (*Vigna mungo*). *Advances in Biological Research*. 2010; **7**(4): 114-121.
- [19] Tharmaraj, K., Ganesh, P., Kalanjunathan, K., Kumar, Suresh, R. and Anandan, A. Influence of vermicompost and vermiwash on physio chemical properties of black gram cultivated soil. *International Journal of Recent Scientific Research*. 2010; **3**: 77-83.
- [20] Varghese, S.M. and Prabha, M.L. Biochemical characterization of vermiwash and its effect on growth of *Capsicum frutescens*. *Malaya Journal of Biosciences*. 2014; **1**(2): 86-91.

[21]Yadav, A.K., Ramawat, N. and Singh, D. Effect of organic manures and biofertilizers on growth and yield parameters of cowpea. Journal of Pharmacognasy and Phytochemistry. 2019; **8**(2): 271-274.

UNDER PEER REVIEW