

## Effect of Zinc and Salicylic acid on growth and yield of green gram (*Vignaradiata*L.)

### ABSTRACT

A field experiment was carried out through the growth season 2022 at Zaid Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj (U.P). To study the effect of Zinc and Salicylic acid on improve growth green gram (*Vignaradiata*L) in sandy loam. The soil of experimental plot was sandy loam in texture, nearly neutral in soil reaction (pH 7.1), low in organic carbon (0.36 %), available N (171.48 kg/ha), available P (15.2 kg/ha) and available K (232.5 kg/ha). The experiment was laid out on Randomized Block Design with nine treatments each replicated thrice on the basis of one year experimentation. The treatments which are T<sub>1</sub>: Zn (75PPM) + Salicylic acid (100PPM), T<sub>2</sub>: Zn (75PPM) + Salicylic acid (150PPM), T<sub>3</sub>: Zn (75PPM) + Salicylic acid (200PPM), T<sub>4</sub>: Zn (100PPM) + Salicylic acid (100PPM), T<sub>5</sub>: Zn (100PPM) + Salicylic acid (150PPM), T<sub>6</sub>: Zn (100PPM) + Salicylic acid (200PPM), T<sub>7</sub>: Zn (125PPM) + Salicylic acid (100PPM), T<sub>8</sub>: Zn (125PPM) + Salicylic acid (150PPM), T<sub>9</sub>: Zn (125PPM) + Salicylic acid (200PPM) are used. The results showed that treatment T<sub>9</sub>- Zn (125PPM) + Salicylic acid (200PPM). Result was recorded significantly higher growth parameters i.e. Plant height (35.14 cm), Plant dry weight (6.64g/plant) and Crop growth rate (2.99g/m<sup>2</sup>/day). However, yield attributes and yield parameters like No. of Pods/plant (21.22), No. of seeds/pod (11.35), Test weight (34.04 g), Seed yield (1100.97 kg/ha), straw yield (4471.00 kg/ha) were affected with the treatment T<sub>9</sub>- Zn (125PPM)+Salicylic acid (200PPM) than other treatments. Higher gross returns (Rs.82572.75/ha), net return (Rs. 56228.32/ha) and benefit cost ratio (2.13) was obtained in the treatment T<sub>9</sub>- Zn (125PPM) + Salicylic acid (200PPM) as compared to other treatments.

**Key words:** *Growth, Salicylic acid, yield, Zinc.*

## Introduction

Mung bean (*Vignaradiata*L.) is also known as green gram, it is an important pulse crop of India and grown in Rabi (South India), Kharif and Zaid seasons. It is green with husk and yellow when dehusked. The beans are small, ovoid in shape and green in color. “The mung bean (*Vignaradiata*), alternatively known as the moongbean, monggo, green gram, or mungSanskritmugd,isaplantspeciesinthelegumefamily.Themungbeanismainlycultivated in India, Pakistan, Bangladesh, Nepal, China, Korea, South Asia and Southeast Asia. It is used as an ingredient in both savory and sweet dishes. Mungbeanis third most important pulse crop of India after chickpea and pigeon pea. The nutritive value of mungbeanis a high with easily digestible protein (approximately 25-28%), oil 1.0-1.5%, fiber 3.5-4.5%, ash 4.5-5.5%, carbohydrate62-65%,water9.1%,andvitaminsondryweightbasis.Mungbean(*Vignaradiata* L. Wilczek) is a summer pulse crop with short duration (70-90 days) and high nutritive value.It has many effective uses, green pod is cooked as peas, sproutrichin vitamins and amino acids” (Sharma.,2012).

“Micronutrients are essential for the normal growth of plants, deficiencies of which adverselyaffectthegrowth,metabolismandreproductivephaseinplants.Inmanypartsofthe country, zinc (Zn) as a plant nutrient stands third in importance i.e., next to nitrogen and phosphorus. In the recent years, zinc is considered as one of the constraints in the optimum production of crops. It plays a vital role in synthesis of chlorophyll, protein and nucleic acid and helps in the utilization of nitrogen and phosphorus by plants as it acts as an activator of dehydrogenaseandproteinaseenzymes,directlyorindirectlyinsynthesisofcarbohydratesand protein” (Katyal and Sharma1991).

“Plantgrowthhormonesregulate almosteveryphaseofplantgrowthanddevelopment. Plant growth and development are known to beunder the control of internal and external factors. The effect of growth regulators is found to be largely dependent upon various factors as concentration and type of growth regulator, method of application, time of application, soil type and other conditions. Salicylic acid (SA) is an endogenous plant hormone that has been found to play a major role in the regulation of plant growth and development, including seed germination, organ differentiation, stomatal movement, photoperiodic responses, and senescence mediation”(Hayat *et al.*,2010).

## Materials and Methods

The present examination was carried out during 2022 at Zaid Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj, UP, which is located at 25.28°N latitude, 81.54°E longitude and 98 m altitude above the mean sea level. The experiment laid out in Randomized Block Design which consisting of ten treatments with T<sub>1</sub>: Zn (75PPM) + Salicylic acid (100PPM), T<sub>2</sub>: Zn (75PPM) + Salicylic acid (150PPM), T<sub>3</sub>: Zn (75PPM) + Salicylic acid (200PPM), T<sub>4</sub>: Zn (100PPM) + Salicylic acid (100PPM), T<sub>5</sub>: Zn (100PPM) + Salicylic acid (150PPM), T<sub>6</sub>: Zn (100PPM) + Salicylic acid (200PPM), T<sub>7</sub>: Zn (125PPM) + Salicylic acid (100PPM), T<sub>8</sub>: Zn (125PPM) + Salicylic acid (150PPM), T<sub>9</sub>: Zn (125PPM) + Salicylic acid (200PPM), are used. The experimental site was uniform in topography and sandy loam in texture, nearly neutral in soil reaction (pH 7.1), low in Organic carbon (0.38%), medium available N (225 kg ha<sup>-1</sup>), higher available P (19.50 kg ha<sup>-1</sup>) and medium available K (213.7 kg ha<sup>-1</sup>). In the period from germination to harvest several plant growth parameters were recorded at frequent intervals along with it after harvest several yield parameters were recorded those parameters are growth parameters, plant height, and plant dry weight are recorded. The yield parameters like Pods/plant, No. of grains/pod, Test weight, seed yield, straw yield and harvest index were recorded.

Study was statistically analyzed using analysis of variance (ANOVA) as applicable to Randomized Block Design (Gomez and Gomez 1984).

## Results and Discussion

### Growth attributes

#### Plant height

Significantly maximum Plant height (35.14 cm) was recorded with the treatment with Zn (125PPM) + Salicylic acid (200PPM) over the other treatments. However, treatments with Zn(125PPM)+Salicylicacid(150PPM)(34.99cm)whichwerefoundtobestatisticallyatpar with Zn (125PPM) + Salicylic acid (200PPM) as compared to othertreatments.

Zinc application in different physiological processes like enzyme activation, electron transport, chlorophyll formation, stomatal regulation, etc. With the increase in levels of zinc the plant height gradually increased, which might be attributable to greater photosynthetic activity and chlorophyll synthesis due to zinc fertilization resulting into better vegetative growth. Similar results were reported by **Masihet *al.* (2020)**.

### Plant dry weight (g/plant)

Data in Table (1) show that the effect of Zn (125PPM) + Salicylic acid (200PPM) was significantly maximum dry weight (6.64g/plant) compared the other treatments. However, treatments with Zn (125PPM) + Salicylic acid (150PPM) (6.56 g/plant), Zn (100PPM) + Salicylic acid (200PPM) (6.40 g/plant), Zn (100PPM) + Salicylic acid (150PPM) (6.39 g/plant), Zn (125PPM) + Salicylic acid (100PPM) (6.26 g/plant) and Zn (75PPM) + Salicylic acid (200PPM) (6.19 g/plant) which was found to be statistically at par with Zn (125PPM) + Salicylic acid (200PPM) as compared to other treatments.

The reported positive effect of application of Zn on an enhanced branching in green gram and mainly attributed to promotion of shoot development by the auxins whereas Zn application ultimately increased the availability of other nutrients and accelerated the translocation of photo assimilates which ultimately helped in increase of plant dry weight of millet. Similar results were reported by Kumar *et al.* (2016).

The increase in dry weight with increasing levels of salicylic acid might be due to the promoted effect of salicylic acid on morphological characters which enhanced the photosynthetic rate and also maintained the stability of membrane thereby improving the dry matter production. The reports were in accordance to Keykhaet *al.* (2014).

**Table 1: Effect of Zinc and Salicylic acid on growth attributes of Green gram**

<b>Treatments</b>	<b>Plant height (cm)</b>	<b>Dry weight (g/plant)</b>
1. Zn (75PPM) + Salicylic acid (100PPM)	32.66	5.79
2. Zn (75PPM) + Salicylic acid (150PPM)	32.93	5.88
3. Zn (75PPM) + Salicylic acid (200PPM)	33.48	6.19
4. Zn (100PPM) + Salicylic acid (100PPM)	33.03	6.07
5. Zn (100PPM) + Salicylic acid (150PPM)	33.98	6.26
6. Zn (100PPM) + Salicylic acid (200PPM)	34.52	6.40
7. Zn (125PPM) + Salicylic acid (100PPM)	34.17	6.39
8. Zn (125PPM) + Salicylic acid (150PPM)	34.99	6.56
9. Zn (125PPM) + Salicylic acid (200PPM)	35.14	6.64
<b>F- test</b>	S	S
<b>S. EM (<math>\pm</math>)</b>	0.14	0.16
<b>C. D. (P = 0.05)</b>	0.41	0.49

### **Yield attributes and Yield Number of Pods/Plant**

Result in Table (2) found that the application of Zn (125PPM) + Salicylic acid (200PPM) was significantly Maximum Number of Pods/Plant (21.22) than other the treatments. However, the treatments Zn (125PPM) + Salicylic acid (150PPM) (20.79) which were found to be statistically at par with Zn (125PPM) + Salicylic acid (200PPM).

The increase in number of pods per plant under increasing levels of salicylic acid might be due to reduced drop due to efficient translocation of photosynthesis from source to sink. These findings were obtained by Sruthiet al. (2020).

### **Number of Grains/pods**

Result in Table (2) show that the significantly maximum Number of Grains/pod (11.35) as affected with the application of Zn (125PPM) + Salicylic acid (200PPM) over all the treatments. However, the treatments Zn (125PPM) + Salicylic acid (150PPM) (11.16) which were found to be statistically at par with Zn (125PPM) + Salicylic acid (200PPM).

Result in Table (2) indicator that the application of Zinc to green gram crop generally improves ear head growth by synthesizing tryptophan and auxin. The enhancement effect on pods/plant, seeds/pod and attributed to the favorable influence of the Zn application to crops on nutrient metabolism, biological activity and growth parameters and hence, applied zinc resulted in taller and higher enzyme activity which in turn encouraged more number of pods/plants, seeds/pod and seeds test weight. Similar findings have been reported earlier by Mahilane and Singh (2018).

### **Test Weight (g)**

Significantly Maximum Test Weight (34.04 g) was recorded with the treatment of application of Zn (125PPM) + Salicylic acid (200PPM) over all the treatments. However, the treatments Zn (125PPM) + Salicylic acid (150PPM) (33.73 g) which were found to be statistically at par with Zn (125PPM) + Salicylic acid (200PPM).

### **Seed yield (kg/ha)**

Significantly highest Seed Yield (1100.97 kg/ha) was recorded with the treatment of application of Zn (125PPM) + Salicylic acid (200PPM) over all the treatments. However, the treatments Zn (125PPM) + Salicylic acid (150PPM) (1081.50 kg/ha) which were found to be statistically at par with Zn (125PPM) + Salicylic acid (200PPM).

Zinc plays a vital role in increasing seed yield because zinc takes place in many physiological process of plant such as chlorophyll formation, stomatal regulation, starch utilization which enhance seed yield. Zinc also converts ammonia to nitrate in crops which

contribute to yield. These results are in confirmatory with the work of **Debnath *et al.* (2016)**.

**Straw yield (kg/ha)**

Significantly highest Seed Yield (4471.00 kg/ha) was recorded with the treatment of application of Zn (125PPM) + Salicylic acid (200PPM) over all the treatments. However the treatments Zn (125PPM) + Salicylic acid (150PPM) (4370.59 kg/ha) were found to be statistically at par with Zn (125PPM) + Salicylic acid (200PPM).

Increased in straw yield might be due to the growth promoting effect of salicylic acid which increased the level of cell division within the apical meristem of seedling root and caused higher plant growth and increased the dry matter production. Similar findings were also corroborated by Laishram *et al.* (2020).

**Table 2: Effect of Zinc and Salicylic acid on Yield attributes and Yield of Green gram**

<b>Treatments</b>	<b>Pods/plant</b>	<b>Grains/pod</b>	<b>Test Weight (g)</b>	<b>Seed Yield (kg/ha)</b>	<b>Stover Yield (kg/ha)</b>
1. Zn (75PPM) + Salicylic acid (100PPM)	16.67	9.05	31.43	796.30	3560.41
2. Zn (75PPM) + Salicylic acid (150PPM)	17.39	9.35	31.63	825.04	3678.90
3. Zn (75PPM) + Salicylic acid (200PPM)	18.39	9.72	32.42	926.37	3944.34
4. Zn (100PPM) + Salicylic acid (100PPM)	17.94	9.53	32.17	857.21	3772.54
5. Zn (100PPM) + Salicylic acid (150PPM)	19.02	10.05	32.68	973.58	4012.50
6. Zn (100PPM) + Salicylic acid (200PPM)	19.98	10.83	33.32	1031.80	4272.03
7. Zn (125PPM) + Salicylic acid (100PPM)	19.58	10.48	33.13	1000.78	4153.63
8. Zn (125PPM) + Salicylic acid (150PPM)	20.79	11.16	33.73	1081.50	4370.59
9. Zn (125PPM) + Salicylic acid (200PPM)	21.22	11.35	34.04	1100.97	4471.00
F test	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>
S. EM ( $\pm$ )	0.15	0.06	0.11	19.28	37.69
CD (P = 0.05)	0.44	0.19	0.32	57.81	155.70

**CONCLUSION**

It is concluded that application of treatment T<sub>9</sub>- Zn (125PPM) + Salicylic acid (200PPM) was recorded significantly higher Grain yield (1100.97 kg/ha), higher gross returns (Rs. 82572.75/ha), net return (Rs. 56228.32/ha) and benefit cost ratio (2.13) of Green gram as compared to other treatments. Since, the findings based on the research done in one season, further trails may be required for further confirmation.

## REFERENCES

- Debnath, P., Hemalatha, S. and Bhowmik, S. 2016. Effect of yield, Quality parameters and nutrient uptake studies of biofortified corn with soil and foliar applied zinc. *Ecology, Environment and conservation Paper*, 22; 97-103.
- Hayat, Q., Hayat, S., Irfan, M. and Ahmad, A. (2010) Effect of exogenous salicylic acid under changing environment: A review. *Environmental and Experimental Botany*. 68,14-25.
- Katyal JC, Sharma.(1991). DTPA extractable and total Zn, Cu, Mn and Fe in indian soils and their association with soil properties. *Geoderma*.1991; 49:165-179.
- Keykha, M., Ganjali, H. R. and Mobasser, H. R. (2014). Effect of SA and GA, on Mungbean. *International Journal of Biosciences*. 5(11):70-75.
- Kumar, R., Rathore, D.K., Singh, M., Kumar, P. and Khippal, A. 2016. Effect of phosphorous and zinc nutrition on growth and yield of fodder cowpea. *Legume Research*, 39 (2): 262-267.
- Laishram, B., Basanta Singh, T., Athokpam Kalpana, Merinda Wangkheirakpam, Sunil Kumar Chongtham and Jiten Singh, W. 2020. Effect of Salicylic Acid and Potassium Nitrate on Growth and Yield of Lentil (*Lens culinaris* L.) under Rainfed Condition. *International Journal of Current Microbiology and Applied Sciences*, 9(11): 2779- 2791.
- Masih, A., Dawson, J. and Singh, R. E. 2020. Effect of Levels of Phosphorus and Zinc on Growth and Yield of Greengram (*Vigna radiata* L.). *International Journal of Current Microbiology and Applied Sciences*, 9(10); 3106-3112.
- Mahilane, C. and Vikram Singh. 2018. Effect of Zinc and Molybdenum on Growth, Yield Attributes, Yield and Protein in Grain on Summer Blackgram (*Vigna mungo* L.). *International Journal of Current Microbiology and Applied Sciences*, 7(01): 1156-1162.
- Sharma RK. Effect of salicylic acid and gibberellic acid on seed germination and growth of pea. *Internat. J Plant Sci*. 2012; 7(2):322-324.

Sruthi, D. S. V. S., Singh, S., Tiwari, D. and Kavya, P. 2020. Effect of Foliar spray of nutrient and plant growth regulators on growth and yield of green gram (*Vignaradiata*L.).*The Bioscan*,15(2); 253-2