

Original Research Article
**Nutritional and Antioxidant Properties of
Resistant Starch-Based Flour Blends from
Unripe Plantain, Pigeon Pea and Rice-Bran**

ABSTRACT

Aim:

The nutritional potentials of resistant starch-based flour blends from unripe plantain, pigeon pea and rice bran as functional food products were investigated.

Place and Duration of Study:

Food Chemistry laboratory, Department of Food Science and Technology, and Functional Foods Laboratory, Department of Biochemistry, Federal University of Technology, Akure, Nigeria. From January, 2022 to January, 2023.

Methodology:

Native plantain starch was isolated and its resistant starch was prepared via acetylation process. The flours of native plantain starch, pigeon pea and rice bran were prepared, and their blends obtained by using Mixture design expert 11.0.0. The dependent variables were proximate and minerals compositions, amino acid and nutritional qualities profiles, α -amylase and α -glucosidase activities and antioxidant potentials of the blends. Resistant starch-based flour from unripe plantain, pigeon pea and rice bran flour blends coded; PLF – (100% Plantain flour) were used as control; PLPRF – Plantain flour:Pigeon pea flour:Rice bran flour (73.08:15.15:11.76); PSPRF – Native Plantain starch:Pigeon pea flour:Rice bran flour (73.08:15.15:11.76); AP1SPRF – Acetylated Plantain starch:Pigeon pea flour:Rice bran flour (73.08:15.15:11.76); AP2SPRF – Acetylated Plantain starch:Pigeon pea flour:Rice bran flour (70.67:18.15:11.18). The proximate, mineral, Amino acid profile, α -amylase and α -glucosidase and in-vitro antioxidant potentials of the flour were compared with the commercial flour product CRF – 100% Cerolina flour (wheat and soybean produced by More Foods Lagos, Nigeria).

Results:

The overall results showed that resistant starch-based flour blends had reduced moisture content, as low as 2.40%, increased ash content of 4.63% and reduced fat content of 2.42%. K, Cu, Fe, Na, Zn and P were more relatively abundant in the flour blends than the 100% plantain flour. The modified resistant starch-based flour blends were able to reduce Fe^{2+} to Fe^{3+} , scavenge DPPH, FRAP and ABTS and inhibited α -amylase and α -glucosidase activities. The chemical compositions improved with the addition of quality protein which extensively improved the mineral contents in terms of calcium, potassium, iron, zinc and phosphorus.

Conclusion:

The flour blends of AP1SPRF and AP2SPRF studied were a good source of resistant starch and bioactive ingredients that would be useful in a variety of dietary functional food products.

Keywords: Resistant starch, acetylated starch, antioxidant, α -amylase, α -glucosidase

1. INTRODUCTION

The quest for quality functional foods is influenced by lot of factors such as health issues (obesity, diabetes, cholesterol, cancer, etc.), developmental stages in life and lifestyles. The development of novel foods products, using new technologies has influenced consumers' perception on the awareness of the nutritional value and health benefits of novel foods [1].

Recently, tremendous importance has been given to functional foods, because it provides health benefits by reducing the risk of chronic diseases apart from the nutritional benefits [1]. Resistant starch (RS), basically recognized as a type of dietary fiber, is one of the most widely used ingredients in functional foods. RS is a dietary starch with unique property of not being absorbed intact by the gut, but hydrolyzed by intestinal enzymes and fermented by bacteria in the colon to produce short-chain fatty acids like butyrate [2]. Recently, RS has drawn interest of so many researchers due to its health benefits, functional peculiarities and promising physiological benefits for non-diabetic and diabetic people, which include reduction in the amplified levels of blood glucose and insulin, restraining of colonic cancer and risks of ulcerative colitis, positive impact on bowel movements, increase in the assimilation of minerals, prebiotic actions, laxation, hypocholesterolemic and hypoglycemic effects, and enhanced fat oxidation [2,3]. Several studies have revealed that the starches modified via acetylation result in the production of food starch products with enough resistant starch content, capable of serving as raw materials for diabetes treatment [4,5]. In the acetylation process, the hydroxyl groups of the glucose monomers are converted to the groups CH_3COO^- , leading to a decrease in the gelatinization temperature, an increase or decrease in the swelling power, and solubility along with the storage stability [6].

Antioxidants are compounds that prevent deterioration of lipids during or after food processing by scavenging the free radicals [7], thereby increasing the oxidative stability of food, controlling the pro-oxidants and other oxidative intermediates [8]. They are phytochemicals, vitamins and nutrients that protect our cells from damage caused by free radicals by stabilizing or deactivating free radicals before they attack cells [8]. Consumption of foods rich in antioxidant plays a special role in prevention and management of several human ailments including diabetes mellitus [7].

Plantain (*Musa paradisiaca*) has a thick cell-wall that cannot be easily penetrated by enzymes, leading to low glycemic index. It is a cheap source of energy, and also, medically recommended for diabetic patients, due to its high resistant starch [9]. Its fruit pulp has been reported to possess high antioxidant properties [10]. The extract of unripe plantain has potential to inhibit the key enzymes linked with type 2-diabetes [11].

Pigeon pea (*Cajanus cajan*) cultivars have good quality resistant starch, minerals, medicinal and antimicrobial properties, and 36.5% protein, rich in sulphur containing amino acids [12] in addition to excellent water retention (250.3 mL/100g), fat absorption (130 mL/100g), emulsification capacity (120%) and foaming capacity (130%) [13].

Rice bran is a by-product of the rice milling process, and it contains various antioxidants, 12-13% oil and 4.3% highly unsaponifiable components. It is also rich in dietary fibers (α -glucan, pectin, and gum), non-lignified cell walls, and tocotrienol, γ -oryzanol, and α -sitosterol, which lower the plasma levels of the various parameters of the lipid profile [14].

Despite the increasing research interest in the production of functional foods from local crop materials to manage chronic diseases [15], there is scanty information on the combination of plantain, pigeon pea and rice bran flours for food products formulation. Hence, the aim of this study was to produce and evaluate nutritional composition and antioxidants properties of resistant starch based-dough meal from plantain, pigeon pea and rice bran.

2. MATERIALS AND METHODS

2.1 Sources of Materials

Matured unripe plantains (*Musa ABB*) were obtained from a local farm in Sabongida-Ora, Owan-West LGA, pigeon peas were obtained from Jattu Market, Etsako-West LGA, and rice bran was obtained from Pemos Foods, Aviele, Etsako-West LGA, Edo State. The food materials were authenticated at the Department of Crop Soil and Pest Management, Federal University of Technology, Akure, Nigeria.

2.2 Flour Sample Preparation

The unripe plantain was processed into flour using the method of Oluwajuyitan et al. [16] with slight modifications. The plantain was manually peeled, sliced (0.5 cm thick), washed, oven dried (Plus 11 Sanyo Gallenkamp PLC, Loughborough, Leicestershire, UK) at 60 °C for 24 h,

cooled in a desiccator, milled (Rico MG 1803 Mixer Grinder, 1000 watts, India), sieved through 200 μm mesh sieve and packaged in airtight polythene prior to analyses. The pigeon peas were handpicked, oven dried (Plus11 Sanyo Gallenkamp PLC, Loughborough, Leicestershire, UK) at 60 °C for 24 h, cooled in a desiccator, milled (Rico MG 1803 Mixer Grinder, 1000watts, India), sieved through 200 μm mesh sieve and packaged in airtight polythene prior to analyses. The rice bran was washed with distilled water, drained and oven dried (Plus11 Sanyo Gallenkamp PLC, Loughborough, Leicestershire, UK) at 60 °C for 24 h, cooled in a desiccator, milled (Rico MG 1803 Mixer Grinder, 1000watts, India), sieved through 200 μm mesh sieve and packaged in airtight polythene prior to analyses.

2.3 Acetylation of Plantain Starch

Prior to acetylation of plantain starch, the native starch of the plantain was isolated according to the method described by Oladebeye et al. [17] with slight modifications. The plantains were blended, sieved through muslin cloth, supernatant decanted, the cake dried (Plus11 Sanyo Gallenkamp PLC, Loughborough, Leicestershire, UK) at 60 °C for 24 h, cooled, milled (Rico MG 1803 Mixer Grinder, 1000watts, India), sieved through 200 μm mesh sieve and packaged in airtight polythene prior to modification via acetylation. For acetylation process, 100 g (dwb) of the native starch was dispersed in 200 mL of acetic anhydride in a reaction flask and stirred at 500 rpm with a mechanical stirrer for 5 min in the presence of NaOH solution (50 g NaOH/100 g water) as a catalyst at 100 °C and stirred for 1 h. To precipitate the starch, 100 mL of ethyl alcohol solution (96%) was added followed by filtration by suction with a Buchner filter funnel (Whatman filter No. 4). The residue was washed with ethyl alcohol and then with distilled water till most of the acetic anhydride was removed. The resulting paste produced by these washes was dried in an oven (Plus11 Sanyo Gallenkamp PLC, Loughborough, Leicestershire, UK) at 40 °C for 16 h up to approximately 9% moisture content [18].

2.4 Preparation of Resistant Starch-Based Flour Blends

Flour samples of plantain, pigeon pea and rice-bran were blended with reference to 14 g/day protein and 5 g/day fibre (i.e. 25% of recommended daily intakes of an adult requirements), using Optimal Mixture Design Methodology. The flour samples were blended as CRF – 100% Cerolina flour (wheat and soybean produced by More Foods Lagos, Nigeria); PLF – (100% Plantain flour) were used as control; PLPRF – Plantain flour:Pigeon pea flour:Rice bran flour (73.08:15.15:11.76); PSPRF – Native Plantain starch:Pigeon pea flour:Rice bran flour (73.08:15.15:11.76); AP1SPRF – Acetylated Plantain starch:Pigeon pea flour:Rice bran flour (73.08:15.15:11.76); AP2SPRF – Acetylated Plantain starch:Pigeon pea flour:Rice bran flour (70.67:18.15:11.18).

2.5 Determination of Proximate Compositions

The moisture, crude protein, fat, fiber and ash contents of the food samples were determined according to AOAC methods [19]. Total carbohydrate content was calculated by Difference.

2.6 Determination of Mineral Compositions

Calcium, magnesium, iron, zinc, copper and manganese were determined using Atomic Absorption Spectrophotometer (AAS Model SP9). Sodium and potassium contents were determined using flame photometer (Sherwood, UK) using NaCl and KCl as the standard. Phosphorus was determined by the vanado-molybdate colorimetric method of AOAC [19].

2.7 Amino Acid Compositions

The amino acid compositions of the food samples were determined using Automated Amino Acid Analyzer (Model 6300; Beckman Coulter Inc., Fullerton, Calif., USA) as described by Mansouri et al. [20] and Hernández et al. [21].

2.8 Antioxidant Activities

2.8.1 In-Vitro Antioxidant Assay

Prior to in-vitro analyses, 50 g each of the samples was dissolved in 250 mL flask containing distilled water. The mixture was sieved and filtered, to obtain clear solution. The solutions extracted were used for the in-vitro after concentrating using freeze drying methods. The in-vitro antioxidant assays of the aqueous extracts of the food samples were carried out using standard methods. The free radical scavenging activity of the food samples was determined against 2,2-azino-bis(3-ethylbenzthiazoline-6-sulphonate (ABTS) [22]. The 2,2-diphenyl-1-picrylhydrazyl (DPPH) free radical scavenging was determined as described by Aluko and Monu [23]. Metal chelating activity of the formulated food aqueous extract was determined using the method of Xie et al. [24]. Ferric reducing antioxidant activity (FRAP) was determined according to Zhang and Lin [25].

2.8.2 Inhibition of α -amylase Activity

α -amylase activity was used to measure in-vitro by hydrolysis of starch in presence of α -amylase enzyme [26]. α -amylase activity was carried out by starch-iodine method. 10 μ L of α -amylase solution (0.025 mg/mL) was mixed with 390 μ L of phosphate buffer (0.02 M containing 0.006 M NaCl, pH 7.0) containing 20 μ L of 50 μ g/ml concentration of extracts. After incubation at 37 °C for 10 min, 100 μ L of starch solution (1%) was added and the mixture was re-incubated for 1 h. Additions of 0.1 mL of 1% iodine solution and 5 mL distilled water were done before taking the absorbance at 565 nm. Sample, substrate and α -amylase blank determinations were carried out under the same reaction conditions.

$$\text{Inhibition of enzyme activity (\%)} = \frac{A - C}{B - C} \times 100$$

where, A = absorbance of the sample, B = absorbance of blank (without α -amylase), and C = absorbance of control (without starch).

2.8.3 Inhibition of α -glucosidase Activity

The α -glucosidase inhibition activity was performed according to the method of Tiwari et al. [27] with slight modification. The final volume of the reaction mixture was 100 μ L, which contained 70 μ L of phosphate buffer saline (50 mM, pH 6.8), 10 μ L of 50 μ g/ml concentration of extracts, and 10 μ L (0.057 U) enzyme. The content was mixed, pre-incubated at 37 °C for 10 min and pre-read against the reagent blank value by spectrophotometry at 400 nm. The reaction was initiated using 10 μ L of 0.5 mM substrate (i.e., p-nitrophenol glucopyranoside). Acarbose was used as a positive control. After incubation at 37 °C for 30 min, optical absorbance was measured against the reagent blank value by spectrophotometry at 400 nm. The percentage of enzyme inhibition was calculated.

$$\% \text{ scavenging activity} = \frac{A_c - A_s}{A_c} \times 100$$

where, A_c is the absorbance of control and A_s the absorbance of the extract.

3. STATISTICAL ANALYSIS

The statistical analysis of the data obtained was carried out with IBM SPSS 26.0 software for mean comparison, using Duncan's least significant test and one-way analysis of variance (ANOVA) at 5% significance level.

4. RESULTS AND DISCUSSION

4.1 Proximate Compositions

Proximate composition of the flour blends is presented in Table 1. The moisture content ranged between 2.40% in AP2SPRF to 10.43% in PLPRF. All the values are lower than the 10% recommended for a longer storage stability of food products with no adverse effect on the quality attributes of the flour blends except for sample PLPRF, which is higher by 0.43 [28], indicating higher shelf-life in the Acetylated flour blends than the other blends [29]. The

highest protein content observed in CRF being the positive control sample with wheat and soy beans compositions. The other flour blend ranges between 13.85% and 6.91%. This is similar to the report that protein content can be improved with a better source of protein like bambara groundnut, soybeans and pigeon peas and this will help to boost the protein intake of consumers who cannot afford animal-based protein [30]. The crude fibre content ranges between 2.40% in CRF to 14.28% in PLPRF, and this high crude fibre could be due to the addition of the rice bran in the flour blends. Fat contents of the flour blends range from 2.42% in AP1SPRF to 7.74% in CRF, which may be due to the addition of soybeans, an oil seed crop, in the formulation of CRF. Also, the ash content ranges from 1.30% in PSPRF to 4.63% in AP2SPRF. High ash content in AP2SPRF (acetylated starch-based flour sample) is an indication of relative abundance and nature of minerals [31], and the degree of digestibility of the sample [32].

Table 1. Proximate compositions of the blends from plantain flour, starch, acetylated starch with pigeon pea and rice bran and commercial product

Sample	Moisture (%)	Ash (%)	Crude Protein (%)	Fat Content (%)	Crude Fibre (%)	Carbohydrates By Difference (%)	Energy (Kcal)
PLF	8.25 ^b ±0.02	1.97 ^d ±0.02	4.38 ^f ±0.02	2.96 ^g ±0.02	3.18 ^e ±0.02	78.26 ^a ±1.71	377.73 ^b ±0.18
CRF	5.77 ^d ±0.02	1.90 ^d ±0.02	18.31 ^a ±0.02	7.74 ^a ±0.02	2.40 ^f ±0.02	63.88 ^c ±0.02	415.01 ^a ±0.33
PLPRF	10.43 ^a ±0.03	3.48 ^c ±0.02	13.85 ^b ±0.02	4.98 ^c ±0.02	14.28 ^a ±0.02	52.98 ^d ±0.02	325.51 ^f ±0.35
PSPRF	6.88 ^c ±0.02	1.30 ^e ±0.02	10.30 ^c ±0.02	5.35 ^b ±0.02	12.91 ^b ±0.02	63.26 ^c ±0.02	357.10 ^e ±0.18
AP1SPRF	2.59 ^e ±0.03	4.24 ^b ±0.15	8.21 ^d ±0.03	2.42 ^f ±0.02	7.23 ^d ±0.10	75.31 ^b ±0.06	372.58 ^c ±0.27
AP2SPRF	2.40 ^f ±0.12	4.63 ^a ±0.06	6.91 ^e ±0.06	2.65 ^e ±0.03	9.13 ^c ±0.06	74.27 ^b ±0.26	365.00 ^d ±1.02

PLF = 100% Plantain flour; CRF = 100% Cerolina flour; PLPRF = Plantain flour:Pigeon pea flour:Rice bran flour (73.08:15.15:11.76); PSPRF = Plantain starch:Pigeon pea flour:Rice bran flour (73.08:15.15:11.76); AP1SPRF = Acetylated Plantain starch:Pigeon pea flour:Rice bran flour (73.08:15.15:11.76); AP2SPRF = Acetylated Plantain starch:Pigeon pea flour:Rice bran flour (70.67:18.15:11.18)

Results are the means of triplicate determination ± standard deviation values in the same column with the same superscripts letters (a > b > c > d) are not significantly different (P < 0.05)

The carbohydrate content ranges from 52.98% in PLPRF to 78.26% in PLF. The carbohydrate content decreases with an increase in the percentage of protein, ash and crude fibre. The energy value ranges from 325.51 kcal/100 g in PLPRF to 415.01 kcal/100 g in CRF. High energy value has been attributed to fat content, and this is an index of the total amount of energy the human body could generate during its metabolism [33]. Recently, both high-carbohydrate, low-fat (HCLF) diets and low-carbohydrate, high-fat (LCHF) diets have been identified as having non-significant difference in terms of changes in lean mass, fat mass, systolic blood pressure, diastolic blood pressure, triglycerides and glucose, implying that both diets are effective for weight control and reduction of cardiovascular risk factors [34].

4.2 Mineral Compositions

The minerals compositions determined in the flour blends are sodium (Na), potassium (K), calcium (Ca), iron (Fe), copper (Cu), Phosphorus (P) and zinc (Zn) Lead was not detected in all the samples (Table 2). Dietary minerals (also known as mineral nutrients) are the chemical elements required by living organism. Potassium is more abundant in all the minerals determined. This is the confirmation of the relative abundance of ash contents in flour sample as indicated in (Table 1). Potassium, being the most abundant intracellular cation, maintains intracellular fluids balance and transmission of nerves in relation with sodium [35]. Dietary sodium is needed by all ages in very small quantity [36] with the view to effectively controlling the blood pressure [37]. Importantly, lowering the blood pressure implies lowering the sodium intake and simultaneously increasing the potassium intake [38]. Getting enough calcium and phosphorus at infancy to adulthood is through adequate diet, which will prevent the risk of weak teeth and bones, and fractures in later life. Additional roles played by calcium in the body are clotting of blood, neuro-sensing, contraction and relaxation of muscles, release of

hormones and maintaining normal heartbeat of appetite, fatigue, nausea, muscle cramps, sleeplessness, hyperactivity, irritability, anxiety in infants and children [39].

Table 2. Mineral compositions of the blends from plantain flour, starch, acetylated starch with pigeon pea and rice bran and commercial product

Sample	Mineral (mg/100g)						
	K	Ca	Cu	Fe	Na	Zn	P
PLF	57.53 ^d ±0.04	3.23 ^c ±0.02	0.04 ^b ±0.02	0.33 ^a ±0.03	17.25 ^e ±0.02	0.69 ^e ±0.02	27.63 ^d ±0.03
CRF	27.75 ^f ±0.02	6.19 ^a ±0.02	0.11 ^a ±0.02	0.55 ^d ±0.02	28.53 ^a ±0.04	0.99 ^a ±0.02	34.04 ^a ±0.02
PLPRF	67.00 ^a ±0.02	2.88 ^e ±0.02	0.11 ^a ±0.02	1.41 ^a ±0.02	18.75 ^b ±0.02	0.86 ^b ±0.02	32.81 ^b ±0.02
PSPRF	51.00 ^e ±0.02	2.93 ^d ±0.03	0.08 ^a ±0.02	0.84 ^c ±0.02	18.25 ^d ±0.02	0.70 ^e ±0.03	31.60 ^c ±0.02
AP1SPRF	62.75 ^b ±0.02	3.38 ^b ±0.02	0.10 ^a ±0.02	1.14 ^b ±0.02	17.00 ^f ±0.02	0.74 ^d ±0.02	31.60 ^c ±0.54
AP2SPRF	62.23 ^c ±0.25	3.38 ^b ±0.03	0.09 ^a ±0.01	1.17 ^b ±0.06	18.57 ^c ±0.06	0.81 ^c ±0.01	31.55 ^c ±0.09

Results are the means of triplicate determination ± standard deviation values in the same column with the same superscripts letters (a > b > c > d) are not significantly different (P < 0.05)

The Iron and the zinc are the least abundant out of all the mineral analyzed ranging from 0.33 mg/100g Fe (PLF) to 1.41 mg/100g Fe (PLPRF) and from 0.69 mg/100g Zn (PLF) to 0.99 mg/100gZn (CRF). Flour blends with high calories, nutrient density and adequate iron content as well as high nutrient bioavailability can be used to prevent iron deficiency in all ages [40].

4.3 Amino Acid Compositions

The amino acid profiles of flour blends are depicted in Table 3. The total amino acids in each of the flour blends is total 100 (g/100g protein). The total non-essential amino acids of flour blends range from 58.31 (g/100g protein) in PLF to 64.82 (g/100g protein) in CRF while the total essential amino acids of flour blends range from 35.18 (g/100g protein) in CRF to 41.81 (g/100g protein) in PLF. There is a significant increase in TEAAs, that is, lysine, leucine, isoleucine, tryptophan, arginine, threonine, valine, histidine, phenylalanine, tyrosine, and methionine, obtained in all the resistant starch-based flour samples compared to the control sample CRF (wheat and soybean-based). The increase in the amino acid profile of resistant starch-based flour blends is due to the high quality of protein present in pigeon pea flour, because legumes are protein-rich crops and have higher amino-acid composition than plantain [30].

Table 3a. Essential amino acids (g/100g protein)

Essential Amino Acid	Sample					
	CRF	PLF	PLPRF	PSPRF	AP1SPRF	AP2SPRF
Histidine	2.92 ^e	8.95 ^a	5.80 ^b	4.14 ^d	4.55 ^c	4.28 ^{cd}
Isoleucine	4.25 ^{bc}	3.91 ^d	3.89 ^d	4.12 ^c	4.43 ^a	4.27 ^b
Leucine	7.44 ^a	6.08 ^c	7.07 ^b	7.45 ^a	7.47 ^a	7.58 ^a
Lysine	4.84 ^d	5.88 ^c	6.55 ^a	5.99 ^c	6.43 ^{ab}	6.34 ^b
Methionine	1.60 ^c	2.11 ^a	1.86 ^b	1.71 ^{bc}	1.87 ^b	1.75 ^{bc}
Phenylalanine	5.11 ^d	4.73 ^e	7.33 ^a	7.07 ^b	6.74 ^c	7.26 ^{ab}
Threonine	3.37 ^c	3.95 ^a	3.71 ^b	3.67 ^b	3.89 ^a	3.72 ^b
Tryptophan	1.27 ^{ab}	1.37 ^a	1.05 ^c	1.10 ^c	1.15 ^{bc}	1.15 ^{bc}
Valine	4.38 ^d	4.71 ^{ab}	4.55 ^c	4.59 ^{bc}	4.79 ^a	4.65 ^{abc}

Means along the row with different superscripts are significantly different at 5% level with a > b > c. Mean separation done by Duncan Multiple Range Test

Same trend is observed in terms of hydrophilic, hydrophobic, positively charged, branched chain and aromatic amino acids in resistant starch-based samples, especially the native and acetylated starch-based samples compared to CRF (wheat-based). This is an added advantage as this sample can be eaten as a functional food, because hydrophobic amino acids act as antioxidants by increasing the solubility of peptides in lipids, which then facilitates better interaction with free radicals [30]. The ArAA (aromatic amino acids) are made up phenylalanine, tryptophan, and tyrosine. They are essential amino acids, which the body

cannot synthesize and must therefore be consumed in foods. Except PLF (100% plantain flour), all the plantain-based, plantain starch-based flour blends are more abundant in ArAA than CRF. However, the extent of acetylation does not have significant difference ($P < 0.05$) on the relative abundance of ArAA in the food samples.

Table 3b. Non-essential amino acids (g/100g protein)

Non-Essential Amino Acid	Sample					
	CRF	PLF	PLPRF	PSPRF	AP1SPRF	AP2SPRF
Alanine	3.61 ^f	5.08 ^a	4.44 ^c	4.22 ^e	4.67 ^b	4.30 ^d
Arginine	5.77 ^c	8.25 ^a	6.76 ^a	6.32 ^{bc}	6.25 ^{bc}	6.10 ^c
Asparagine	8.95 ^d	12.53 ^a	11.26 ^b	10.34 ^c	11.29 ^b	11.06 ^b
Cysteine	2.02 ^a	1.85 ^b	1.43 ^c	1.43 ^c	1.23 ^d	1.38 ^c
Glutamine	25.68 ^a	13.79 ^f	18.20 ^d	20.70 ^b	17.42 ^e	19.44 ^c
Glycine	3.77 ^c	4.42 ^a	3.75 ^c	3.64 ^c	4.00 ^b	3.64 ^c
Proline	7.22 ^a	3.92 ^f	4.36 ^e	5.26 ^b	4.53 ^d	4.82 ^c
Serine	4.89 ^d	5.60 ^b	5.22 ^c	5.07 ^c	6.01 ^a	5.21 ^c
Tyrosine	2.93 ^a	2.86 ^a	2.76 ^a	3.19 ^a	3.30 ^a	3.07 ^a

Means along the row with different superscripts are significantly different at 5% level with $a > b > c$. Mean separation done by Duncan Multiple Range Test

The respective values of EAAI, PER, BV and NI of the flour blends range from 99.92 (CRF) to 118.49% (PLF), 1.99 (PLF) to 2.65 g/100 g (AP2SPRF), 97.21 (CRF) to 117.45% (PLF) and 7.87 (AP2SPRF) to 18.30% (CRF). The values of EAAI, PER and BV show that all the plantain-based and plantain starch-based flour blends are capable of improving the adequate intake of nutrients and amino acids by infants and adults.

Table 3c. Nutritional Quality

Parameter	Sample					
	CRF	PLF	PLPRF	PSPRF	AP1SPRF	AP2SPRF
TEAA	35.18 ^d	41.69 ^{ab}	41.81 ^a	39.84 ^c	41.29 ^{ab}	41.00 ^b
TNEA	64.82 ^a	58.31 ^{bc}	58.19 ^c	60.16 ^b	58.71 ^{bc}	59.00 ^c
TEAA/TNEA	0.54 ^d	0.72 ^{ab}	0.72 ^a	0.66 ^c	0.70 ^b	0.69 ^b
HAA-1	16.98 ^b	18.68 ^a	16.87 ^b	17.01 ^b	18.43 ^a	17.02 ^b
HAA-2	33.60 ^b	30.55 ^c	33.51 ^b	34.41 ^a	34.48 ^a	34.62 ^a
PCAA	11.36 ^f	19.92 ^a	16.79 ^b	14.35 ^e	15.64 ^c	14.93 ^d
NCAA	34.62 ^a	26.32 ^e	29.46 ^c	31.03 ^b	28.71 ^d	30.49 ^b
BCAA	16.06 ^c	14.70 ^e	15.51 ^d	16.16 ^{bc}	16.68 ^a	16.50 ^{ab}
SCAA	3.61 ^b	3.96 ^a	3.29 ^c	3.14 ^c	3.10 ^c	3.13 ^c
ArAA	9.31 ^b	8.95 ^b	11.14 ^a	11.36 ^a	11.18 ^a	11.47 ^a
EAAI (%)	99.92 ^d	118.49 ^a	115.49 ^{ab}	110.53 ^c	115.83 ^{ab}	113.84 ^{bc}
PER (g/100 g)	2.60 ^{ab}	1.99 ^c	2.45 ^b	2.58 ^{ab}	2.58 ^{ab}	2.65 ^a
BV (%)	97.21 ^d	117.45 ^a	114.19 ^{ab}	108.78 ^c	114.55 ^{ab}	112.39 ^{bc}
NI (%)	18.30 ^a	5.19 ^f	16.00 ^b	11.38 ^c	9.51 ^d	7.87 ^e

Means along the row with different superscripts are significantly different at 5% level with $a > b > c$. Mean separation done by Duncan Multiple Range Test

TEAA - total essential amino acids = Leusine + Lysine + Isoleucine + Phenylalanine + Tryptophan + Valine + Methionine + Histidine + Threonine; TNEA - total non-essential amino acids = Tyrosine + Cystine + Alanine + Glutamic acid + Glycine + Aspartic acid + Serine + Proline + Arginine; HAA-1, Hydrophilic amino acids = Glycine + Tyrosine + Serine + Threonine + cysteine; HAA-2, Hydrophobic amino acids = Methionine + Alanine + Valine + Leucine + Isoleucine + Proline + Phenylalanine; PCAA – positively charged amino acids = Basic amino acids = amino acids with basic group in their side chains = Lysine + Histidine + Arginine; NCAA – negatively charged amino acids = Acidic amino acids = amino acids with carboxyl group in their side chains = Glutamic acid + Aspartic acid OR ASX (asparagine +aspartic acid) + GLX (glutamine+glutamic acid); BCAA- Branch chain amino acids = amino acids having aliphatic side chains with a branch = valine, isoleucine, leucine; SCAA- sulphur containing amino acids = amino acids with sulphur in their side chains = cysteine and methionine; ArAA- aromatic amino acids = amino acids with benzene ring in their side chains = phenylalanine, tryptophan and tyrosine; EAAI (%) – Essential amino acids index; PER (g/100 g) – Protein efficiency ratio; BV (%) – Biological value; NI (%) – Nutritional Index.

4.4 Antioxidant Activities

4.4.1 In-Vitro Antioxidant Assay

The scavenging activities of the acetylated starch-based samples are significantly different from other samples (Figure 1a-d). The scavenging activity of AP1SRF is significantly higher than the native starch, 100% PLF, the control sample, CRF and other non-acetylated samples. The results suggest that acetylation increases the DPPH radical scavenging activity of the samples. These observations are consistent with the findings of the antioxidant activities of polysaccharide, which improve after acetylation [41–43]. The ABTS radical scavenging activities of the flour samples show that the resistant starch-based flour blends exhibit higher radical scavenging activities compared to the 100% CRF. The FRAP and Fe-chelating scavenging activities also follow the same trend with the acetylated samples AP1SRF and AP2SRF having the highest scavenging activity, indicating that the composite flours may have better protective effect against free radicals compared to 100% CRF (wheat-soybean based flour).




Figure 1. (a) DPPH radical scavenging ability (%), (b) ABTS radical scavenging ability (mmol.TEAC/g), (c) Fe chelation ability (%), (d) Ferric reducing antioxidant property (mg.AAE/g)

The addition of acetyl group had a significantly lifting effect on the DPPH free radical scavenging activity [44], implying a possibility of acetyl groups reducing DPPH to DPPH₂, thereby preventing lipid oxidation.

4.4.2 α -amylase and α -glucosidase Inhibition Activities

The inhibitions of α -amylase and α -glucosidase of the resistant starch-based flour blends are shown in Figure 2a, b. α -amylase and α -glucosidase are enzymes that hydrolyze carbohydrates [45]. The addition of resistant starch in the flour blends increases the percentage inhibition of the α -amylase from 38% in CRF (commercial control sample) to 40% in PLPRF, and as high as 80% in AP1SPRF. Generally, the acetylated starch-based samples, AP1SPRF and AP2SPRF, however have higher α -amylase inhibition than CRF, PLF, PLPRF and PSPRF. Heating produces resistant starches (RS 2 and 3) hence, this may account for higher inhibition by the modified starches [46,47]. The same trend is observed in terms of α -glucosidase, with the highest α -glucosidase inhibition (30% and above) in acetylated starch-based flour blends (AP1SPRF and AP2SPRF) and less than 20% in the control sample (CRF). This agrees with the observations on the inhibition of α -amylase and α -glucosidase by cooked rice [46,47]. The α -glucosidase inhibitory effect exhibited by the resistant starch-based flour blends may indicate their effective potentials in managing type-2 diabetes mellitus (T2DM) [47].




Figure 2. (a) α -amylase inhibition (%), (b) α -Glucosidase inhibition (%)

5. CONCLUSION

This study shows that resistant starch-based flour blends (RS-type 2) have high beneficial antioxidant properties, coupled high inhibitory effect against the α -amylase and α -glucosidase activities. The increase in the amino acid and nutritional quality profile of resistant starch-based flour blends is due to the high quality of protein present in pigeon pea flour, because legumes are protein-rich crops and have higher amino-acid compositions than plantain. The proximate and mineral compositions of the samples show appreciably indications of quality nutrition (protein, ash, fibre, energy, potassium, calcium, sodium iron and zinc). Hence, this study provides evidence for the application of resistant starch-based flour blends (AP1SPRF and AP2SPRF) as dietary functional foods.

CONSENT

Not applicable to this manuscript

ETHICAL APPROVAL

Not applicable to this manuscript

REFERENCES

- [1] Ashwar BA, Gani A, Shah A, Wani IA, Masoodi FA. Preparation, health benefits and applications of resistant starch - A review. *Starch/Staerke* 2016;68. <https://doi.org/10.1002/star.201500064>.
- [2] Hedemann MS, Hermansen K, Pedersen S, Bach Knudsen KE. Resistant starch but not enzymatically modified waxy maize delays development of diabetes in Zucker diabetic fatty rats. *J Nutr* 2017;147:825–34. <https://doi.org/10.3945/jn.116.243899>.
- [3] Batey I, Wrigley C, Miskelly D. Industrial Processing of Grains Into Co-Products of Protein, Starch, Oils and Fibre. *Cereal Grains Assess. Manag. Qual. Second Ed.*, 2017. <https://doi.org/10.1016/B978-0-08-100719-8.00023-1>.
- [4] Singh H, Sodhi NS, Singh N. Structure and functional properties of acetylated Sorghum starch. *Int J Food Prop* 2012;15. <https://doi.org/10.1080/10942912.2010.483633>.
- [5] Sha XS, Xiang ZJ, Bin L, Jing L, Bin Z, Jiao YJ, et al. Preparation and physical characteristics of resistant starch (type 4) in acetylated indica rice. *Food Chem* 2012;134. <https://doi.org/10.1016/j.foodchem.2012.02.081>.
- [6] Colussi R, El Halal SLM, Pinto VZ, Bartz J, Gutkoski LC, Zavareze E da R, et al. Acetylation of rice starch in an aqueous medium for use in food. *LWT - Food Sci Technol* 2015;62:1076–82. <https://doi.org/10.1016/J.LWT.2015.01.053>.
- [7] Sodipo MA, Oyeleye SI, Oboh G, Agbede JO, Oluwamukomi MO. Hypoglycemic effect of biscuits produced from flour blends of three medicinal foods on high-fat diet-streptozotocin-induced diabetic rats. *J Food Biochem* 2021;45. <https://doi.org/10.1111/jfbc.13334>.
- [8] Khan AN, Khan RA, Ahmad M, Mushtaq N, Khan A. Role of antioxidant in oxidative stress and diabetes mellitus. ~ 217 ~ *J Pharmacogn Phytochem* 2015;3:217–20.
- [9] Eleazu CO, Okafor PN, Ahamefuna I. Total antioxidant capacity, nutritional composition and inhibitory activity of unripe plantain (*Musa paradisiaca*) on oxidative stress in alloxan induced diabetic rabbits. *Pakistan J Nutr* 2010;9:1052–7. <https://doi.org/10.3923/pjn.2010.1052.1057>.
- [10] Shodehinde SA. In vitro antioxidant activities and inhibitory effects of aqueous extracts of unripe plantain pulp (*Musa paradisiaca*) on enzymes linked with type 2 diabetes and hypertension. *J Toxicol Environ Heal Sci* 2012;4. <https://doi.org/10.5897/jtehs11.087>.
- [11] Shodehinde SA, Ademiluyi AO, Oboh G, Akindahunsi AA. Contribution of *Musa paradisiaca* in the inhibition of α -amylase, α -glucosidase and Angiotensin-I converting enzyme in streptozotocin induced rats. *Life Sci* 2015;133. <https://doi.org/10.1016/j.lfs.2015.03.026>.
- [12] Narina SS, Xu Y, Hamama AA, Phatak SC, Bhardwaj HL. Effect of Cultivar and Planting Time on Resistant Starch Accumulation in Pigeonpea Grown in Virginia. *ISRN Agron* 2012;2012. <https://doi.org/10.5402/2012/576471>.
- [13] Eltayeb ARSM, Ali AO, Haron R. The chemical composition of pigeon pea (*Cajanus cajana*) seed and functional properties of protein isolate. *Pakistan J Nutr* 2010;9:1069–73. <https://doi.org/10.3923/pjn.2010.1069.1073>.
- [14] Giugliano D. Dietary antioxidants for cardiovascular prevention. *Nutr Metab Cardiovasc Dis* 2000;10.
- [15] Oluwajuyitan TD, Ijarotimi OS, Fagbemi TN. Nutritional, biochemical and organoleptic properties of high protein-fibre functional foods developed from plantain, defatted soybean, rice-bran and oat-bran flour. *Nutr Food Sci* 2020;51. <https://doi.org/10.1108/NFS-06-2020-0225>.

- [16] Oluwajuyitan TD, Ijarotimi OS. Nutritional, antioxidant, glycaemic index and Antihyperglycaemic properties of improved traditional plantain-based (Musa AAB) dough meal enriched with tigernut (*Cyperus esculentus*) and defatted soybean (*Glycine max*) flour for diabetic patients. *Heliyon* 2019;5:1504. <https://doi.org/10.1016/j.heliyon.2019>.
- [17] Oladebeye AO, Oshodi AA, Amoo IA, Karim AA. Gaseous Ozonation of Pigeon Pea, Lima Bean, and Jack Bean Starches: Functional, Thermal, and Molecular Properties; Gaseous Ozonation of Pigeon Pea, Lima Bean, and Jack Bean Starches: Functional, Thermal, and Molecular Properties 2018. <https://doi.org/10.1002/star.201700367>.
- [18] Halal SLM El, Colussi R, Pinto VZ, Bartz J, Radunz M, Carreño NLV, et al. Structure, morphology and functionality of acetylated and oxidised barley starches. *Food Chem* 2015;168:247–56. <https://doi.org/10.1016/J.FOODCHEM.2014.07.046>.
- [19] AOAC. Official Methods of Analysis of Association of Official Analytical Chemists. AOAC 2012;19st ed:121–30.
- [20] Mansouri F, Ben Moumen A, Richard G, Fauconnier ML, Sindic M, Elamrani A, et al. Proximate composition, amino acid profile, carbohydrate and mineral content of seed meals from four safflower (*Carthamus tinctorius* L.) varieties grown in north-eastern Morocco. *OCL - Oilseeds Fats, Crop Lipids* 2018;25. <https://doi.org/10.1051/ocl/2018001>.
- [21] Hernandez H, Lynn H, Bates S. A modified method for rapid tryptophan analysis of maize CENTRO Internacional Oe Mejoramiento De Maiz Y Trigo International Maize Ano Wheat Improvement Center Mexico. 1969.
- [22] Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radic Biol Med* 1999;26. [https://doi.org/10.1016/S0891-5849\(98\)00315-3](https://doi.org/10.1016/S0891-5849(98)00315-3).
- [23] Aluko RE, Monu E. Functional and bioactive properties of quinoa seed protein hydrolysates. *J Food Sci* 2003;68. <https://doi.org/10.1111/j.1365-2621.2003.tb09635.x>.
- [24] Xie Z, Huang J, Xu X, Jin Z. Antioxidant activity of peptides isolated from alfalfa leaf protein hydrolysate. *Food Chem* 2008;111. <https://doi.org/10.1016/j.foodchem.2008.03.078>.
- [25] Zhang L liang, Lin Y ming. Tannins from *Canarium album* with potent antioxidant activity. *J Zhejiang Univ Sci B* 2008;9. <https://doi.org/10.1631/jzus.B0820002>.
- [26] Hossain SJ, Tsujiyama I, Takasugi M, Islam MA, Biswas RS, Aoshima H. Total phenolic content, antioxidative, anti-amylase, anti-glucosidase, and antihistamine release activities of bangladeshi fruits. *Food Sci Technol Res* 2008;14:261–8. <https://doi.org/10.3136/fstr.14.261>.
- [27] Tiwari AK, Swapna M, Ayesha SB, Zehra A, Agawane SB, Madhusudana K. Identification of proglycemic and antihyperglycemic activity in antioxidant rich fraction of some common food grains. *Int Food Res J* 2011;18.
- [28] Bello M, Oluwamukomi M, Enujiugha V. Nutrient Composition and Sensory Properties of Biscuit from Mushroom-Wheat Composite Flours. *Arch Curr Res Int* 2017;9. <https://doi.org/10.9734/acri/2017/35686>.
- [29] Oduje AA, Ayodele AJ, Stephen AA. Assessment of the Nutritional, Anti nutritional and Antioxidant capacity of Uripe, ripe, and over ripe Plantain (*Musa paradisiaca*) Peels. *Int J Adv Res JournalwwwJournalijarCom Int J Adv Res* 2015;3.
- [30] Arise AK, Dauda AO, Awolola GV, Akinlolu-ojo TV. Physico-chemical, functional and pasting properties of composite flour made from wheat, plantain and bambara for biscuit production. vol. 18. 2017; 18(4):616–624
- [31] Harris GK, Marshall MR. Ash Analysis, 2017, p. 287–97. https://doi.org/10.1007/978-3-319-45776-5_16.
- [32] Shirley RB, Parsons CM. Effect of Ash Content on Protein Quality of Meat and Bone Meal. *Poult Sci* 2001;80:626–32. <https://doi.org/10.1093/PS/80.5.626>.
- [33] Omoba O. Optimisation of Plantain - Brewers' Spent Grain Biscuit Using Response Surface Methodology. *J Sci Res Reports* 2013;2. <https://doi.org/10.9734/jsrr/2013/4944>.

- [34] Yang Q, Lang X, Li W, Liang Y. The effects of low-fat, high-carbohydrate diets vs. low-carbohydrate, high-fat diets on weight, blood pressure, serum lipids and blood glucose: a systematic review and meta-analysis. *Eur J Clin Nutr* 2022;76. <https://doi.org/10.1038/s41430-021-00927-0>.
- [35] Akinyeye RO, Oluwadunsin A, Omoyeni A. Proximate, Mineral, Anti-Nutrients, Phyto-Chemical Screening and Amino Acid Compositions of the Leaves of *Pterocarpus Mildbraedi* Harms. *Electron J Environ Agric Food Chem* 2011;10:1848–57.
- [36] Gowrishankar M, Blair B, Rieder MJ. Dietary intake of sodium by children: Why it matters 2020. <https://doi.org/10.1093/pch/pxz153>.
- [37] Frazier KM, Bender BF, Julien G. (Digital Presentation) Point-of-Care Diagnostic Device for Na/K Urinalysis. *ECS Meet Abstr* 2022;MA2022-02:2233–2233. <https://doi.org/10.1149/ma2022-02612233mtgabs>.
- [38] Stone MS, Martyn L, Weaver CM. Potassium intake, bioavailability, hypertension, and glucose control. *Nutrients* 2016;8. <https://doi.org/10.3390/nu8070444>.
- [39] Gröber U, Werner T, Vormann J, Kisters K. Myth or reality-transdermal magnesium? *Nutrients* 2017;9. <https://doi.org/10.3390/nu9080813>.
- [40] Faleiros FTV, da Silva VN, de Assis Carvalho M, Machado NC. Intake, bioavailability, and absorption of iron in infants aged 6 to 36 months: an observational study in a Brazilian Well Child Clinic. *Nutrire* 2016;41. <https://doi.org/10.1186/s41110-016-0011-0>.
- [41] Liu J, Luo J, Ye H, Zeng X. Preparation, antioxidant and antitumor activities in vitro of different derivatives of levan from endophytic bacterium *Paenibacillus polymyxa* EJS-3. *Food Chem Toxicol* 2012;50. <https://doi.org/10.1016/j.fct.2011.11.016>.
- [42] Song Y, Yang Y, Zhang Y, Duan L, Zhou C, Ni Y, et al. Effect of acetylation on antioxidant and cytoprotective activity of polysaccharides isolated from pumpkin (*Cucurbita pepo*, lady godiva). *Carbohydr Polym* 2013;98. <https://doi.org/10.1016/j.carbpol.2013.06.049>.
- [43] Xie J-H, Zhang F, Wang ZJ, Shen MY, Nie SP, Xie MY. Preparation, characterization and antioxidant activities of acetylated polysaccharides from *Cyclocarya paliurus* leaves. *Carbohydr Polym* 2015;133:596–604. <https://doi.org/10.1016/j.carbpol.2015.07.031>.
- [44] Xue Y, Ding X, Zhang Z, Wu X, Cao X, Zhang W, Zhang J, Chen Y. Acetylation process and antioxidant activity of a polysaccharide from loquat (*Eriobotrya japonica*) leaves. *Am J Biochem Biotechnol* 2020;16. <https://doi.org/10.3844/AJBBS.2020.590.601>.
- [45] Jenkins DJA, Kendall CWC, Augustin LSA, Franceschi S, Hamidi M, Marchie A, Jenkins AL, Axelsen M. Glycemic index: overview of implications in health and disease. *Am J Clin Nutr* 2002;76:266S-273S. <https://doi.org/10.1093/AJCN/76.1.266S>.
- [46] Donkor ON, Stojanovska L, Ginn P, Ashton J, Vasiljevic T. Germinated grains - Sources of bioactive compounds. *Food Chem* 2012;135. <https://doi.org/10.1016/j.foodchem.2012.05.058>.
- [47] Olugbuyi AO, Oladipo GO, Malomo SA, Ijarotimi SO, Fagbemi TN. Biochemical Ameliorating Potential of Optimized Dough Meal from Plantain (*Musa AAB*), Soy cake (*Glycine max*) and Rice bran (*Oryza sativa*) Flour Blends in Streptozotocin Induced Diabetic Rats. *Appl Food Res* 2022;2. <https://doi.org/10.1016/j.afres.2022.100097>.