

Original Research Article

Chairux Algorithm for Divisibility Test



Abstract

This paper presents a novel Chairux Algorithm for the divisibility test. The test is based on an arbitrary integer via the concepts of Bezout's identity and the Euclidean algorithm. Examples demonstrating the effectiveness of the proposed algorithm indicate its simplicity, efficiency, and flexibility compared to the existing state-of-the-art algorithms.

1 Introduction

The problem of testing the divisibility of one integer by another has been of interest to mathematicians for centuries [3], Tiebekabe, Pagdame and Diouf, Ismaïla journal paper [6], Bennett, Albert B and Nelson, L Ted [2], Mahmood, M Khalid [5], Cherniavsky, Yonah and Mouftakhov, Artour [4], Barkema, JR [1]. Traditional methods, such as long division, require complex calculations and repeated subtractions, which can be time-consuming and error-prone. In this paper, I present a new method for testing divisibility, called Chairux Algorithm. The algorithm is based on the well-known concepts of Bezout's identity and the Euclidean algorithm, and provides a simple and efficient approach to testing divisibility.

2 Background



Before presenting Chairux Algorithm, I review the two fundamental concepts on which it is based: Bezout's identity and the Euclidean algorithm.

2.1 Bezout's Identity

Bezout's identity is a fundamental theorem in number theory that states that for any two integers a and b , there exist integers x and y such that $ax + by = \gcd(a, b)$, where $\gcd(a, b)$ is the greatest common divisor of a and b . This identity is useful in many areas of mathematics, including cryptography and algebraic geometry.

2.2 Euclidean Algorithm

The Euclidean algorithm is a classic algorithm for finding the greatest common divisor of two integers. The algorithm starts by dividing the larger number by the smaller number and taking the remainder. The smaller number is then

replaced with the remainder, and the process is repeated until the remainder is zero. The last non-zero remainder is the greatest common divisor of the original two numbers.

3 Chairux Algorithm

Definition: Let N be a divisor and given by $N = 10x + y$. Then \exists a pair of integers (α, β) s.t $\alpha x + \beta y = N, \forall \alpha, \beta \in \mathbb{Z}$ and $|\alpha| < 10$. The integers (α, β) are known as **Chairux coefficient for (x, y)** .

Theorem 3.1. Let $N = 10x + y \in \mathbb{N}$.

if $\alpha = 10 - y$ and $\beta = x + 1$, then

$$\alpha x + \beta y = N$$

Proof. $\alpha x + \beta y = (10 - y)x + (x + 1)y$

$$= 10x - yx + yx + y$$

$$= 10x + y$$

□

A function f_N is known as **chairux identity function** if:

$$f_N(\alpha, \beta) = N.$$

chairux identity table

Recall $N = 10x + y = \alpha x + \beta y$

$$10x - \alpha x = \beta y - y$$

$$x(10 - \alpha) = y(\beta - 1)$$

$$xy = yx$$



$$\therefore y = 10 - \alpha \text{ and } x = \beta - 1$$

proof 1

$$N = 10x + y$$

$$N = 10(\beta - 1) + 10 - \alpha$$

$$N = 10\beta - 10 + 10 - \alpha$$

$$\therefore N = 10\beta - \alpha$$

proof 2

$$N = \alpha x + \beta y$$

$$N = \alpha(\beta - 1) + \beta(10 - \alpha)$$

$$N = \alpha\beta - \alpha + 10\beta - \beta\alpha$$

$$\therefore N = 10\beta - \alpha$$

Table 1. chairux identity table

(α, β)	1	2	3	4	5	6	7	8	9	10
1	09	19	29	39	49	59	69	79	89	99
2	08	18	28	38	48	58	68	78	88	98
3	07	17	27	37	47	57	67	77	87	97
4	06	16	26	36	46	56	66	76	86	96
5	05	15	25	35	45	55	65	75	85	95
6	04	14	24	34	44	54	64	74	84	94
7	03	13	23	33	43	53	63	73	83	93
8	02	12	22	32	42	52	62	72	82	92
9	01	11	21	31	41	51	61	71	81	91
10	00	10	20	30	40	50	60	70	80	90

3.1 chairux minimal iteration

Defination: Let N be a divisor and given by $N = 10x + y$, then chairux minimal iteration coefficient for N happen when then number of steps of testing divisibility is minimal.

Chairux minimum iteration is given by:

$$(\alpha, \beta) = \begin{cases} [(10 - y), (1 + x)] & : y \geq 5 \\ [(y^2 - 6y + 10), (6x - xy + 1)] & : y \leq 5 \end{cases}$$

Proof. recall

1) $\alpha = 10 - ky$ and $\beta = 1 + kx$ for $y \geq 5, k = 1$

$\therefore (\alpha, \beta) = [(10 - y), (1 + x)]$ for $y \geq 5$

2) for $y \leq 5, k = (6 - y)$

i $\alpha = 10 - (6 - y)y$ and $\beta = 1 + (6 - y)x$

$$\alpha = 10 - 6y + y^2$$

$$\alpha = y^2 - 6y + 10 \dots *$$

ii $\beta = 1 + (6 - y)x$

$$\beta = 1 + 6x - xy$$

$$\beta = 6x - xy + 1$$

$$\therefore (\alpha, \beta) = [(y^2 - 6y + 10), (6x - xy + 1)] \quad \square$$

Example

find chairux minimum coefficient of the following number.

a) 14 c) 17

b) 23 d) 54

Solution

a) 14

$$\text{Recall } N = 10x + y$$

$$14 = 10 \times 1 + 4$$

$$y = 4 < 5$$

$$\text{since } y < 5$$

$$\alpha = y^2 - 6y + 10 = 4^2 - 6 \times 4 + 10 = \mathbf{2}$$

$$\beta = 6x - xy + 1 = 6 \times 1 - 1 \times 4 + 1 = \mathbf{3}$$

$$\therefore (\alpha, \beta) = \mathbf{(2,3)}$$

$$\mathbf{\text{Proof } } N = \alpha x + \beta y = 2 \times 1 + 3 \times 4 = 14$$

b) 23

$$N = 10x + y$$

$$23 = 10 \times 2 + 3$$

$$y = 3 < 5$$

$$\text{since } y < 5$$

$$\alpha = y^2 - 6y + 10 = 3^2 - 6 \times 3 + 10 = \mathbf{1}$$

$$\beta = 6x - xy + 1 = 6 \times 2 - 2 \times 3 + 1 = \mathbf{7}$$

$$\therefore (\alpha, \beta) = (1, 7)$$

Proof $N = \alpha x + \beta y = 1 \times 2 + 7 \times 3 = 23$

c) 17

$$N = 10x + y$$

$$17 = 10 \times 1 + 7$$

$$y = 7 > 5$$

since $y > 5$

$$\alpha = 10 - y = 10 - 7 = 3$$

$$\beta = x + 1 = 1 + 1 = 2$$

$$\therefore (\alpha, \beta) = (3, 2)$$

Proof $N = \alpha x + \beta y = 3 \times 1 + 2 \times 7 = 17$

3.2 Properties of Chairux Identity

1) $f_N(\alpha_a, \beta_a) \pm f_N(\alpha_b, \beta_b) = f_N[(\alpha_a \pm \alpha_b), (\beta_a \pm \beta_b)]$.

2) $k f_N(\alpha, \beta) = f_N(k\alpha, k\beta)$.

3) $f_N(\alpha, -\beta) = -f_N(-\alpha, \beta) = |f_N(-\alpha, \beta)|$.

Proof. suppose $A = (\alpha_a, \beta_a)$ and $B = (\alpha_b, \beta_b)$ s.t $A, B \in N$

then

$$f_N(\alpha_a, \beta_a) + f_N(\alpha_b, \beta_b) = \lambda N \quad \forall \lambda \in \mathbb{Z}$$

Recall $f_N(\alpha, \beta) = \alpha x + \beta y = N$

$$f_N(\alpha_a, \beta_a) = \alpha_a x + \beta_a y = N \dots *$$

$$f_N(\alpha_b, \beta_b) = \alpha_b x + \beta_b y = N \dots **$$

by adding eqn * and ** we have:

$$f_N(\alpha_a, \beta_a) + f_N(\alpha_b, \beta_b) = (\alpha_a x + \beta_a y) + (\alpha_b x + \beta_b y)$$

$$f_N(\alpha_a, \beta_a) + f_N(\alpha_b, \beta_b) = (\alpha_a x + \alpha_b x) + (\beta_a y + \beta_b y)$$

$$f_N(\alpha_a, \beta_a) + f_N(\alpha_b, \beta_b) = (\alpha_a + \alpha_b)x + (\beta_a + \beta_b)y \cdots ***$$

recall $f_N(\alpha, \beta) = \alpha x + \beta y$

$$\text{let } (\alpha_a + \alpha_b) = \alpha_m \text{ and } (\beta_a + \beta_b) = \beta_m$$

then eqn *** become:

$$f_N(\alpha_a, \beta_a) + f_N(\alpha_b, \beta_b) = \alpha_m x + \beta_m y$$

$$\alpha_m x + \beta_m y = f_N(\alpha_m, \beta_m)$$

$$\therefore f_N(\alpha_m, \beta_m) = \alpha_m x + \beta_m y = N$$

□

Different chairux coefficient may have the same divisor.

Principle of convergence depend on α , that is as $\alpha \rightarrow 0$ chairux coefficient converge faster to N.

1) **alpha**

$$\alpha_0 = 2^0 \alpha_0 - (2^0 - 1)10 \implies 2^0 \alpha_0 - 10 \times 2^0 + 10$$

$$\alpha_1 = 2^1 \alpha_0 - (2^1 - 1)10 \implies 2^1 \alpha_0 - 10 \times 2^1 + 10$$

$$\alpha_2 = 2^2 \alpha_0 - (2^2 - 1)10 \implies 2^2 \alpha_0 - 10 \times 2^2 + 10$$

$$\alpha_3 = 2^3 \alpha_0 - (2^3 - 1)10 \implies 2^3 \alpha_0 - 10 \times 2^3 + 10$$

$$\alpha_n = 2^n \alpha_0 - (2^n - 1)10 \implies 2^n \alpha_0 - 10 \times 2^n + 10$$

$$\therefore \alpha_n \implies 2^n(\alpha_0 - 10) + 10$$

2) **beta**

$$\beta_0 = 2^0 \beta_0 - (2^0 - 1) \implies 2^0 \beta_0 - 2^0 + 1$$

$$\beta_1 = 2^1 \beta_0 - (2^1 - 1) \implies 2^1 \beta_0 - 2^1 + 1$$

$$\beta_2 = 2^2 \beta_0 - (2^2 - 1) \implies 2^2 \beta_0 - 2^2 + 1$$

$$\beta_3 = 2^3 \beta_0 - (2^3 - 1) \implies 2^3 \beta_0 - 2^3 + 1$$

$$\beta_n = 2^n \beta_0 - (2^n - 1) \implies 2^n \beta_0 - 2^n + 1$$

$$\therefore \beta_n \implies 2^n (\beta_0 - 1) + 1$$

In general $(\alpha_n, \beta_n) = [(2^n(\alpha_0 - 10) + 10), (2^n(\beta_0 - 1) + 1)] \forall n \in \mathbb{Z}$

Table 2. Chairux coefficient with same identity

(α_0, β_0)	(α_1, β_1)
(6,2)	(2,3)
(6,3)	(2,5)
(6,4)	(2,7)
(6,5)	(2,9)
(7,2)	(4,3)
(7,3)	(4,5)
(7,4)	(4,7)
(7,5)	(4,9)
(8,2)	(2,5)
(8,3)	(2,9)
(8,4)	(2,13)
(8,5)	(2,17)
(9,2)	(2,9)
(9,3)	(2,17)
(9,4)	(2,25)

definition Let a_0 be the tested value for chairux coefficient (α, β) . Then a function f , is known as **chairux iteration function of N**, if $f_{a_n}(\alpha, \beta) = f_{a_m}(\alpha, \beta) \forall n \neq m$.

definition 2: let $f_{a_0}(\alpha, \beta), f_{a_1}(\alpha, \beta), f_{a_2}(\alpha, \beta), \dots, f_{a_n}(\alpha, \beta), f_{a_{n+1}}(\alpha, \beta) \dots, f_{a_{n+k}}(\alpha, \beta), f_{a_1}(\alpha, \beta) \dots$

Since $f_{a_n}(\alpha, \beta) = f_{a_{n+k}}(\alpha, \beta)$ then k is known as Chairux iteration period for (α, β)

illustration

Let $a_0 = 10x_0 + y_0$

then $f_{a_0}(\alpha, \beta) = \alpha x_0 + \beta y_0 = a_1$

$a_i = 10x_i + y_i$

$\therefore f_{a_i}(\alpha, \beta) = \alpha x_i + \beta y_i = a_{i+1} \forall i = 0, 1, 2 \dots$

Chairux iteration equation

Let $a_0 = 10x_0 + y_0$

$a_1 = \alpha x_0 + \beta y_0$

$\alpha x_i + \beta y_i = a_{i+1}$

$a_1 = \alpha \left(\frac{a_0 - y_0}{10} \right) + \beta y_0 \implies \left(\frac{\alpha(a_0 - y_0) + 10\beta y_0}{10} \right)$

$a_2 = \alpha \left(\frac{a_1 - y_1}{10} \right) + \beta y_1 \implies \left(\frac{\alpha(a_1 - y_1) + 10\beta y_1}{10} \right)$

$a_3 = \alpha \left(\frac{a_2 - y_2}{10} \right) + \beta y_2 \implies \left(\frac{\alpha(a_2 - y_2) + 10\beta y_2}{10} \right)$

$\therefore a_{n+1} = \alpha \left(\frac{a_n - y_n}{10} \right) + \beta y_n \implies \left(\frac{\alpha(a_n - y_n) + 10\beta y_n}{10} \right)$

Example

find chairux iteration of 4270893 for (1,7):

solution

$f_{4270893}(1, 7) = 427089 \times 1 + 7 \times 3 = 427110$

$f_{427110}(1, 7) = 42711 \times 1 + 7 \times 0 = 42711$

$f_{42711}(1, 7) = 4271 \times 1 + 7 \times 1 = 4278$

$f_{4278}(1, 7) = 427 \times 1 + 7 \times 8 = 483$

$f_{483}(1, 7) = 48 \times 1 + 7 \times 3 = 69$

$f_{69}(1, 7) = 6 \times 1 + 7 \times 9 = 69$

Defination: The is inverse of a given number for chairux coefficient (α, β) and given by:

$[f_{n_0}(\alpha, \beta)]^{-1} = \sum_{k=0}^{n-1} 10^k y_k$ is known as **inversion equation of order one**.

illustration

$a_1 = \alpha \left(\frac{a_0 - y_0}{10} \right) + \beta y_0 \implies \left(\frac{\alpha(a_0 - y_0) + 10\beta y_0}{10} \right)$

$\therefore a_1 \implies \frac{\alpha^0}{10^1} [\alpha a_0 + y_0(10\beta - \alpha)]$

$a_2 \implies \left(\frac{\alpha(a_1 - y_1) + 10\beta y_1}{10} \right) \implies \left(\frac{\alpha \left(\frac{\alpha(a_0 - y_0) + 10\beta y_0}{10} - y_1 \right) + 10\beta y_1}{10} \right)$

$$a_2 \implies \frac{1}{10} [\alpha (\frac{\alpha a_0 - \alpha y_0 + 10\beta y_0}{10}) - \alpha y_1 + 10\beta y_1]$$

$$a_2 \implies [\alpha \frac{\alpha a_0 - \alpha y_0 + 10\beta y_0 - 10\alpha y_1 + 100\beta y_1}{10^2}]$$

$$\therefore a_2 \implies \frac{\alpha^1}{10^2} [\alpha a_0 + (10\beta - \alpha)(y_0 + 10y_1)]$$

$$\therefore a_3 \implies \frac{\alpha^2}{10^3} [\alpha a_0 + (10\beta - \alpha)(y_0 + 10y_1 + 100y_2)]$$

$$\therefore a_n \implies \frac{\alpha^{n-1}}{10^n} [\alpha a_0 + (10\beta - \alpha)(y_0 + 10y_1 + 100y_2 + \dots + 10^{n-1}y_{n-1})]$$

in general chairux inversionl equation is given by:

$$a_n = \frac{\alpha^{n-1}}{10^n} [\alpha a_0 + (10\beta - \alpha) \sum_{k=0}^{n-1} 10^k y_k]$$

let $(10\beta - \alpha)$ be γ then:

$$a_n = \frac{\alpha^{n-1}}{10^n} [\alpha a_0 + \gamma \sum_{k=0}^{n-1} 10^k y_k]$$

$$\therefore \sum_{k=0}^{n-1} 10^k y_k = \frac{\frac{10^n \times a_n}{\alpha^{n-1}} - \alpha a_0}{\gamma}$$

Example

find chairux inversion of $f_{4270893}(1, 7)$.

Solution

Chairux inversion is given by:

$$\sum_{k=0}^{n-1} 10^k y_k = \frac{\frac{10^n \times a_n}{\alpha^{n-1}} - \alpha a_0}{\gamma}$$

$$\gamma = 10\beta - \alpha$$

$$\gamma = 10 \times 7 - 1 = 69$$

$$\frac{\gamma}{k} \leq N \leq \gamma \quad \forall k = 1, 2, 3, \dots$$

suppose $k=1$

$$N=69$$

$$[f_{4270893}(1, 7)]^{-1} = \frac{10^5 \times 69 - 4270893}{69} = 38, 103$$

$$[f_{4270893}(1, 7)]^{-1} = \frac{10^6 \times 69 - 4270893}{69} = 938, 103$$

$$[f_{4270893}(1, 7)]^{-1} = \frac{10^7 \times 69 - 4270893}{69} = 9, 938, 103$$

$$[f_{4270893}(1, 7)]^{-1} = \frac{10^{10} \times 69 - 4270893}{69} = 9, 999, 938, 103$$


in general

$$[f_{4270893}(1, 7)]^{-1} = \frac{10^t \times 69 - 4270893}{69} = 9_{t-5}38, 103$$

∴ chairux inversion=38, 103

The graph of y_n against n is known [Chairux inversion graph](#).

Fig 1. **chairux inversion graph of $f_{4270893}(1, 7)$**



3.3 Chairux period length (λ_l)

Let $f_{a_i}(\alpha, \beta) = a_{i+1}$ be an iteration function for $i=0,1,2,3,\dots n,\dots m,\dots$ if

$f_{a_n}(\alpha, \beta) = f_{a_m}(\alpha, \beta)$ then:

$|m - n|$ is known as [the period length of \$\(\alpha, \beta\)\$](#) and denoted as:

$$\lambda_l = |m - n|$$

Table 3. Chairux period length (λ_l) table

(α, β)	1	2	3	4	5	6	7	8	9	10
1	01	18	28	06	42	58	22	13	44	02
2	01	06	06	09	04	14	16	04	05	42
3	01	08	01	18	46	09	66	10	14	96
4	01	01	03	03	22	02	10	06	13	01
5	01	04	20	06	06	10	14	20	08	18
6	01	03	02	04	05	01	04	12	03	44
7	01	10	11	05	42	26	03	03	41	15
8	01	02	05	01	06	12	11	02	04	22
9	01	06	06	05	20	01	60	35	01	06

Theorem 3.2. *let $|m - n|$ be chairux period length and a_0 be the tested value of N . Then a_0 is divisible by N if and only if the period length is one.*

i.e $\lambda_l = |m - n| = 1$

Not divisible by N if:

$|m - n| > 1$

Example

1) Find wheather the following are divisible by 37.

a) 2181964 c) 5881964

b) 236754321 d) 95421

2) Find wheather the following are divisible by 23.

a) 42348903 c) 7234597

b) 28395041 d) 181574903

3) Find wheather the following are divisible by 84.

a) 81964 c) 4786740

b) 626547684 d) 2157894

Solution

1.a) 37

$N = 10x + y$

$$37 = 10 \times 3 + 7$$

$$y = 7 > 5$$

since $y > 5$

$$\alpha = 10 - y = 10 - 7 = \mathbf{3}$$

$$\beta = x + 1 = 3 + 1 = \mathbf{4}$$

$$\therefore (\alpha, \beta) = \mathbf{(3,4)}$$

$$f_{2181964}(3, 4) = 3 \times 218196 + 4 \times 4 = 654604$$

$$f_{654604}(3, 4) = 3 \times 65460 + 4 \times 4 = 196396$$

$$f_{196396}(3, 4) = 3 \times 19639 + 4 \times 6 = 58941$$

$$f_{58941}(3, 4) = 3 \times 5894 + 4 \times 1 = 17686$$

$$f_{17686}(3, 4) = 3 \times 1768 + 4 \times 6 = 5328$$

$$f_{5328}(3, 4) = 3 \times 532 + 4 \times 8 = 1628$$

$$f_{1628}(3, 4) = 3 \times 162 + 4 \times 8 = 518$$

$$f_{518}(3, 4) = 3 \times 51 + 4 \times 8 = 185$$

$$f_{185}(3, 4) = 3 \times 18 + 4 \times 5 = 74$$

$$f_{74}(3, 4) = 3 \times 7 + 4 \times 4 = 37$$

$$f_{37}(3, 4) = 3 \times 3 + 4 \times 7 = 37$$

since $f_{74}(3, 4) = f_{37}(3, 4) = 37$

$\therefore 2181964$ is divisible by 37.

1.c)

$$N = 10x + y$$

$$37 = 10 \times 3 + 7$$

$$y = 7 > 5$$

since $y > 5$

$$\alpha = 10 - y = 10 - 7 = \mathbf{3}$$

$$\beta = x + 1 = 3 + 1 = \mathbf{4}$$

$$\therefore (\alpha, \beta) = (\mathbf{3,4})$$

$$f_{5881964}(3, 4) = 3 \times 588196 + 4 \times 4 = 1764604$$

$$f_{1764604}(3, 4) = 3 \times 176460 + 4 \times 4 = 529396$$

$$f_{529396}(3, 4) = 3 \times 52939 + 4 \times 6 = 158841$$

$$f_{158841}(3, 4) = 3 \times 15884 + 4 \times 1 = 47656$$

$$f_{47656}(3, 4) = 3 \times 4765 + 4 \times 6 = 14319$$

$$f_{14319}(3, 4) = 3 \times 1431 + 4 \times 9 = 4329$$

$$f_{4329}(3, 4) = 3 \times 432 + 4 \times 9 = 1332$$

$$f_{1332}(3, 4) = 3 \times 133 + 4 \times 2 = 407$$

$$f_{407}(3, 4) = 3 \times 40 + 4 \times 7 = 148$$

$$f_{148}(3, 4) = 3 \times 14 + 4 \times 8 = 74$$

$$f_{74}(3, 4) = 3 \times 7 + 4 \times 4 = 37$$

$$\text{since } f_{74}(3, 4) = f_{37}(3, 4) = 37$$

$\therefore 5881964$ is divisible by 37.

3.4 Chairux iteration bound

Let $f_N(\alpha, \beta)$ be chairux function of N . Then chairux iteration bound is given by:

$$a_i = N \pm k \quad \forall 1 \leq k < N$$

chairux iteration is either upper or lower

Chairux iteration bound graph


Let $N = 10x + y$ and $a_{i+1} = \alpha x_i + \beta y_i$. Then Chairux differential iteration is given by:

$$\Delta_i = \frac{a_{i+1} - a_i}{y} \quad \forall i=0,1,2,\dots$$

A graph of Δ_i against i is known as chairux iteration graph.

The graph of chairux iteration below zero is known as lower bound and above is known as upper bound graph.

Fig 2. Chairux iteration bound graph of $f_{22,23}$ and $24(1,7)$



3.5 Chairux divisor N for (α, β)

This are various method of finding N given chairux coefficient (α, β) . This methods include:

- 1) Structural method

This is the simplest method of finding N given Chairux coefficient. this method is also known as inverse method.

Recall

$$\alpha = 10 - ky, \beta = 1 + kx \text{ and } N = 10x + y$$

$$ky = 10 - \alpha \implies y = \frac{10 - \alpha}{k}$$

$$kx = \beta - 1 \implies x = \frac{\beta - 1}{k}$$

But we know: $N = 10x + y$

$$N = 10\left(\frac{\beta - 1}{k}\right) + \left(\frac{10 - \alpha}{k}\right)$$

$$N = \frac{1}{k}(10\beta - 10 + 10 - \alpha)$$

$$N = \frac{1}{k}(10\beta - \alpha)$$

k is equal to gcd $[(10\beta - \alpha), y]$

y is the last digit of $(10\beta - \alpha)$

$$\therefore N_{min} = \frac{(10\beta - \alpha)}{\gcd[(10\beta - \alpha), y]} \text{ of } (\alpha, \beta)$$

2) Iteration method

Let a_m be the inverse of a_n and $f_{a_m}(\alpha, \beta) = f_{a_n}(\alpha, \beta)$ s.t $a_m \neq a_n$.

$$\therefore N_{min} = \frac{(|a_m - a_n|)}{\gcd[(|a_m - a_n|), y]} \text{ of } (\alpha, \beta)$$

Example

Given that $a_m = 36$ for chairux coefficient (4, 5). Calculate chairux divisor N.

Solution

$$a_n = (a_m)^{-1} \text{ s.t } a_m \neq a_n$$

$$f_{m+1}(4, 5) = f_{36}(4, 5) = 3 \times 4 + 5 \times 6 = 42$$

$$4x + 5y = 42$$

$$4x = 42 - 5y$$

$$x = \frac{42 - 5y}{4}$$

x is a whole nuber when y is either 2 or 6

i) when y=2

$$x = \frac{42 - 5 \times 2}{4} = 8$$

$$a_n = 10x + y = 82$$

ii) when y=6

$$x = \frac{42 - 5 \times 6}{4} = 3$$

$$a_n = 10x + y = 36$$

since $a_m \neq a_n$

$$\therefore a_n = 82$$

$$36=(82)^{-1}$$

$$f_{82}(4, 5) = f_{36}(4, 5) = 42$$

$$|a_m - a_n|=|36 - 82|=46$$

$$\therefore N_{min} = \frac{f_{46}}{gcd(46,6)} = \frac{46}{2} = 23$$

3) Bi-section method

Let a_0 and b_0 be 2 ivp s.t $b_0 = ka_0$ for $k > 1$, If the function $f_{a_n}(\alpha, \beta) > f_{b_n}(\alpha, \beta)$, Then Chairux divisor N is given by:

$$N = ka_n - b_n$$

chairux minimum divisor is given by;

(α, β)	(α, β)
$a_0 = \Omega$	$b_0 = k\Omega$
$f_{a_0}(\alpha, \beta) = a_1$	$f_{b_0}(\alpha, \beta) = b_1$
$f_{a_1}(\alpha, \beta) = a_2$	$f_{b_1}(\alpha, \beta) = b_2$
\vdots	\vdots
$f_{a_{n-1}}(\alpha, \beta) = a_n$	$f_{b_{n-1}}(\alpha, \beta) = b_n$

$$\therefore N_{min} = \frac{(ka_n - b_n)}{gcd[(ka_n - b_n), y]} \text{ of } (\alpha, \beta)$$

Example

Using bisection method, find chairux minimum divisor (N_{min}) given chairux coefficient (3, 6).

Solution

(3, 6)	(3, 6)
$a_0 = 10$	$b_0 = 20$
$a_1 = 3 \times 1 + 0 \times 6 = 03$	$b_1 = 3 \times 2 + 0 \times 6 = 06$
$a_2 = 3 \times 0 + 3 \times 6 = 18$	$b_2 = 3 \times 0 + 6 \times 6 = 36$
$a_3 = 3 \times 1 + 8 \times 6 = 51$	$b_3 = 3 \times 3 + 6 \times 6 = 45$
$a_4 = 3 \times 5 + 1 \times 6 = 21$	$b_4 = 3 \times 4 + 5 \times 6 = 42$

since $a_3 > b_3$

$$\gamma = 2a_3 - b_3$$

$$\gamma = 2 \times 51 - 45 = 57$$

$$N_{min} = \frac{f_{\gamma}(3,6)}{gcd[f_{\gamma}(3,6), y]} \implies \frac{f_{57}(3,6)}{gcd[f_{57}(3,6), 7]} = \frac{57}{gcd(57, 7)} = 57$$

chairux order

Let (a,b) be a given pair of Chairux coefficient, Then **Chairux order** denoted as $\ll N \gg$ is given by:

$$\ll N \gg = \gamma^\theta \gcd\left(\frac{(a,b)}{[\gamma^\theta, (10-y)]}\right)$$

$$\gamma^\theta = \frac{\gamma}{\gcd[\gamma, (10-y)]}$$

$$\gamma = N = ka_n - b_n$$

Theorem 3.3. Let (a,b) be a given pair of chairux coefficient s.t $b > a$. if the function $f_{a_n}(\alpha, \beta) > f_{b_n}(\alpha, \beta)$. Then Chairux order denoted as $\ll N \gg$ is given by:

$$\ll N \gg = \frac{a_n \times b_0 - b_n \times a_0}{\gcd[,y]}$$

3.6 Chairux polynomial mean

The average mean of Chairux identity denoted as $P_n(k)$, where n and k are the rate and order of chairux identity respectively is known as **Chairux polynomial mean**.

Theorem 3.4. Every chairux identity with the same factor, have the same polynomial mean.

Proof. since all divisor having the common multiple factor have the same polynomial mean.

Chairux polynomial mean can be illustrated by table.

Table 4. Chairux polynomial mean $P_n(k)$ table

(n, k)	$\alpha = 1$									
	1	2	3	4	k
1	00 ₀ 1	10 ₀ 1	20 ₀ 1	30 ₀ 1	$(k-1)0_01$
2	00 ₁ 1	10 ₁ 1	20 ₁ 1	30 ₁ 1	$(k-1)0_11$
3	00 ₂ 1	10 ₂ 1	20 ₂ 1	30 ₂ 1	$(k-1)0_21$
4	00 ₃ 1	10 ₃ 1	20 ₃ 1	30 ₃ 1	$(k-1)0_31$
⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮
⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮
n	00 _{$n-1$} 1	10 _{$n-1$} 1	20 _{$n-1$} 1	30 _{$n-1$} 1	$(k-1)0_{n-1}1$

$$\alpha = 2$$

(n, k)	1	2	3	4	k
1	01 ₀ 2	11 ₀ 2	21 ₀ 2	31 ₀ 2	$(k-1)1_02$
2	01 ₁ 2	11 ₁ 2	21 ₁ 2	31 ₁ 2	$(k-1)1_12$
3	01 ₂ 2	11 ₂ 2	21 ₂ 2	31 ₂ 2	$(k-1)1_22$
4	01 ₃ 2	11 ₃ 2	21 ₃ 2	31 ₃ 2	$(k-1)1_32$
⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮
⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮
n	01 _{n-1} 2	11 _{n-1} 2	21 _{n-1} 2	31 _{n-1} 2	$(k-1)1_{n-1}2$

⋮

$$\alpha = \Omega$$

(n, k)	1	2	3	k
1	$0(\Omega-1)_0\Omega$	$1(\Omega-1)_0\Omega$	$2(\Omega-1)_0\Omega$	$(k-1)(\Omega-1)_0\Omega$
2	$0(\Omega-1)_1\Omega$	$1(\Omega-1)_1\Omega$	$2(\Omega-1)_1\Omega$	$(k-1)(\Omega-1)_1\Omega$
3	$0(\Omega-1)_2\Omega$	$1(\Omega-1)_2\Omega$	$2(\Omega-1)_2\Omega$	$(k-1)(\Omega-1)_2\Omega$
4	$0(\Omega-1)_3\Omega$	$1(\Omega-1)_3\Omega$	$2(\Omega-1)_3\Omega$	$(k-1)(\Omega-1)_3\Omega$
⋮	⋮	⋮	⋮	⋮	⋮	⋮
⋮	⋮	⋮	⋮	⋮	⋮	⋮
n	$0(\Omega-1)_{n-1}\Omega$	$1(\Omega-1)_{n-1}\Omega$	$2(\Omega-1)_{n-1}\Omega$	$(k-1)(\Omega-1)_{n-1}\Omega$

$$\therefore P_n(k) = (k-1)_1(\Omega-1)_{n-1}\Omega_1$$

$$P_n(k) = 10^{1+(n-1)}(k-1)_1 + 10^1(\Omega-1)_{n-1} + \Omega_1$$

$$P_n(k) = 10^n(k-1)_1 + 10^1(\Omega-1)_{n-1} + \Omega_1$$

But $10^n = 1_{n+1} - 1_n$, $10_{n-1} = 1_n - 1$ and $A_n B_m = A_m B_n$, from chairux repeatal statement.

$$P_n(k) = 10^n(k-1)_1 + 10_{n-1}(\Omega-1) + \Omega_1$$

$$P_n(k) = 10^n(k-1)_1 + (1_n - 1)(\Omega-1) + \Omega_1$$

$$P_n(k) = 10^n(k-1)_1 + \Omega_n - 1_n - \Omega_1 + 1 + \Omega_1$$

$$P_n(k) = 10^n(k-1)_1 + (\Omega-1)_n + 1$$

$$P_n(k) = (1_{n+1} - 1_n)(k-1)_1 + (\Omega-1)_n + 1$$

$$P_n(k) = k_{n+1} - 1_{n+1} + \Omega_n - k_n + 1$$

$$P_n(k) = k(10^n + 1_n) - 1_{n+1} - k_n + 1_n + \Omega_n 1_n + 1$$

$$P_n(k) = 10^n k + k_n - k_n + \Omega_n - (1_{n+1} - 1)$$

$$P_n(k) = 10^n k + \Omega_n - 10_n$$

$$\therefore P_n(k) = 10^n k + (\Omega - 10)_n$$

3.7 Properties of Chairux polynomial mean

$$1) P_0(k) = k$$

$$2) P_n(0) = (\Omega - 10)_n$$

$$3) P_n(1) = (\Omega - 10)_n + 1 = P_n(0) + 1$$

$$4) P_1(1) = \Omega$$

$$5) P_1(k) = 10(k - 1) + \Omega$$

$$6) P_n(k) = 10^n k + (\Omega - 10)_n$$

When the rate of chairux polynomial mean countinuously increase by one unit is known as [chairux unitizol mean](#).

Recall:

$$P_n(k) = 10^n k + (\Omega - 10)_n \dots *$$

$$P_{n+1}(k) = 10^{n+1} k + (\Omega - 10)_{n+1} = 10^n \cdot 10k + (\Omega - 10) \cdot 1_{n+1}$$

$$\text{but: } 1_{n+1} = 10^n + 1_n$$

$$P_{n+1}(k) = 10^n \cdot 10k + (\Omega - 10)(10^n + 1_n) = 10^n \cdot 10k + 10^n(\Omega - 10)_n$$

$$P_{n+1}(k) = 10^n [10k + (\Omega - 10)_1] + (\Omega - 10)_n$$

$$P_{n+2}(k) = 10^{n+1} [10k + (\Omega - 10)_1] + (\Omega - 10)_{n+1}$$

$$P_{n+2}(k) = 10^n \cdot 10 [10k + (\Omega - 10)_1] + 10^n \cdot (\Omega - 10)_1 + (\Omega - 10)_n$$

$$P_{n+2}(k) = 10^n [[10^2 k + 10(\Omega - 10)_1] + (\Omega - 10)_1] + (\Omega - 10)_n$$

$$P_{n+2}(k) = 10^n [10^2 k + 10(\Omega - 10)_1] + (\Omega - 10)_1 + (\Omega - 10)_n$$

$$P_{n+2}(k) = 10^n [10^2 k + (\Omega - 10)_2] + (\Omega - 10)_n$$

$$P_{n+3}(k) = 10^{n+1} [10^2 k + (\Omega - 10)_2] + (\Omega - 10)_{n+1}$$

$$P_{n+3}(k) = 10^n \cdot 10 [10^2 k + (\Omega - 10)_2] + 10^n \cdot (\Omega - 10)_1 + (\Omega - 10)_n$$

$$P_{n+3}(k) = 10^n [[10^3 k + 10(\Omega - 10)_2] + (\Omega - 10)_1] + (\Omega - 10)_n$$

$$P_{n+3}(k) = 10^n [10^3 k + 10(\Omega - 10)_2] + (\Omega - 10)_1 + (\Omega - 10)_n$$

$$P_{n+3}(k) = 10^n [10^3 k + (\Omega - 10)_3] + (\Omega - 10)_n$$

Given that n is the rate, k is the order and b is the unitizol constant then Chairux unitizol mean is given by:

$$\therefore P_{n+b}(k) = 10^n [10^b k + (\Omega - 10)_b] + (\Omega - 10)_n$$

Let 'b' be negative

$$P_{n-b}(k) = 10^n [10^{-b} k + (\Omega - 10)_{-b}] + (\Omega - 10)_n$$

Remark

$$X_{-\rho} = \frac{-X_\rho}{10^\rho}$$

$$(\alpha - 10)_{-\rho} = \frac{-(\alpha - 10)_\rho}{10^\rho}$$

$$P_{n-b}(k) = 10^n \left[\frac{k}{10^b} - \frac{(\Omega - 10)_b}{10^b} \right] + (\Omega - 10)_n$$

$$P_{n-b}(k) = \frac{10^n}{10^b} [k - (\Omega - 10)_b] + (\Omega - 10)_n$$

$$\therefore P_{n-b}(k) = 10^{n-b} [k - (\Omega - 10)_b] + (\Omega - 10)_n$$

Theorem 3.5. *let X be a universal set $\forall a, b \in X$ and let $Y = a + b$ then;*

$$(10^X - 1)_Y = (10^Y - 1)_X$$

Proof. we know that $X = a^c + a = b^c + b$ and $Y = a + b$

$$(10^m)_n + 1_m = 1_{m+n}$$

$$(10^{(a+b)^c})_{a+b} + 1_{(a+b)^c} = 1_{(a+b)^c + (a+b)} = 1_X$$

but $(a + b)^c = a^c - b$

$$(10^{(a+b)^c})_{a+b} = (10^{a^c-b})_{a+b} = \frac{(10^{a^c})_{a+b}}{10^b}$$

we know that: $1_{(a+b)^c} = 1_{a^c-b} = \frac{1_{a^c}-1_b}{10^b}$

$$\frac{(10^{a^c})_{a+b}}{10^b} + \frac{1_{a^c}-1_b}{10^b} = 1_X$$

$$(10^{a^c})_{a+b} + 1_{a^c} - 1_b = 10^b_X$$

$$(10^{a^c})_Y + 1_{a^c} - 1_b = 10^b_X$$

$$(10^{X-a})_Y + 1_{X-a} - 1_b = 10^b_X$$

$$\frac{(10^X)_Y}{10^a} + \frac{1_X-1_a}{10^a} = (10^b)_X + 1_b$$

$$(10^X)_Y + 1_X - 1_a = 10^a(10^b)_X + (10^a)_b$$

$$(10^X)_Y + 1_X = 10^a(10^b)_X + (10^a)_b + 1_a$$

$$(10^X)_Y + 1_X = (10^{a+b})_X + (10^a)_b + 1_a$$

$$(10^X)_Y + 1_X = (10^Y)_X + (10^a)_b + 1_a$$

$$(10^X)_Y + 1_X = (10^Y)_X + 1_{a+b}$$

$$(10^X)_Y + 1_X = (10^Y)_X + 1_Y$$

$$(10^X)_Y - 1_Y = (10^Y)_X - 1_X$$

$$(10^X - 1)_Y = (10^Y - 1)_X \forall X, Y \in \mathbb{C}$$

□

Theorem 3.6. let X be a universal set $\forall a, b \in X$ and let $Y = a - b$ then;

$$(10^X - 1)_Y = (10^Y - 1)_X$$

Proof. we know that $X = a^c + a = b^c + b$ and $Y = a - b$

$$(10^m)_n + 1_m = 1_{m+n}$$

$$(10^{(a-b)^c})_{a-b} + 1_{(a-b)^c} = 1_{(a-b)^c+(a-b)} = 1_X$$

but $(a - b)^c = a^c + b$

$$(10^{a^c+b})_{a-b} + 1_{a^c+b} = 1_X$$

$$10^b(10^{a^c})_{a-b} + (10^b)_{a^c} + 1_b = 1_X$$

$$10^b[(10^{a^c})_{a-b} + 1_{a^c}] = 1_X - 1_b$$

$$10^b[(10^{X-a})_{a-b} + 1_{X-a}] = 1_X - 1_b$$

$$10^b\left[\frac{(10^X)_{a-b}}{10^a} + \frac{1_{X-1_a}}{10^a}\right] = 1_X + 1_b$$

$$10^{b-a}[(10^{X-a})_{a-b} + 1_{X-a}] = 1_X - 1_b$$

$$10^{-Y}[(10^{X-a})_{a-b} + 1_{X-a}] = 1_X - 1_b$$

$$[(10^{X-a})_{a-b} + 1_{X-a}] = (10^Y)_X - (10^Y)_b$$

$$1_a = 1_{Y+b} \implies (10^Y)_b + 1_Y$$

$$(10^X)_Y + 1_X - ((10^Y)_b + 1_Y) = (10^Y)_X - (10^Y)_b$$

$$(10^X)_Y - 1_Y = (10^Y)_X - 1_X$$

$$(10^X - 1)_Y = (10^Y - 1)_X \forall X, Y \in \mathbb{C}$$

□

3.8 chairux reptal function

This is a technique involving arrangement of common series of an integers from statement to column then to row.

The number of times the main repeat itself is known as [Reptal number](#)

Statement which give the main data 'X' and the number of time's it repeat itself (n) is known as [reptal statement](#) and denoted as (X_n) .

a) Chairux column: This is arrangement of reptal statement in a vertical order.

$$X_n = \begin{pmatrix} X \\ X \\ \vdots \\ X \end{pmatrix}$$

b) Chairux row: This is arrangement of reptal statement in a horizontal order.

$$X_n = X \left(\sum_{k=1}^n 10^{k-1} \right) = X f_{10}(n) = X (1 \ 1 \ 1 \ \dots \ 1)$$

Properties of Repeatal Statement

1) $X_n = 10^{n-b}X_b + X_{n-b}$

2) $X_0 = 0$

3) $X_1 = X$

4) $X_n \pm Y_n = (X \pm Y)_n$

5) $X_n \pm Y_m = X_{n-m}10^m + (X \pm Y)_m$

6) $X_n \times Y_m = X_m \times Y_n = (XY)_{n|m} = (XY)_{m|n}$

7) $X_{-b} = -\frac{X_b}{10^{-b}}$

Proof. $X_n = 10^{n-b}X_b + X_{n-b}$

let $b = n$ then:

$$X_n = 10^{n-n}X_n + X_{n-n}$$

$$X_n = 10^0X_n + X_0$$

$$X_0 = X_n - X_n$$

$$X_0 = 0$$

□

Proof. let $n = 0$ then:

$$X_0 = 10^{-b}X_b + X_{-b}$$

but $X_0 = 0$

$$10^{-b}X_b + X_{-b} = 0$$

$$X_{-b} = -10^{-b}X_b$$

$$X_{-b} = -\frac{X_b}{10^{-b}}$$

□

Proof. Recall: $X_n = 10^{n-b}X_b + X_{n-b}$

let $n = a + b$ then:

$$X_{a+b} = 10^aX_b + X_a$$

let $a = a - b$ then:

$$X_{a-b+b} = 10^{a-b}X_b + X_{a-b}$$

$$X_a = 10^{a-b}X_b + X_{a-b}$$

$$X_{a+b} = 10^bX_a + X_b = 10^aX_b + X_a$$

$$X_{a+b} = 10^bX_a + X_b$$

□

Example

Use repeatal statement to solve:

a) $7_4 \times 5_3$ b) $5_4 \times 7_3$

Solution

$$\text{a) } (7 \times 5)_{4||3} = (35)_{4||3} = (35)_4 = \begin{pmatrix} 3 & 5 & & & \\ & 3 & 5 & & \\ + & & 3 & 5 & \\ & & & 3 & 5 \\ & & & & 3 & 5 \end{pmatrix} = (38885)_3 = \begin{pmatrix} 3 & 8 & 8 & 8 & 5 & \\ + & 3 & 8 & 8 & 8 & 5 \\ & & 3 & 8 & 8 & 5 \\ & & & 3 & 8 & 5 \\ & & & & 3 & 5 \end{pmatrix} =$$

4316235

$$\text{a) } (5 \times 7)_{3||4} = (35)_{3||4} = (35)_3 = \begin{pmatrix} 3 & 5 & & & \\ + & 3 & 5 & & \\ & & 3 & 5 & \\ & & & 3 & 5 \\ & & & & 3 & 5 \end{pmatrix} = (3885)_4 = \begin{pmatrix} 3 & 8 & 8 & 5 & \\ + & 3 & 8 & 8 & 5 \\ & & 3 & 8 & 5 \\ & & & 3 & 8 & 5 \\ & & & & 3 & 5 \end{pmatrix} =$$

4316235

$$\therefore X_n \times Y_m = X_m \times Y_n = (XY)_{n||m} = (XY)_{m||n}$$

definition: Chairux repeatal function f_{10} is known as **chairux repeatal linear function** if it have more than one rate and given as:

$$f_{10}(a + b + c + \dots)$$

Properties of Repeatal function

1) $f_{10}(0) = 0$

2) $f_{10}(1) = 1$

3) $f_{10}(-n) = \frac{-f_{10}(n)}{10^n}$

4) $f_{10}(a + b) = 10^a f_{10}(b) + f_{10}(a) = 10^b f_{10}(a) + f_{10}(b)$

5) $f_{10}(n) = 1_n = \sum_{k=1}^n 10^{k-1}$

Chairux Repeatal Table

This is a table of chairux function of base 10 and rate less to absolute value of one.

When the rate is more than one we use chairux linearity property.

$$\text{i.e } f_{10}(a + b) = 10^a f_{10}(b) + f_{10}(a) = 10^b f_{10}(a) + f_{10}(b)$$

Example

Use repeatal function of a whole number rate to solve:

$$a) f_{10}(3) \quad b) f_{10}(1) \quad c) f_{10}(-4) \quad d) f_{10}(0)$$

Solution

$$a) f_{10}(3) = 1_3 = 111$$

$$b) f_{10}(1) = 1_1 = 1$$

$$c) f_{10}(-4) = 1_{-4} = \frac{-f_{10}(4)}{10^4} = \frac{-1_4}{10^4} = -0.1111$$

$$d) f_{10}(0) = 1_0 = 0$$

Table 5. Chairux Repeatal ($f_{10}(n)$) Table

n	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	0.0000	0.0026	0.0052	0.0079	0.0107	0.0136	0.0165	0.0194	0.0225	0.0256
0.1	0.0288	0.0321	0.0354	0.0388	0.0423	0.0458	0.0495	0.0533	0.0571	0.0610
0.2	0.0650	0.0691	0.0733	0.0776	0.0820	0.0866	0.0912	0.0959	0.1007	0.1057
0.3	0.1106	0.1158	0.1211	0.1265	0.1320	0.1377	0.1435	0.1495	0.1555	0.1618
0.4	0.1680	0.1745	0.1812	0.1880	0.1950	0.2021	0.2094	0.2169	0.2246	0.2324
0.5	0.2403	0.2485	0.2569	0.2655	0.2743	0.2832	0.2924	0.3019	0.3115	0.3213
0.6	0.3312	0.3415	0.3521	0.3629	0.3739	0.3852	0.3968	0.4086	0.4208	0.4332
0.7	0.4458	0.4588	0.4721	0.4857	0.4996	0.5138	0.5284	0.5433	0.5585	0.5742
0.8	0.5900	0.6063	0.6231	0.6402	0.6577	0.6756	0.6939	0.7127	0.7319	0.7516
0.9	0.7715	0.7921	0.8131	0.8347	0.8567	0.8793	0.9023	0.9260	0.9501	0.9747

Example

1) Use repeatal function of a rate less than one from repeatal table to solve:

$$a) f_{10}(0.3) \quad b) f_{10}(0.71) \quad c) f_{10}(-0.58) \quad d) f_{10}(0.07)$$

2) Use chairux linearity function to show that.

$$a) f_{10}(0.7) = f_{10}(0.3 + 0.4) \quad b) f_{10}(-0.71) = f_{10}(0.29 - 1)$$

3) Use chairux linearity function to solve.

$$a) f_{10}(3.7) \quad b) f_{10}(2.71)$$

Solution

$$1) a) f_{10}(0.3) = 1_{0.3} = 0.1106$$

$$b) f_{10}(0.71) = 1_{0.71} = 0.4588$$

$$c) f_{10}(-0.58) = 1_{-0.58} = \frac{-f_{10}(0.58)}{10^{0.58}} = \frac{-0.3115}{10^{0.58}} = -0.0819$$

$$d) f_{10}(0.07) = 1_{0.07} = 0.0194$$

$$2) a) f_{10}(0.7) = f_{10}(0.3 + 0.4)$$

$$f_{10}(0.7) = 0.4458$$

$$f_{10}(0.3 + 0.4) = 10^{0.3} f_{10}(0.4) + f_{10}(0.3) = 10^{0.3}(0.1680) + 0.1106 = 0.4458$$

$$f_{10}(0.3 + 0.4) = 10^{0.4} f_{10}(0.3) + f_{10}(0.4) = 10^{0.4}(0.1106) + 0.1680 = 0.4458$$

$$\therefore f_{10}(0.7) = f_{10}(0.3 + 0.4)$$

$$b) f_{10}(-0.71) = f_{10}(0.29 - 1)$$

$$f_{10}(-0.71) = \frac{-f_{10}(0.71)}{10^{0.71}} = \frac{-0.4588}{10^{0.71}} = -0.0895$$

$$f_{10}(0.29 - 1) = 10^{0.29} f_{10}(-1) + f_{10}(0.29) = 10^{0.29}(-0.1) + 0.1057 = -0.0893$$

$$f_{10}(0.29 - 1) = 10^{-1} f_{10}(0.29) + f_{10}(-1) = 10^{-1}(0.1057) + -0.1 = -0.0894$$

$$\therefore f_{10}(-0.71) = f_{10}(0.29 - 1)$$

$$3) a) f_{10}(3.7) = f_{10}(3 + 0.7) = f_{10}(4 - 0.3)$$

$$f_{10}(3.7) = f_{10}(3 + 0.7) = 10^3 f_{10}(0.7) + f_{10}(3) = 1000(0.4458) + 111 = 556.8$$

$$f_{10}(4 - 0.3) = 10^{-0.3} f_{10}(4) + f_{10}(-0.3) = 0.5012(1111) + -0.0554 = 556.7778$$

$$\therefore f_{10}(3.7) = 556.8$$

ExampleFind the value of p in $3_p = 1000$ **Solution**

$$3_p = 3(1_p) = 1000$$

$$3(1_p) = 1000$$

$$1_p = \frac{1000}{3} = 333.3\dots$$

$$1_3 \leq 333.33 \leq 1_4$$

method (i)

$$1_{3+k} = 333.33333 \quad s.t \quad 3 + k = p$$

$$1_{3+k} = 10^3(f_{10}(k)) + f_{10}(3) = 1000(f_{10}(k)) + 111 = 333.3\dots$$

$$1_{3+k} = 1000(f_{10}(k)) = 333.3\dots - 111 = 222.3\dots$$

$$1_k = \frac{222.3\dots}{1000} = 0.2223\dots$$

$$k = (1_k)^{-1} = (0.2223\dots)^{-1} = 0.48$$

$$p = 3 + k = 3 + 0.48 = 3.48$$

method (ii)

$$1_{4-k} = 333.33333 \quad s.t \quad 4 - k = p$$

$$1_{4-k} = 10^4(f_{10}(-k)) + f_{10}(4) = 10000(f_{10}(-k)) + 1111 = 333.3\dots$$

$$1_{4-k} = 10000(f_{10}(-k)) = 333.3\dots - 1111 = -777.6\dots$$

$$1_{-k} = \frac{-777.6\dots}{1000} = -0.07776\dots$$

$$k = (1_{-k})^{-1} = (-0.07776\dots)^{-1} = \text{STUNK!!!}$$

Since this is difficult to solve i provide a negative repeatal table.

Table 6. Chairux Repeatal ($f_{10}(-n)$) Table

$-n$.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	0.0000	0.0025	0.0050	0.0074	0.0098	0.0121	0.0144	0.0165	0.0187	0.0208
0.1	0.0229	0.0249	0.0269	0.0288	0.0306	0.0324	0.0342	0.0360	0.0377	0.0394
0.2	0.0410	0.0426	0.0442	0.0457	0.0472	0.0487	0.0501	0.0515	0.0528	0.0542
0.3	0.0554	0.0567	0.0580	0.0592	0.0603	0.0615	0.0626	0.0638	0.0648	0.0659
0.4	0.0669	0.0679	0.0689	0.0698	0.0708	0.0717	0.0726	0.0735	0.0744	0.0752
0.5	0.0760	0.0768	0.0776	0.0784	0.0791	0.0798	0.0805	0.0813	0.0819	0.0826
0.6	0.0832	0.0838	0.0845	0.0851	0.0857	0.0862	0.0868	0.0874	0.0879	0.0884
0.7	0.0889	0.0895	0.0900	0.0904	0.0909	0.0914	0.0918	0.0923	0.0927	0.0931
0.8	0.0935	0.0939	0.0943	0.0947	0.0951	0.0954	0.0958	0.0961	0.0965	0.0968
0.9	0.0971	0.0974	0.0978	0.0981	0.0984	0.0987	0.0989	0.0992	0.0995	0.0997

$$k = (1-k)^{-1} = (-0.07776\dots)^{-1} = -0.53$$

$$p = 4 - k = 4 - 0.53 = 3.47$$

Application Question

- 1) An equation of diffusion of a certain gas is given as K_{2+c} .
- i) Write chairux equation of K_{2+c} .
- ii) If k is the rate of diffusion and given by 0.7 iro's.

Find the value of 'c' if a mass of 10g is released with a maximum catalyst of c gram.

Solution

i) $K_{2+c} = k_2 \times 10^c + k_c$

OR

$$K_{2+c} = k_c \times 10^2 + k_2$$

ii) $k = 0.7$ and $k_{2+c} = 10g$

$$0.7_c \times 100 + 0.7_2 = 10$$

$$0.7_c \times 100 + 7.7 = 10$$

$$0.7_c \times 100 = 10 - 7.7$$

$$0.7_c \times 100 = 2.3$$

$$0.7c = \frac{2.3}{100} = 0.023$$

$$0.7 \times 1_c = 0.023$$

$$1_c = \frac{0.023}{0.7} = 0.0329$$

$$c = 0.0329_c = 0.11$$

4 conclusion

To sum it all, the Chairux algorithm helps us in finding the divisibility test of any integers. It also helps us to find the periodical length of a given integer which is equal to the period length of reciprocal of integers. It can also be used to determine the repeated function of any data and also prove the geometric series, whereby $f_z(a+b) = z^a f_z(b) + f_z(a) = z^b f_z(a) + f_z(b)$ s.t $z = x + iy$. This method will be proved in the next paper s.t $f_z(a) = \frac{z^a - 1}{z - 1}$



5 References

References

- [1] JR Barkema. Extending stein's gcd algorithm and a comparison to euclid's gcd algorithm. B.S. thesis, 2019.
- [2] Albert B Bennett and L Ted Nelson. Divisibility tests: So right for discoveries. *Mathematics Teaching in the Middle School*, 7(8):460–465, 2002.
- [3] Lewis Berenson. A divisibility test for amateur discoverers. *The Arithmetic Teacher*, 17(1):39–41, 1970.
- [4] Yonah Cherniavsky and Artour Mouftakhov. Zbikowski's divisibility criterion. *The College Mathematics Journal*, 45(1):17–21, 2014.
- [5] M Khalid Mahmood. An appraisal algorithm for tests of divisibility using modular arithmetic. *Pakistan Journal of Science*, 67(4), 2015.
- [6] Pagdame Tiebekabe and Ismaila Diouf. New divisibility tests. *Far East Journal of Mathematical Education*, 21(1):31–41, 2021.

