

Original Research Article

Effect of Herbicides Combination on Weed Management and Yield of Direct Wet Seeded Rice (*Oryza sativa* L.)

ABSTRACT

Rice is the most stable food crop in the world, especially in Southeast Asia. Different types of methods are used in rice cultivation *viz.*, conventional method and the no-till (direct seeding) method. Nowadays, the direct seeding method of rice cultivation is gaining popularity among farmers due to its lower labour and energy requirements. In direct seeding rice weeds are the major hindrance, which compete with crops for important factors such as light, space, nutrient and moisture. To manage the weeds population in direct-seeded rice, a study was conducted in the summer, 2022 to find out the best herbicide combination. The design adopted for this experiment was Randomized Block Design (RBD) with 12 treatments and 3 replications. The results of the study revealed that a weed free plot (T₁₁) followed by the application of Pretilachlor 50% EC @ 625 g a.i. ha⁻¹ on 3 DAS + Bispyribac-sodium 10% SC @ 20 g a.i. ha⁻¹ + Ethoxysulfuron 15% WDG @ 15 g a.i. ha⁻¹ (Tank Mix) on 15 DAS and 40 DAS (T₉) recorded higher weed control efficiency, lesser weed density and weed dry weight compared to all other treatments. A similar trend was observed in yield attributes like number of tillers m⁻², number of productive tillers m⁻², panicle length, number of grains panicle⁻¹, test weight, higher grain and straw yield of direct wet seeded rice.

Key words: Herbicide mixtures, direct wet seeded rice, weeds, yield attributes

INTRODUCTION

Rice is the most significant stable food crop nourishing two third of the Southeast Asian population (Reference). In India, rice is extensively grown in an area of 47.70 million ha, with a production of 136 million metric tonnes and productivity of 4.28 metric tonnes per ha [1]. For emergence, rice needs an optimal temperature of 16°C – 27°C and rainfall of 100 cm to 200 cm during its growth period. Though rice can be grown in all kinds of soil right, from light to heavy, clay will be the most appropriate for realising maximum productivity [2]. When compared to other grains like maize, wheat and potatoes, rice is a boundless source of carbs, as well as other minerals like calcium, iron, and thiamine as well as pantothenic acids, folate, and vitamin E. It covers 5.8 g -7.7 g of protein, 0.5 g -2.3 g of fat, 0.01 g -0.78 g of

Formatted: Font: Not Italic, English (United States)

Comment [PCDP1]: Not clear, rewrite the sentence and add the conclusion

crude fiber, 0.12 g -1.22 g of crude ash and 76.80 g -86.56 g of carbohydrates [3]. In India, rice has been cultivated using various methods like transplanting and direct seeding. In transplanting, demands for continuous water stagnation has been maintained throughout the season.

The wet-seeded method of rice cutting down the labour requirement by 97% compared to traditional methods owing to the absence of nursery preparation [4]. In India, pest damage accounts for 33% of total rice yield losses, whereas weeds account for 12.5%. *Kharif* rice showed more weed flora than summer transplanted rice [5]. In the case of direct seeded rice, the emergence of weeds occurs simultaneously with the crop's emergence, which has a substantial impact on the overall quality of the crop and acts as a vector for the main insects and pests. Weeds play a negative role in the emergence of crops at early stages of growth, and they compete with other crops. To reduce the weed flora in direct seeded rice hand weeding is the best one. However, due to a lack of labour and high labour wages, this method is no longer feasible. When compared to manual weeding, chemical weed control is the most effective way to suppress the weeds in direct-sown rice.

Comment [PCDP2]: Write one paragraph on herbicide and herbicide usage of rice and compare the India with other rice cultivating countries

Nevertheless, a single weed control technique may not be sufficient to reduce weeds to the economic threshold level and may lead to a change in weed flora, the development of resistance and environmental threats. Therefore, it is imperative to adopt a variety of technologies for weed management, as weed communities are highly adaptable to management strategies [6]. Hence, an experiment was carried out to study the effect of different herbicide combinations on wet direct seeded rice.

Comment [PCDP3]: Write your hypothesis too

MATERIALS AND METHODS

To evaluate the effect of different combination of herbicides on weed control and yield of direct seeded puddled lowland rice, an experiment was conducted at Wetland farms, Agricultural College and Research Institute, Tamil Nadu Agricultural University, Coimbatore, during summer, 2022. The trial site is located at 11°N latitude and 77°E longitude with an elevation of 426.7 m above mean sea level which is classified under the western agricultural climate zone of Tamil Nadu. The edaphic characteristics of the trial site include clay loam soil type, alkaline pH of 8.10 with an EC of 0.50 (dSm⁻¹), high organic content (1.42%), low available nitrogen (125.60 kg ha⁻¹), high available phosphorus (36.75 kg ha⁻¹) and high available potassium (282.00 kg ha⁻¹).

Formatted: Font: Not Italic, English (United States)

The field experiment was designed in a Randomized Block Design (RBD) with three replications and twelve treatments. The treatment details are, T₁ - Pretilachlor 50% EC @ 625 g a.i. ha⁻¹ on 3 DAS, T₂ - Bispyribac-sodium 10% SC @ 20 g a.i. ha⁻¹ on 15 DAS, T₃ - Ethoxysulfuron 15% WDG @ 15 g a.i. ha⁻¹ on 15 DAS, T₄ - Pretilachlor 50% EC @ 625 g a.i. ha⁻¹ on 3 DAS + Bispyribac-sodium 10% SC @ 20 g a.i. ha⁻¹ on 15 DAS, T₅ - Pretilachlor 50% EC @ 625 g a.i. ha⁻¹ on 3 DAS + Ethoxysulfuron 15% WDG @ 15 g a.i. ha⁻¹ on 15 DAS, T₆ - Bispyribac-sodium 10% SC @ 20 g a.i. ha⁻¹ + Ethoxysulfuron 15% WDG @ 15 g a.i. ha⁻¹ (Tank Mix) on 15 DAS, T₇ - Pretilachlor 50% EC @ 625 g a.i. ha⁻¹ on 3 DAS + Bispyribac-sodium 10% SC @ 20 g a.i. ha⁻¹ + Ethoxysulfuron 15% WDG @ 15 g a.i. ha⁻¹ (Tank Mix) on 15 DAS, T₈ - Bispyribac-sodium 10% SC @ 20 g a.i. ha⁻¹ + Ethoxysulfuron 15% WDG @ 15 g a.i. ha⁻¹ (Tank Mix) on 15 DAS & 40 DAS, T₉ - Pretilachlor 50% EC @ 625 g a.i. ha⁻¹ on 3 DAS + Bispyribac-sodium 10% SC @ 20 g a.i. ha⁻¹ + Ethoxysulfuron 15% WDG @ 15 g a.i. ha⁻¹ (Tank Mix) on 15 DAS & 40 DAS, T₁₀ - Recommended practice (Pretilachlor 50% EC @ 750 g a.i. ha⁻¹ on 3 DAS *fb* one hand weeding on 40 DAS), T₁₁ - Weed free, T₁₂ - Weedy check. The rice variety chosen for the experiment was CO 55 and adopted a spacing of 20 cm × 10 cm. Other management practices were adopted as per the recommendation. Weed control efficiency (WCE) was calculated by using the formula,

$$\text{Weed Control Efficiency} = \frac{W_{pc} - W_{pt}}{W_{pc}} \times 100$$

Where,

W_{pc} = Weed population in the control plot

W_{pt} = Weed population in the treated plot

Observation on weed flora, density and dry weight was recorded at 20, 40 and 60 DAS and yield attributes of direct wet seeded rice was recorded at the time of harvest and the data was statistically analysed by following the analysis of variance for Randomized Block Design (RBD) as suggested by Gomez and Gomez [7]. Statistical significance was tested with the F test at a 5 per cent level of probability and compared with treatment means with the critical difference.

Comment [PCDP4]: Explain why you use particular herbicide and dosages? If you use some previous literatures, cite them

Comment [PCDP5]: Explain what is DAS in beginning for the readers (Day After Sowing)

Comment [PCDP6]: mention the statistic software that you used and post hoc test if need

RESULT AND DISCUSSION

Comment [PCDP7]: discussion is too short

A. Weed flora

Major weeds observed in the experimental field were *Echinochloa crus-galli*, *Echinochloa colona* and *Leptochloa chinensis* among grassy weeds; *Cyperus iria* and *Cyperus difformis* among sedges and *Eclipta prostrata* among broadleaved weeds (Table 1). These findings were in ~~confirmity~~ conformity with Guru [8].

B. Weed Density

Weed density represented the degree of competition from weeds on the crop (Table 2). The treatment showing the lowest weed density was considered the best. Weed control practices had a major impact on weed populations. The total weed population was significantly reduced by various weed control measures. Lower weed density was observed in weed-free plots (T_{11}) with 0.71 m^{-2} at 20, 40 and 60 DAS followed by treatment with Pretilachlor 50% EC @ $625 \text{ g a.i. ha}^{-1}$ on 3 DAS + Bispyribac 10% SC @ $20 \text{ g a.i. ha}^{-1}$ + Ethoxysulfuron 15% WDG @ $15 \text{ g a.i. ha}^{-1}$ (Tank mix) on 15 DAS and 40 DAS (T_9) with 5.92 m^{-2} , 7.40 m^{-2} and 8.87 m^{-2} at 20, 40 and 60 DAS respectively. This might be due to reduced weed infestation, as evident from lower weed density and bio-mass which facilitated better utilization of grown resources and reproductive potential of the crop. These findings were in ~~confirmity~~ conformity with Teja [9].

Comment [PCDP8]: re write the sentence according to the connection of the results.

C. Weed Dry Weight Accumulation

Higher weed dry weight was recorded (Table 3) in weedy check (T_{12}) with 9.69 g m^{-2} , 11.33 g m^{-2} , 13.11 g m^{-2} at 20, 40 and 60 DAS and lower weed dry weight was recorded in weed free (T_{11}) with 0.71 g m^{-2} at 20, 40 and 60 DAS followed by Pretilachlor 50% EC @ $625 \text{ g a.i. ha}^{-1}$ on 3 DAS + Bispyribac-sodium 10% SC @ $20 \text{ g a.i. ha}^{-1}$ + Ethoxysulfuron 15% WDG @ $15 \text{ g a.i. ha}^{-1}$ (Tank Mix) on 15 DAS and 40 DAS (T_9) with 3.44 g m^{-2} , 4.11 g m^{-2} and 6.41 g m^{-2} at 20, 40 and 60 DAS, respectively. Similar findings were also recorded Kumar [10].

Comment [PCDP9]: if someone has already done why did you do the same thing. Combine the existing result

D. Weed Control Efficiency

The weed control efficiency was maximum under the weed free plot (T₁₁) with 100% at 20, 40 and 60 DAS respectively (Fig 1), followed by treatment with Pretilachlor 50% EC @ 625 g a.i. ha⁻¹ on 3 DAS + Bispyribac-sodium 10% SC @ 20 g a.i. ha⁻¹ + Ethoxysulfuron 15% WDG @ 15 g a.i. ha⁻¹ (Tank Mix) on 15 DAS and 40 DAS with 87.52%, 87.12% and 75.49% at 20, 40 and 60 DAS, respectively. The lowest weed control efficiency was recorded in the weedy check (T₁₂). Similar results were also observed Chauhan and Johnson [11].

Comment [PCDP10]: Need to improve the sentence

E. Yield attributes

Results show that yield attributes (Table 4) like number of tillers m⁻² (649) at harvest, number of productive tillers m⁻² (366), panicle length (21.29 cm), number of filled grains panicle⁻¹ (176) and test weight (18.23 g) were found to be higher in the weed free plot (T₁₁) followed by Pretilachlor 50% EC @ 625 g a.i. ha⁻¹ on 3 DAS + Bispyribac-sodium 10% SC @ 20 g a.i. ha⁻¹ + Ethoxysulfuron 15% WDG @ 15 g a.i. ha⁻¹ (Tank Mix) on 15 DAS and 40 DAS (T₉). Which recorded more number of tillers m⁻² (632) at harvest, number of productive tillers m⁻² (350), panicle length (21.18 cm), number of filled grains panicle⁻¹ (164) and test weight (18.19 g). Lower yield attributes like a number of tillers m⁻² (536) at harvest, number of productive tillers m⁻² (150), panicle length (17.09 cm), number of grains panicle⁻¹ (81) and test weight (17.34 g). The lowest yield attributes were recorded in weedy check (T₁₂). Similar results were also reported by Ramana [12]. This was clearly related to weed competition for grown resources immediately from seedling emergence, resulting in poor crop growth and inefficient transfer of ~~photosynthates~~ photosynthesis from source to sink, resulting in decreased yield attributes of rice Umamahesh, Rao [13], [14].

Comment [PCDP11]: compare with the existing results

F. Grain and Straw Yield

Among the treatments weed free check (T₁₁) registered higher grain (5,897 kg ha⁻¹) and straw yield (9,100 kg ha⁻¹) (Fig. 2) followed by treatment Pretilachlor 50% EC @ 625 g a.i. ha⁻¹ on 3 DAS + Bispyribac-sodium 10% SC @ 20 g a.i. ha⁻¹ + Ethoxysulfuron 15%

WDG @ 15 g a.i. ha⁻¹ (Tank Mix) on 15 DAS and 40 DAS (T₉) (5,560 kg ha⁻¹ and 8,970 kg ha⁻¹) of grain and straw yield, respectively. Similarly, higher grain and straw yield were observed by Pusdekar [15].

Comment [PCDP12]: explain it

Table 1. Weed flora in the direct seeded rice during ~~Summers~~ summer season

Formatted: Font: Not Italic

S.No.	Common Name	Scientific Name	Family	Morphological type
1.	Barnyard grass	<i>Echinochloa crus-galli</i>	Poaceae	Grass
2.	Jungle rice	<i>Echinochloa colona</i>	Poaceae	Grass
3.	Chinese sprangletop	<i>Leptochloa chinensis</i>	Poaceae	Grass
4.	Rice flatsedge	<i>Cyperus iria</i>	Cyperaceae	Sedge
5.	Umbrella plant	<i>Cyperus difformis</i>	Cyperaceae	Sedge
6.	False daisy	<i>Eclipta prostrata</i>	Asteraceae	Broad leaved

Table 2. Effect of weed management practices on total weed density of different weed species at 20, 40 and 60 DAS in direct wet seeded rice

Treatments	Total density of weeds (No. m ⁻²)		
	20 DAS	40 DAS	60 DAS
T ₁ –Pretilachlor 50% EC @ 625 g a.i. ha ⁻¹ on 3 DAS	10.30 (105.67)	12.40 (153.33)	14.91 (222.33)
T ₂ –Bispyribac-sodium 10% SC @ 20 g a.i. ha ⁻¹ on 15 DAS	7.47 (55.33)	9.26 (85.33)	11.49 (131.67)
T ₃ –Ethoxysulfuron 15% WDG @ 15 g a.i. ha ⁻¹ on 15 DAS	7.36 (53.67)	8.99 (80.33)	11.07 (122.00)
T ₄ –Pretilachlor 50% EC @ 625 g a.i. ha ⁻¹ on 3 DAS + Bispyribac-sodium 10% SC @ 20 g a.i. ha ⁻¹ on 15 DAS	7.12 (50.33)	8.70 (75.33)	11.19 (125.00)
T ₅ –Pretilachlor 50% EC @ 625 g a.i. ha ⁻¹ on 3 DAS + Ethoxysulfuron 15% WDG @ 15 g a.i. ha ⁻¹ on 15 DAS	7.01 (48.67)	8.57 (73.00)	11.09 (122.67)
T ₆ –Bispyribac-sodium 10% SC @ 20 g a.i. ha ⁻¹ + Ethoxysulfuron 15% WDG @ 15 g a.i. ha ⁻¹ (Tank Mix) on 15 DAS	7.21 (51.67)	8.37 (69.67)	11.19 (125.00)
T ₇ –Pretilachlor 50% EC @ 625 g a.i. ha ⁻¹ on 3 DAS + Bispyribac-sodium 10% SC @ 20 g a.i. ha ⁻¹ + Ethoxysulfuron 15% WDG @ 15 g a.i. ha ⁻¹ (Tank Mix) on 15 DAS	6.59 (43.00)	8.09 (65.00)	10.68 (113.67)
T ₈ –Bispyribac-sodium 10% SC @ 20 g a.i. ha ⁻¹ + Ethoxysulfuron 15% WDG @ 15 g a.i. ha ⁻¹ (Tank Mix) on 15 DAS & 40 DAS	6.91 (47.33)	8.36 (69.67)	10.04 (101.00)
T ₉ –Pretilachlor 50% EC @ 625 g a.i. ha ⁻¹ on 3 DAS + Bispyribac-sodium 10% SC @ 20 g a.i. ha ⁻¹ + Ethoxysulfuron 15% WDG @ 15 g a.i. ha ⁻¹ (Tank Mix) on	5.92 (34.67)	7.40 (54.33)	8.87 (78.33)

15 DAS & 40 DAS			
T ₁₀ – Recommended practice (Pretilachlor 50% EC @ 750 g a.i. ha ⁻¹ on 3 DAS <i>fb</i> one hand weeding on 40 DAS)	8.87 (78.33)	10.89 (118.00)	9.53 (90.33)
T ₁₁ – Weed free	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)
T ₁₂ – Weedy check	10.9 (118.67)	15.08 (227.33)	18.04 (325.00)
S. Ed	0.30	0.34	0.44
CD (P=0.05%)	0.62	0.70	0.91

Comment [PCDP13]: better to show the post hoc comparison too

Figures in parenthesis are original values; Data subjected to square root transformation.

Table 3. Effect of weed management practices on total weed dry weight accumulation (g m⁻²) of different weed species at 20, 40 and 60 DAS in direct wet seeded rice

Comment [PCDP14]: show the post hoc comparison too

Treatments	Total weed dry weight accumulation (g m ⁻²)		
	20 DAS	40 DAS	60 DAS
T ₁ –Pretilachlor 50% EC @ 625 g a.i. ha ⁻¹ on 3 DAS	7.72 (59.07)	9.81 (95.73)	10.59 (111.89)
T ₂ –Bispyribac-sodium 10% SC @ 20 g a.i. ha ⁻¹ on 15 DAS	4.03 (15.72)	7.10 (49.97)	8.24 (67.67)
T ₃ –Ethoxysulfuron 15% WDG @ 15 g a.i. ha ⁻¹ on 15 DAS	3.96 (15.17)	6.58 (43.47)	8.34 (69.07)
T ₄ –Pretilachlor 50% EC @ 625 g a.i. ha ⁻¹ on 3 DAS + Bispyribac-sodium 10% SC @ 20 g a.i. ha ⁻¹ on 15 DAS	3.90 (14.75)	6.73 (44.87)	7.92 (62.33)
T ₅ –Pretilachlor 50% EC @ 625 g a.i. ha ⁻¹ on 3 DAS + Ethoxysulfuron 15% WDG @ 15 g a.i. ha ⁻¹ on 15 DAS	3.94 (15.01)	6.62 (43.4)	7.86 (61.32)
T ₆ –Bispyribac-sodium 10% SC @ 20 g a.i. ha ⁻¹ + Ethoxysulfuron 15% WDG @ 15 g a.i. ha ⁻¹ (Tank Mix) on 15 DAS	3.86 (14.42)	6.37 (40.07)	7.58 (56.97)
T ₇ –Pretilachlor 50% EC @ 625 g a.i. ha ⁻¹ on 3 DAS + Bispyribac-sodium 10% SC @ 20 g a.i. ha ⁻¹ + Ethoxysulfuron 15% WDG @ 15 g a.i. ha ⁻¹ (Tank Mix) on 15 DAS	3.71 (13.32)	6.43 (40.87)	7.73 (59.27)
T ₈ –Bispyribac-sodium 10% SC @ 20 g a.i. ha ⁻¹ + Ethoxysulfuron 15% WDG @ 15 g a.i. ha ⁻¹ (Tank Mix) on 15 DAS & 40 DAS	3.66 (12.89)	6.83 (46.13)	7.58 (56.95)
T ₉ –Pretilachlor 50% EC @ 625 g a.i. ha ⁻¹ on 3 DAS + Bispyribac-sodium 10% SC @ 20 g a.i. ha ⁻¹ + Ethoxysulfuron 15% WDG @ 15 g a.i. ha ⁻¹ (Tank Mix) on 15 DAS & 40 DAS	3.44 (11.36)	4.11 (16.4)	6.41 (40.67)
T ₁₀ – Recommended practice (Pretilachlor 50% EC @ 750 g a.i. ha ⁻¹ on 3 DAS <i>fb</i> one hand weeding on 40 DAS)	7.29 (52.66)	7.94 (62.57)	7.25 (52.09)
T ₁₁ – Weed free	0.71	0.71	0.71

	(0.00)	(0.00)	(0.00)
T ₁₂ - Weedy check	9.69 (94.3)	11.33 (128.13)	13.11 (171.33)
S. Ed	0.29	0.28	0.25
CD (P=0.05%)	0.60	0.59	0.53


Figures in parenthesis are original values; Data subjected to square root transformation.

Table. 4. Effect of weed management practices on Numbers of tillers (m⁻²) at harvest, No. of productive tillers (m⁻²), No. of filled grains panicle⁻¹, Panicle length (cm), Panicle weight (g), Test weight (g), Grain conversion efficiency (%) in direct wet seeded rice

Comment [PCDP15]: better to show the post hoc comparison too


Treatments	Yield components				
	Numbers of tillers (m ⁻²) at harvest	No. of productive tillers (m ⁻²)	No. of filled grains panicle ⁻¹	Panicle length (cm)	Test weight (g)
T ₁ - Pretilachlor 50% EC @ 625 g a.i. ha ⁻¹ on 3 DAS	560.00	166.00	95.00	17.89	17.35
T ₂ - Bispyribac-sodium 10% SC @ 20 g a.i. ha ⁻¹ on 15 DAS	571.00	191.00	110.00	18.17	17.39
T ₃ - Ethoxysulfuron 15% WDG @ 15 g a.i. ha ⁻¹ on 15 DAS	584.00	250.00	127.00	18.9	17.58
T ₄ - Pretilachlor 50% EC @ 625 g a.i. ha ⁻¹ on 3 DAS + Bispyribac-sodium 10% SC @ 20 g a.i. ha ⁻¹ on 15 DAS	579.00	200.00	119.00	18.62	17.44
T ₅ - Pretilachlor 50% EC @ 625 g a.i. ha ⁻¹ on 3 DAS + Ethoxysulfuron 15% WDG @ 15 g a.i. ha ⁻¹ on 15 DAS	597.00	258.00	129.00	19.78	17.62
T ₆ - Bispyribac-sodium 10% SC @ 20 g a.i. ha ⁻¹ + Ethoxysulfuron 15% WDG @ 15 g a.i. ha ⁻¹ (Tank Mix) on 15 DAS	601.00	275.00	135.00	19.92	17.75
T ₇ - Pretilachlor 50% EC @ 625 g a.i. ha ⁻¹ on 3 DAS + Bispyribac-sodium 10% SC @ 20 g a.i. ha ⁻¹ + Ethoxysulfuron 15% WDG @ 15 g a.i. ha ⁻¹ (Tank Mix) on 15 DAS	609.00	325.00	140.00	20.34	17.87
T ₈ - Bispyribac-sodium 10% SC @ 20 g a.i. ha ⁻¹ + Ethoxysulfuron 15% WDG @ 15 g a.i. ha ⁻¹ (Tank Mix) on 15 DAS & 40 DAS	614.00	325.00	143.00	20.79	17.98
T ₉ - Pretilachlor 50% EC @ 625 g a.i.	632.00	350.00	164.00	21.18	18.19

ha ⁻¹ on 3 DAS + Bispyribac-sodium 10% SC @ 20 g a.i. ha ⁻¹ + Ethoxysulfuron 15% WDG @ 15 g a.i.ha ⁻¹ (Tank Mix) on 15 DAS & 40 DAS					
T ₁₀ - Recommended practice (Pretilachlor 50% EC @ 750 g a.i. ha ⁻¹ on 3 DAS <i>fb</i> one hand weeding on 40 DAS)	627.00	338.00	155.00	21.09	18.05
T ₁₁ - Weed free	649.00	366.00	176.00	21.29	18.23
T ₁₂ - Weedy check	536.00	150.00	81.00	17.30	17.34
S. Ed	29.48	12.99	6.44	0.97	0.88
CD (P=0.05%)	61.14	26.93	13.36	2.01	NS



Comment [PCDP16]: keep axis to 100 and remove the decimals

Fig. 1. Effect of weed management practices on Weed Control Efficiency (WCE) 20, 40 and 60 DAS in direct wet seeded rice



Comment [PCDP17]: Show the significance statistically and show the small letters on the bars with significantly different treatments

Fig. 2. Effect of weed management practices on Grain yield (kg ha⁻¹) and Straw yield (kg ha⁻¹) in direct wet seeded rice

F. Conclusion

Based on the results obtained from the study, it could be concluded that the lowest weed density and weed dry matter accumulation, maximum weed control efficiency and with higher grain and straw yield were observed in weed free treatment followed by Pretilachlor 50% EC @ 625 g a.i. ha⁻¹ on 3 DAS + Bispyribac-sodium 10% SC @ 20 g a.i. ha⁻¹ + Ethoxysulfuron 15% WDG @ 15 g a.i.ha⁻¹ (Tank Mix) on 15 DAS & 40 DAS. Which recorded lower weed density, higher weed control efficiency, grain and straw yield of direct wet seeded rice.

Comment [PCDP18]: go with the format of the journal

REFERENCE

1. USDA. (2023). <https://apps.fas.usda.gov/psdonline/circulars/production>.
2. Mahajan G, Chauhan BS, Johnson DE. Weed management in aerobic rice in Northwestern Indo-Gangetic Plains. *Journal of Crop Improvement*. 2009 Oct 1;23(4):366-82.
3. RathnaPriya TS, Eliazer Nelson AR, Ravichandran K, Antony U. Nutritional and functional properties of coloured rice varieties of South India: a review. *Journal of Ethnic Foods*. 2019 Dec;6(1):1-1.
4. Kathia MK, Mati B, Ndiiri J, Wanjogu R. Integrating mechanical weeding and planting for reduced labour input in paddy rice under System of Rice Intensification (SRI). *Agricultural Sciences*. 2019 Feb 18;10(2):121-30.
5. Mondal D, Ghosh A, Roy D, Kumar A, Shamurailatpam D, Bera S, Ghosh R, Bandopadhyay P, Majumder A. Yield loss assessment of rice (*Oryza Sativa* L.) due to different biotic stresses under system of rice intensification (SRI). *Journal of Entomology and Zoology Studies*. 2017;5(4):1974-80.
6. Buhler DD, Hartzler RG, Forcella F. Implications of weed seedbank dynamics to weed management. *Weed Science*. 1997 Jun;45(3):329-36.
7. Gomez KA, Gomez AA. *Statistical procedures for agricultural research*. John Wiley & sons; 1984 Feb 17.
8. Guru RS, Dwivedi SK, Bhambri MC. Evaluation of different weed management approaches in managing weeds in direct seeded rice (*Oryza sativa* L.).
9. Teja KC, Duary B, Bhowmik MK. Efficacy of herbicides on weed management in wet season transplanted rice. *Journal of Crop and Weed*. 2015;11(Special Issue):224-7.

Comment [PCDP19]: Reference are not in same format, information are missing

~~10.~~ Kumar P, Singh Y, Singh UP. Evaluation of cultivars and herbicides for control of barnyard grass and nutsedge in boro rice.

10.

11. Chauhan BS, Johnson DE. Ecological studies on Echinochloa crus-galli and the implications for weed management in direct-seeded rice. *Crop Protection*. 2011 Nov 1;30(11):1385-91.
12. Ramana AV, Naidu GJ, Murthy KV. Integrated weed management in rainfed upland rice (*Oryzasativa*). *Indian Journal of Agronomy*. 2007;52(4):311-4.
13. Umamahesh V, Madhuri KV, Chandrika V, Sagar GK, Adilakshmi G. Studies on Nutrient and Weed Dynamics on Growth and Yield Attributes of Direct Seeded Rice. *International Journal of Environment and Climate Change*. 2022 Nov 12;12(12):20-8.
14. Rao AN, Johnson DE, Sivaprasad B, Ladha JK, Mortimer AM. Weed management in direct- seeded rice. *Advances in agronomy*. 2007 Jan 1;93:153-255.
15. Pusdekar VR, Pagar PC, Atram KS, Kothikar RB, Mairan NR. Effect of pre and post emergence herbicide on weed control in direct seeded lowland rice. *Journal of Pharmacognosy and Phytochemistry*. 2020;9(6):959-62.

Formatted: Font: (Default) Times New Roman, 12 pt, Font color: Custom Color(RGB(34,34,34))

Formatted: Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.25" + Indent at: 0.5"

UNDER PEER REVIEW