

## **Original Research Article**

# **Effect of inorganic fertilizers and manures on yield and soil nutrient status of Maize in calcareous *Inceptisol* over five decades of intensive farming**

### **Abstract**

Long Term Fertilizer Experiment were initiated with an aim to monitor the continuous manuring and fertilization under intensive cropping changes in soil properties with respect to physical, chemical and biological characteristics and crop productivity. Continuous use of inorganic fertilizer along with FYM could increase the soil nutrients such as  $\text{KMnO}_4 - \text{N}$ , Olsen - P and  $\text{NNH}_4\text{OAc} - \text{K}$  over five decades. The experimental soil belongs to sandy clay loam soil (*Vertic Ustropept*) with ten different nutrient management practices are followed and replicated thrice by adoption RBD. The results revealed that the use of inorganic fertilizer conjoint with FYM @  $10 \text{ t ha}^{-1}$  significantly increased the kernel yield of maize by 18.1% and 20.3% higher than 150% NPK and 100% NPK plots in sandy clay loam soil. The important indicator for soil fertility concern is SOC and the build up of SOC was found in NPK+FYM @  $10 \text{ t ha}^{-1}$  treatment recorded from  $3.0 \text{ g kg}^{-1}$  (1972) to  $7.8 \text{ g kg}^{-1}$  (2023). Soil fertility status of the experiment after 50 years had reported as medium in  $\text{KMnO}_4 - \text{N}$ , medium to high in Olsen - P, high in  $\text{NNH}_4\text{OAc} - \text{K}$  and medium in SOC status in an *Inceptisol*. From the study our hypothesis revealed that, application of inorganic fertilizer along with FYM @  $10 \text{ t ha}^{-1}$  achieved the greater yield and better soil health in sustainable manner.

**Key Words:** *Farmyard manure, SOC, Inorganic fertilizers and Soil Fertility*

### **1. INTRODUCTION**

Soil is a vital natural resources and Soil quality is the combined effect of management on most soil properties which influence the crop productivity and sustainability (Sharma *et al.*, 2005). The maintenance of soil health is not only important increasing the agricultural production but also essential for sustaining the higher productivity of crops (Velu and Subramaniam, 2012). Soil fertility is the most important factor which controls the crop productivity. Soil related constraints affecting crop yield such as nutritional disorders can be assessed by evaluating the soil fertility status. Soil testing provides information about the nutrient status of the soil from which fertilizer recommendation can be made for maximizing the crop productivity (Haribhushan Athokpam *et al.*, 2013).

In recent years, intensive farming practices using high yielding cultivars with imbalanced fertilization have resulted in over mining of native soil nutrients that adversely affects the crop production and soil fertility status (Tandon 1999 and Swarup and Ganeshmurthy, 1998). It is important to maintain the nutrient supply in order to sustain productivity. Since large amount of nutrient has to be applied to soil in chemical form which may have impact on soil properties and soil productivity in long term. Long Term Fertilizer Experiment were initiated with an aim to monitor the changes in soil properties as a result of continuous manuring and fertilization under intensive cropping with respect to the physical, chemical and microbiological characteristics of the soil in relation to its productivity. The integrated application of organic manures and inorganic fertilizers is the best recognized strategy of Integrated Nutrient Management to improve the soil fertility and also enhancing the crop yields (Manimaran *et al.*, 2022).

Maize (*Zea mays*) is most important coarse cereal crop belongs to Poaceae family. It is widely used for cattle feed, poultry feed and also for human consumption (Gul *et al.*, 2021). It is adapted to wide range of soil and agro-climatic conditions but sensitive to water logging hence, it is known for Queen of Cereals. It is more nutrient exhausting crop. Globally, India ranks fourth in area and seventh in production among maize-growing countries, Maize is grown on 9.2 million hectares in India, it accounts for around 4% of global maize area and 2% of total production (DACNET, 2020). In India Maize is the third most important cereal crop next to Rice and Wheat.

In this article an attempt has been made to highlight the Importance of INM practices in maintaining soil health and improving in crop yield and productivity.

## 2. MATERIALS AND METHODS

### 2.1 Experimental site and climate

The study area is located at the Research farm of Tamil Nadu Agricultural University, Coimbatore district of Tamil Nadu, India. The Long Term Fertilizer Experiment (LTFE) was started in 1972 at 11° N latitude, 77° E longitude with an elevation of 426.7 m above MSL with Finger millet- Maize cropping sequence under irrigated condition, soil classified as *Inceptisol* having black calcareous sandy clay loam soil (*vertic ustropept*) belongs to Periyanaickenpalayam soil series. The climate of Coimbatore is tropical characterized by hot and humid summers and cold winters. The NE monsoon majorly contributes to Coimbatore, the average annual precipitation ranges between 550 mm to 900 mm.

### 2.2 Experimental details

LTFE also includes two crops per year, Finger Millet – Maize cropping sequence raised during summer and kharif seasons respectively with ten treatment combinations and three replications, the plots are randomized by Randomized Block Design (RBD). The treatment details are T<sub>1</sub>- control, T<sub>2</sub>-100% N, T<sub>3</sub>-100% NP, T<sub>4</sub>- 100% NPK, T<sub>5</sub>-100% NPK+FYM @ 10 t ha<sup>-1</sup>, T<sub>6</sub>-100% NPK+ Zn, T<sub>7</sub>- 150% NPK T<sub>8</sub>- 50% NPK, T<sub>9</sub>-100% NPK+HW and T<sub>10</sub>-100% NPK ( - S ). In this study a Maize hybrid (COHM 6) as a test crop and ten treatments with three replications were taken for investigation.

The recommended dose of fertilizer for Maize hybrid were 250, 75 and 75 kg of N, P and K ha<sup>-1</sup> respectively. At the time of sowing 50% of RDF N were applied as urea and 100% of RDF of P and K were applied as SSP and MOP respectively for all the treatments except for T<sub>9</sub> (100% NPK – S) DAP can be used as a source. Another 50% of N were applied in two equal splits during knee high stage and pre tasselling stages of crop growth. For INM (100% NPK + FYM) treatment plots were applied 10 t ha<sup>-1</sup> of FYM for every crop.

### 2.3 Soil Sampling, Processing and Analysis

The soil samples were collected from 50 years old Long Term Fertilizer Experiment from the individual treatment plots after the harvest of Finger millet and Maize respectively at the depth of 0 – 15 cm. the soils are air dried, ground and sieved to pass through 2 mm sieve. The processed soil samples were used for determination of various soil physical and chemical properties by following standard procedures. Soil texture was determined by international pipette method. pH and EC were determined in the soil and water suspension in the ratio of 1:2.5 by potentiometric and conductometric methods respectively (Jackson 1973), Available N in soil was estimated by alkaline KMnO<sub>4</sub> method (Asija and Subbaiah 1956), Available P in soil was estimated by NaHCO<sub>3</sub> extractions and determined colorimetrically (Olsen *et al* 1954). Available K in soil was estimated by using neutral normal Ammonium Acetate extractions followed by Emission spectroscopy (Standford and English 1949). Soil organic carbon was determined by chromic acid wet digestion method (Walkley and Black 1934).

### 2.4 Statistical Analysis

The recorded data from field experiment were analysed for their significance (p=0.05) by statistical procedure appropriate for Randomized Block Design outlined by Gomez and Gomez 1985.

## 3. RESULTS AND DISCUSSION

### 3.1 Initial physical, physico- chemical and chemical properties of soil during 1972.

The initial properties of experimental soil was analyzed for various physical, physico- chemical and chemical properties. The pH of the soil was found to be alkaline in nature, the primary nutrient status of the soil

were found to be low in  $\text{KMnO}_4\text{-N}$ , medium in Olsen - P, high in  $\text{NNH}_4\text{OAc-K}$  and medium in SOC status in a Sandy Clay Loam soil (Table 1).

**Table 1. Initial physical, physico- chemical and chemical properties of soil during 1972.**

Characteristics of Soil	(1972) Value
<b>Particle size analysis</b>	
Clay (%)	32.6
Silt (%)	11.8
Fine sand (%)	15.1
Coarse Sand (%)	39.4
Soil textural class	Sandy clay loam
<b>Physicochemical and chemical properties</b>	
pH (1:2.5 soil: water suspension)	8.20
Electrical conductivity ( $\text{dS m}^{-1}$ )	0.20
Organic carbon ( $\text{g kg}^{-1}$ )	3.0
Cation exchange capacity [ $\text{c mol (p+) kg}^{-1}$ ]	25.5
Available nitrogen ( $\text{kg N ha}^{-1}$ )	178.0
Available phosphorus ( $\text{kg ha}^{-1}$ )	11.0
Available potassium ( $\text{kg ha}^{-1}$ )	810.0

### 3.2 Long term fertilization effects on soil physico chemical properties of Maize crop in an *inceptisol* (post-harvest samples of 115<sup>th</sup> crop)

#### 3.2.1 Soil reaction (pH) and Electrical Conductivity ( $\text{d Sm}^{-1}$ )

The soil reaction pH and EC were changes slightly from 1972, currently soil reaction of the research soil was alkaline in nature and pH ranges from 8.07 to 8.40 (Table 3.) specifies that there is no significant influence of pH on soil due to buffering capacity of the soil. The slight changes in the pH was due to release organic acids and continuous addition of inorganic fertilizers and organic manures. Likewise, EC ranges from 0.51 to 0.63 (Table 3.) indicates that there is no significant impact of EC on different fertilization treatments (Malarkodi *et al.*, 2019).

### 3.3 Long term fertilization effects on soil chemical properties of Maize crop in an *inceptisol* (post-harvest samples of 115<sup>th</sup> crop)

#### 3.3.1 Soil Organic Carbon SOC ( $\text{g kg}^{-1}$ )

Soil Organic Carbon (SOC) is the important factor which governs the soil fertility (Prabakaran *et al.*, 2023). The continuous application of inorganic fertilizers and organic manures over five decades in a Finger millet – Maize sequence has increased the SOC content from  $3.0 \text{ g kg}^{-1}$  to  $7.8 \text{ g kg}^{-1}$  (Nayak *et al.*, 2012). The SOC concentration has increased significantly under different fertilized and unfertilized treatments from control to 150% NPK ( $4.0 \text{ g kg}^{-1}$  to  $7.8 \text{ g kg}^{-1}$ ) (Table 2). The maximum SOC concentration was recorded in 100% NPK + FYM @  $10 \text{ t ha}^{-1}$  plot at the rate of  $7.8 \text{ g kg}^{-1}$  followed by 150% NPK at  $7.1 \text{ g kg}^{-1}$  (Bhattacharyya *et al.*, 2010) and (Srinivasarao *et al.*, 2011). The treatments like 100% NPK > 100% NPK +Zn >100% NPK +HW are on par with each other and 100% NP & 100% NPK- S are also on par with each other. The minimum SOC concentration was recorded in control plot at the rate of  $4.0 \text{ g kg}^{-1}$

#### 3.3.2 Soil Available Nitrogen ( $\text{kg ha}^{-1}$ )

Nitrogen is one of the major plant nutrient which imparts green colour to plant and encourages vegetative growth. The continuous application of inorganic fertilizers and organic manures has significantly influenced the available N status in the soil under different management treatments and the available N content in the Experimental soil was found to be medium and ranged from 260 to 408 kg ha<sup>-1</sup> (Table 3). The maximum available N content was recorded in (100% NPK + FYM @ 10 t ha<sup>-1</sup>) plot of 408 kg ha<sup>-1</sup> followed by 150% NPK plot of 378kg ha<sup>-1</sup> (Bairwa *et al.*, 2021). However, the treatments like 100% NPK > 100% NPK +Zn >100% NPK +HW are on par with each other. The 100% NPK plots showed that declined available N content than other treatments. The minimum available N content was recorded in control plots of 260kg ha<sup>-1</sup>. The combined application of inorganic nitrogenous fertilizer along with FYM has increased the mineralization of organic form of N to Inorganic forms.

### 3.3.3 Soil Available Phosphorous (kg ha<sup>-1</sup>)

Phosphorous is also a primary macronutrient which plays important role in energy storage and transfer. The available P content in research area was found to be medium to high under different treatments and the values ranged between 16.13 to 29.32 kg ha<sup>-1</sup> (Table 3). The maximum available P content was recorded in (100% NPK + FYM @ 10 t ha<sup>-1</sup>) plot of 29.32 kg ha<sup>-1</sup> followed by 150% NPK plot of 28.27 kg ha<sup>-1</sup> (Kalaiselvi *et al.*, 2021) and (Mitran *et al.*, 2016). Other treatments like 100% NP > 100% NPK >100% NPK +Zn are on par with each other. The minimum available P content was recorded in control plots of 16.13 kg ha<sup>-1</sup>. The soil pH is a most important factor which influences the P fixation in the soil. Under acidic condition (pH < 6) P fixed as AlPO<sub>4</sub> and FePO<sub>4</sub> and in alkaline (pH > 8) condition fixed as a CaPO<sub>4</sub>. The greater P content in INM treatment plot was due to presence Organic Matter leads to the formation of organophosphate compounds are more soluble and also increase the quantity of organic P to inorganic P.

### 3.3.4 Soil Available Potassium (kg ha<sup>-1</sup>)

Potassium is the third most important primary nutrient absorbed by the plants in larger amount than the other nutrients. The available K content in the research soil was found to be high since from the start of experiment (1972). The use of high yield cultivars and inorganic fertilizers has significantly exhausted the soil K under different treatments and ranges between 236 to 642 kg ha<sup>-1</sup> (Table 3). The maximum available K content was recorded in (100% NPK + FYM @ 10 t ha<sup>-1</sup>) plot of 642 kg ha<sup>-1</sup> followed by 150% NPK plot of 564 kg ha<sup>-1</sup> (Mazumdar *et al.*, 2014). Other treatments like 100% NPK > 100% NPK +Zn >100% NPK +HW are on par with each other. The minimum available P content was recorded in control plots of 236 kg ha<sup>-1</sup>. The application of NPK with FYM has increased the K availability in soil due to release of fixed K ions in the exchangeable sites (Muneshwar *et al.*, 2014).

**Table 2. Long term fertilization effects on soil Physico– chemical and chemical properties of Maize (kg ha<sup>-1</sup>) in an *Inceptisol* (Initial samples of 115<sup>th</sup> crop)**

Treatments	pH	EC dSm <sup>-1</sup>	SOC g kg <sup>-1</sup>	N Kg ha <sup>-1</sup>	P Kg ha <sup>-1</sup>	K Kg ha <sup>-1</sup>
Control	8.09	0.52	0.40	227	11.09	259
50 % NPK	8.21	0.58	0.54	276	16.76	318
100% N	8.25	0.62	0.51	285	12.98	295
100% NP	8.46	0.58	0.60	289	18.94	307
100% NPK	8.23	0.57	0.63	321	19.08	476
100% NPK+ FYM	8.30	0.65	0.78	357	21.99	538
100% NPK +Zn	8.22	0.59	0.66	309	18.67	482
100% NPK +HW	8.15	0.56	0.64	302	18.54	475
100% NPK- S	8.36	0.63	0.61	293	17.91	434
150% NPK	8.18	0.62	0.71	336	20.76	502


<b>SEd</b>	<b>1.222</b>	<b>0.0097</b>	<b>0.0148</b>	<b>7.3533</b>	<b>0.4369</b>	<b>10.2525</b>
<b>CD (P= 0.05)</b>	<b>2.5673</b>	<b>0.0205</b>	<b>0.0311</b>	<b>15.449</b>	<b>0.918</b>	<b>21.54</b>

**Table 3. Long term fertilization effects on soil Physico–chemical and chemical properties of Maize (kg ha<sup>-1</sup>) in an *Inceptisol* (post-harvest samples of 115<sup>th</sup> crop)**

Treatments	pH	EC	N Kg ha <sup>-1</sup>	P Kg ha <sup>-1</sup>	K Kg ha <sup>-1</sup>
<b>control</b>	8.07	0.51	260.3	16.13	236
<b>50 % NPK</b>	8.20	0.57	301.3	21.27	376
<b>100% N</b>	8.25	0.63	314.3	19.53	312
<b>100% NP</b>	8.40	0.59	339.7	24.77	327
<b>100% NPK</b>	8.23	0.57	363.7	25.30	558
<b>100% NPK+ FYM</b>	8.27	0.67	408.7	29.32	642
<b>100% NPK +Zn</b>	8.21	0.60	366.7	25.50	497
<b>100% NPK +HW</b>	8.16	0.53	356.0	24.43	508
<b>100% NPK- S</b>	8.34	0.62	338.3	24.25	478
<b>150% NPK</b>	8.15	0.63	378.0	28.27	564
<b>SEd</b>	<b>0.1977</b>	<b>0.0154</b>	<b>6.6191</b>	<b>0.4255</b>	<b>11.4556</b>
<b>CD (P=0.05)</b>	<b>0.4154</b>	<b>0.03</b>	<b>13.095</b>	<b>0.8939</b>	<b>24.068</b>

### 3.4 Long term fertilization effects on Kernel and Stover yields of Maize (kg ha<sup>-1</sup>) in an *Inceptisol*

The continuous application of inorganic fertilizers and manures caused significant variation in kernel and straw yield of Maize in an *inceptisol* (Fig.1). The maximum kernel and Stover yield of Maize has recorded in the 100%NPK + FYM @ 10 t ha<sup>-1</sup> plot of 6898 kg ha<sup>-1</sup> and 9356 kg ha<sup>-1</sup> respectively followed by 150% NPK plot 5643 kg ha<sup>-1</sup> and 8978 kg ha<sup>-1</sup> (Choudhary *et al.*, 2021). The minimum kernel and straw yield of Maize was recorded in the control plot of 2872 kg ha<sup>-1</sup> and 4317 kg ha<sup>-1</sup> respectively. The yield increase in 100% NPK + FYM @ 10 t ha<sup>-1</sup> plot obtained was 18.1% and 20.3% higher than 150% NPK and 100% NPK plots respectively this might be due to conjoint application of NPK + FYM has significantly improved the soil physical, chemical and biological properties of soil leads to increase the yield and productivity of the soil in a sustained manner (Majhi *et al.*, 2019) and (Sharma *et al.*, 2018).



## Fig.1 Long term fertilization effects on kernel and Straw yields of Maize (kg ha<sup>-1</sup>) in an *Inceptisol*

### 4. CONCLUSION

The present study revealed that the application of recommended dose of fertilizer along with the organic manure (i.e 100% NPK + FYM @ 10 ha<sup>-1</sup>) was considered to be effective among different nutrient management treatments under sandy clay loam of Maize crop. Balanced application of nutrients would improve the physical, chemical and biological properties of the soil as well as maintain the soil fertility and also enhance the productivity of Maize crop under irrigated condition. With this result, our hypothesis revealed that, application of inorganic fertilizer along with FYM@10 t ha<sup>-1</sup> achieved the greater yield and better soil fertility in sustainable manner.

### REFERENCE

1. Sharma, K. L., Mandal, U. K., Srinivas, K., Vittal, K. P. R., Mandal, B., Grace, J. K., & Ramesh, V. (2005). Long-term soil management effects on crop yields and soil quality in a dryland Alfisol. *Soil and Tillage Research*, 83(2), 246-259.
2. Velu, V., & Subramaniam, P. (2012). Soil health and its sustenance through enriched compost. In *National seminar on recycling of solid wastes through composting held during March 8th and 9th at the Department of Soil and Environment, Agricultural College and Research Institute, Madurai* (pp. 5-11).
3. Haribhushan, A., Shabir, H. W., David, K., Herojit, S. A., Jyotsna, N., Deepak, K., ... & Lamalakshmi, D. (2013). Soil macro-and micro-nutrient status of Senapati district, Manipur (India). *African Journal of Agricultural Research*, 8(39), 4932-4936.
4. Tandon, H. L. S. (1995). Micronutrients in soils, crops and fertilisers-a sourcebook-cum-directory (1995)-. *Micronutrients in soils, crops and fertilisers-a sourcebook-cum-directory (1995)*.
5. Swarup, A., & Ganeshamurthy, A. N. (1998). Emerging Nutrient Deficiencies under Intensive Cropping Systems and Remedial Measures for Sustainable High Productivity. *Fertiliser news*, 43(7).
6. Manimaran, G., Jayanthi, D., Janaki, P., Amirtham, D., & Gokila, B. (2022). Long Term Impact of Fertilization and Intensive Cropping on Maize Yield and Soil Nutrient Availability under Sandy Clay Loam Soil (Inceptisol). *International Journal of Plant & Soil Science*, 34(20), 795-801.
7. Gul, H., Rahman, S., Shahzad, A., Gul, S., Qian, M., Xiao, Q., & Liu, Z. (2021). Maize (*Zea mays* L.) productivity in response to nitrogen management in Pakistan. *American Journal of Plant Sciences*, 12(8), 1173-1179.
8. Malarkodi, M., Elayarajan, M., Arulmozhiselvan, K., & Gokila, B. (2019). Long-term impact of fertilizers and manures on crop productivity and soil fertility in an alfisol. *The Pharma Innovation Journal*, 8(7), 252-256.
9. Prabakaran, S., Kaleeswari, R. K., Backiyavathy, M. R., Jagadeeswaran, R., Selvi, R. G., & Bama, K. S. (2023). Estimation of soil carbon pools under major cropping systems of Mayiladuthurai district of Cauvery Delta Zone, Tamil Nadu, India. *Journal of Applied and Natural Science*, 15(2), 802-810.
10. Nayak, A. K., Gangwar, B., Shukla, A. K., Mazumdar, S. P., Kumar, A., Raja, R., ... & Mohan, U. (2012). Long-term effect of different integrated nutrient management on soil organic carbon and its fractions and sustainability of rice-wheat system in Indo Gangetic Plains of India. *Field Crops Research*, 127, 129-139.
11. Bhattacharyya, R., Kundu, S., Srivastva, A. K., Gupta, H. S., Prakash, V., & Bhatt, J. C. (2011). Long term fertilization effects on soil organic carbon pools in a sandy loam soil of the Indian sub-Himalayas. *Plant and soil*, 341, 109-124.
12. Srinivasarao, C. H., Venkateswarlu, B., Lal, R., Singh, A. K., Kundu, S., Vittal, K. P. R & Patel, M. M. (2014). Long-term manuring and fertilizer effects on depletion of soil organic carbon stocks under pearl millet-cluster bean-castor rotation in Western India. *Land Degradation & Development*, 25(2), 173-183.

13. Bairwa, J., Dwivedi, B. S., Rawat, A., Thakur, R. K., & Mahawar, N. (2021). Long-term effect of nutrient management on soil microbial properties and nitrogen fixation in a vertisol under soybean-wheat cropping sequence. *Journal of the Indian Society of Soil Science*, 69(2), 171-178.
14. Kalaiselvi, K., Jayanthi, D., Santhy, P., Gnanachitra, M., & Gokila, B. (2021). Effect of long term fertilization on phosphorus dynamics in root zone environment under finger millet-Maize cropping sequence. *Journal of Applied and Natural Science*, 13(4), 1383-1389.
15. Mitran, T., Mani, P. K., Basak, N., Mazumder, D., & Roy, M. (2016). Long-term manuring and fertilization influence soil inorganic phosphorus transformation vis-a-vis rice yield in a rice-wheat cropping system. *Archives of Agronomy and Soil Science*, 62(1), 1-18.
16. Mazumdar, S. P., Kundu, D. K., Ghosh, D., Saha, A. R., Majumdar, B., & Ghorai, A. K. (2014). Effect of long-term application of inorganic fertilizers and organic manure on yield, potassium uptake and distribution of potassium fractions in the new Gangetic alluvial soil under jute-rice-wheat cropping system. *International Journal of Agriculture and Food Science Technology*, 5(4), 297-306.
17. Singh, M., Ram, S., Wanjari, R. H., & Sharma, P. (2014). Balance and forms of potassium under rice-wheat system in a 40-year-old long-term experiment on Mollisols of Pantnagar. *Journal of the Indian Society of Soil Science*, 62(1), 38-44.
18. Choudhary, M., Meena, V. S., Panday, S. C., Mondal, T., Yadav, R. P., Mishra, P. K. & Pattanayak, A. (2021). Long-term effects of organic manure and inorganic fertilization on biological soil quality indicators of soybean-wheat rotation in the Indian mid-Himalaya. *Applied Soil Ecology*, 157, 103754.
19. Majhi, P., Rout, K. K., Nanda, G., & Singh, M. (2021). Long term effects of fertilizer and manure application on productivity, sustainability and soil properties in a rice-rice system on Inceptisols of Eastern India. *Communications in Soil Science and Plant Analysis*, 52(14), 1631-1644.
20. Sharma, K. L., Reddy, K. S., Chary, G. R., Rao, C. S., Singh, D. N., Lal, M & Haindavi, P. (2018). Effect of Long Term Conjunctive Nutrient Management Practices on Soil Quality Indicators and Indices in Oxisol Soils under Rice (*Oryza sativa* L.)-Black gram (*Vigna radiata* L.) Cropping System in Ranchi Region. *Indian Journal of Dryland Agricultural Research and Development*, 33(2), 1-9.