

Original Research Article

Effect of inorganic fertilizers and manures on yield and soil nutrient status of Maize in calcareous *Inceptisol* over five decades of intensive farming in India

Abstract

Long Term Fertilizer Experiment was initiated with an aim to monitor the continuous manuring and fertilization under intensive cropping changes in soil properties with respect to concerning physical, chemical and biological characteristics and crop productivity. Continuous use of inorganic fertilizer along with FYM could increase the soil nutrients such as $\text{KMnO}_4 - \text{N}$, Olsen - P and $\text{NNH}_4\text{OAc} - \text{K}$ over five decades. The experimental soil belongs to sandy clay loam soil (*Vertic Ustropept*) with ten different nutrient management practices are followed and replicated thrice by the adoption of RBD. The results revealed that the use of inorganic fertilizer conjoint with FYM @ 10 t ha⁻¹ significantly increased the kernel yield of maize by 18.1% and 20.3% higher than 150% NPK and 100% NPK plots in sandy clay loam soil. The important indicator for soil fertility concern is SOC and the build-up of SOC was found in NPK+FYM @ 10 t ha⁻¹ treatment recorded from 3.0 g kg⁻¹ (1972) to 7.8 g kg⁻¹ (2023). Soil fertility status of the experiment after 50 years had reported as medium in $\text{KMnO}_4 - \text{N}$, medium to high in Olsen - P, high in $\text{NNH}_4\text{OAc} - \text{K}$ and medium in SOC status in an *Inceptisol*. From the study, hypothesis revealed that, the application of inorganic fertilizer along with FYM @ 10 t ha⁻¹ achieved the a greater yield and better soil health in a sustainable manner.

Key Words: Farmyard manure, SOC, Inorganic fertilizers and Soil Fertility

1. INTRODUCTION

Soil is a vital natural resources and Soil quality is the combined effect of management on most soil properties which influences the crop productivity and sustainability (Sharma *et al.*, 2005). The maintenance of soil health is not only important in increasing the agricultural production but also essential for sustaining the higher productivity of crops (Velu and Subramaniam, 2012). Soil fertility is the most important factor which controls the crop productivity. Soil-Soil-related constraints affecting crop yield such as nutritional disorders can be assessed by evaluating the soil fertility status. Soil testing provides information about the nutrient status of the soil from which fertilizer recommendations can be made for maximizing the crop productivity (Haribhushan Athokpam *et al.*, 2013).

In recent years, intensive farming practices using high-yielding cultivars with imbalanced fertilization have resulted in over mining of native soil nutrients that adversely affects the crop production and soil fertility status (Tandon 1999 and Swarup and Ganeshmurthy, 1998). It is important to maintain the nutrient supply in order to sustain productivity. Since a large amount of nutrients has to be applied to soil in a chemical form which may have an impact on soil properties and soil productivity in the long term. Long Term Fertilizer Experiment was initiated with an aim to monitor the changes in soil properties as a result of the continuous addition of organic manures and inorganic fertilization under intensive cropping with respect to concerning the physical, chemical and microbiological characteristics of the soil in relation to concerning its productivity. The integrated application of organic manures and inorganic fertilizers is the best-recognized strategy of Integrated Nutrient Management to improve the soil fertility and also enhancing the crop yields (Manimaran *et al.*, 2022).

Maize (*Zea mays*) is the most important coarse cereal crop that belongs to the Poaceae family. It is widely used for cattle feed, poultry feed and also for human consumption (Gul *et al.*, 2021). Maize is adapted to a wide range of soil and agro-climatic conditions but is sensitive to water logging hence, it is known for as the Queen of Cereals. It is a more nutrient-exhausting crop. Globally, India ranks fourth in area and seventh in production among maize-growing countries, Maize is grown on 9.2 million hectares in India, it

Comment [PP1]: What next after this aim?

Comment [PP2]: Write in full followed by the abbreviation and this should apply to the whole document, where the Abbreviation/acronyms has been used for the first time in the document

Comment [PP3]: Refer to comment under PP2

Comment [PP4]: Not sure if this is applicable only after this period. You may wish to revisit this statement of fact.

Comment [PP5]: Find a better word that professional

Comment [PP6]: Refer to comment under PP2

Comment [PP7]: Refer to comment under PP2

Comment [PP8]:

Comment [PP9R8]: Avoid using possessive words in the whole document

Comment [PP10]: What was the hypothesis?

Comment [PP11]: How and why is it vital

Comment [PP12]: Any difference between increasing the agricultural production and sustaining higher productivity of crops?

Comment [PP13]: Need for reference

Comment [PP14]: Not only fertilizer I guess even manure. As such, consider rephrasing

Comment [PP15]: Where recent years shall mean which years?

Comment [PP16]: Mining in what way?

Formatted: Highlight

Comment [PP17]: How large?

Comment [PP18]: What is the message here? It does not sound to be complete

Comment [PP19]: Need for reference

accounts for around 4% of global maize area and 2% of total production (DACNET, 2020). In India, Maize is the third most important cereal crop next to Rice and Wheat.

Comment [PP20]: Consider rephrasing

Comment [PP21]: Need for Reference

In this article, an attempt has been made to highlight the Importance of INM practices in maintaining soil health and improving in-crop yield and productivity.

Comment [PP22]: Make reference to comment under PP2

2. MATERIALS AND METHODS

2.1 Experimental site and climate

The study area is located at the Research farm of Tamil Nadu Agricultural University, Coimbatore district of Tamil Nadu, India. The Long Term Fertilizer Experiment (LTFE) was started in 1972 at 11° N latitude, 77° E longitude with an elevation of 426.7 m above MSL with Finger millet- Maize cropping sequence under irrigated condition, soil classified as *Inceptisol* having black calcareous sandy clay loam soil (*vertic ustropept*) belongs to Periyanaickenpalayam soil series. The climate of Coimbatore is tropical characterized by hot and humid summers and cold winters. The NE monsoon majorly contributes to Coimbatore, the average annual precipitation ranges between 550 mm to 900 mm.

Comment [PP23]: Too long and mixed. Consider rephrasing and splitting.

Comment [PP24]: Consider giving statistic on the wether

Comment [PP25]: Which months of the year do you expect this?

2.2 Experimental details

LTFE also includes two crops per year, Finger Millet – Maize cropping sequence raised during summer and kharif seasons; respectively with ten treatment combinations and three replications, the plots are randomized by Randomized Block Design (RBD). The treatment details were T₁- control, T₂-100% N, T₃-100% NP, T₄- 100% NPK, T₅-100% NPK+FYM at 10 t ha⁻¹, T₆-100% NPK+ Zn, T₇- 150% NPK T₈- 50% NPK, T₉-100% NPK+HW and T₁₀-100% NPK (- S). In this study, a Maize hybrid (COHM 6) as a test crop and ten treatments with three replications were taken for investigation.

Comment [PP26]:

Comment [PP27R26]: Why was this picked?

The recommended dose of fertilizer for Maize hybrid were 250, 75 and 75 kg of N, P and K ha⁻¹ respectively. At the time of sowing 50% of RDF N were applied as urea and 100% of RDF of P and K were applied as SSP and MOP respectively for all the treatments except for T₉ (100% NPK – S) DAP can be used as a source. Another 50% of N was applied in two equal splits during knee high stage and pre-pre-tasselling stages of crop growth. For INM (100% NPK + FYM) treatment plots were applied with 10 t ha⁻¹ of FYM for every crop.

Comment [PP28]: Who recommended this?

Comment [PP29]: Please refer to comment under PP2

Comment [PP30]: Refer to comment under PP2

2.3 Soil Sampling, Processing and Analysis

The soil samples were collected from 50 years old Long Term Fertilizer Experiment from the individual treatment plots after the harvest of Finger millet and Maize; respectively at a depth of 0 – 15 cm. The soils were air dried, ground and sieved to pass through 2 mm sieve. The processed soil samples were used for the determination of various soil physical and chemical properties by following standard procedures. Soil texture was determined by the international pipette method. The power of hydrogen (pH) and Electrical Conductivity (EC) were determined in the soil and water suspension in the ratio of 1:2.5 by potentiometric and conductometric methods; respectively (Jackson 1973). Available N in soil was estimated by alkaline KMnO₄ method (Asija and Subbaiah 1956). Available P in soil was estimated by NaHCO₃ extractions and determined colorimetrically (Olsen *et al* 1954). Available K in soil was estimated by using neutral normal Ammonium Acetate extractions followed by Emission spectroscopy (Standford and English 1949). Soil organic carbon was determined by the chromic acid wet digestion method (Walkley and Black 1934).

Comment [PP31]: Be specific on this by even providing reference(s) for easy followup

Formatted: Highlight

Formatted: Highlight

2.4 Statistical Analysis

The recorded data from the field experiment were analysed for their significance (p=0.05) by statistical procedure appropriate for Randomized Block Design outlined by Gomez and Gomez 1985.

Comment [PP32]: Consider using latest references instead of 1934, 1949 and 1950s

Comment [PP33]:

Comment [PP34R33]: How often was the sampling conducted and for how long?

3. RESULTS AND DISCUSSION

3.1 Initial physical, physico- chemical and chemical properties of soil during 1972.

The initial ~~properties of experimental soil was the property of experimental soil was~~ analyzed for various physical, physico-chemical and chemical properties. The pH of the soil was found to be alkaline ~~in nature~~, and the primary nutrient status of the soil ~~were was~~ found to be low in KMnO_4 -N, medium in Olsen-P, high in NNH_4OAc -K and medium in SOC status in a Sandy Clay Loam soil (Table 1).

Table 1. Initial physical, physico-chemical and chemical properties of soil during 1972.

Characteristics of Soil	(1972) Value
Particle size analysis	
Clay (%)	32.6
Silt (%)	11.8
Fine sand (%)	15.1
Coarse Sand (%)	39.4
Soil textural class	Sandy clay loam
Physicochemical and chemical properties	
pH (1:2.5 soil: water suspension)	8.20
Electrical conductivity (dS m^{-1})	0.20
Organic carbon (g kg^{-1})	3.0
Cation exchange capacity [cmol (p+) kg^{-1}]	25.5
Available nitrogen (kg N ha^{-1})	178.0
Available phosphorus (kg ha^{-1})	11.0
Available potassium (kg ha^{-1})	810.0

3.2 Long term fertilization effects on soil physico-chemical properties of Maize crop in an *inceptisol* (post-harvest samples of 115th crop)

3.2.1 Soil reaction (pH) and Electrical Conductivity (dS m^{-1})

The soil ~~reaction~~ pH and EC ~~were~~ changed ~~s~~ slightly from 1972, currently, soil reaction of the research soil was alkaline ~~in nature~~ and pH ranges from 8.07 to 8.40 (Table 3.) specifies that there is no significant influence of pH on soil due to buffering capacity of the soil. The slight changes in the pH was due to ~~the~~ release of organic acids and ~~the~~ continuous addition of inorganic fertilizers and organic manures. Likewise, EC ranges from 0.51 to 0.63 (Table 3.) ~~indicates indicating~~ that there is no significant impact of EC on different fertilization treatments (Malarkodi *et al.*, 2019).

3.3 Long term fertilization effects on soil chemical properties of Maize crop in an *inceptisol* (post-harvest samples of 115th crop)

3.3.1 Soil Organic Carbon SOC (g kg^{-1})

Soil Organic Carbon (SOC) is the important factor which governs ~~the~~ soil fertility (Prabakaran *et al.*, 2023). The continuous application of inorganic fertilizers and organic manures over five decades in a Finger millet – Maize sequence has increased the SOC content from 3.0 g kg^{-1} to 7.8 g kg^{-1} (Nayak *et al.*, 2012). The SOC concentration has increased significantly under different fertilized and unfertilized treatments from control to 150% NPK (4.0 g kg^{-1} to 7.8 g kg^{-1}) (Table 2). The maximum SOC concentration was recorded in 100% NPK + FYM ~~at~~ 10 t ha^{-1} plot at the rate of 7.8 g kg^{-1} followed by 150% NPK at 7.1 g kg^{-1} (Bhattacharyya *et al.*, 2010) and (Srinivasarao *et al.*, 2011). The treatments like 100% NPK > 100% NPK + Zn > 100% NPK + HW are on par with each other and 100% NP & 100% NPK - S are also on par with each other. The minimum SOC concentration was recorded in ~~the~~ control plot at the rate of 4.0 g kg^{-1} .

Comment [PP35]: Since the data is available, it will be very much appreciated if such data was included in the explanation followed by referencing Table 1

Comment [PP36]: Is this one off data or average and if so, say something about it. I am saying this because there is a need for presentation of values in ranges instead of absolute values like in the case of pH and EC

Comment [PP37]: What is the meaning of this? There is need for an opening statement regarding this sub-section 3.2. Explaining what this sub-section is all about. Consider rephrasing the title as it is long and confusing.

Comment [PP38]: Consider rephrasing this sentence by including the values of both pH and EC. The use of graphs could help in showing trend.

Comment [PP39]: How slight was it. Consider including the figures

Comment [PP40]: Not connecting well why are you saying its not significant, basing on what?

Comment [PP41]: Please make reference to comment PP35 as the titles 3.2 and 3.3 could be confusing. Consider rephrasing this part as its confusing, Consider the main issue of the presentation i.e. what is it that you want the reader to understand

3.3.2 Soil Available Nitrogen (kg ha⁻¹)

Nitrogen is one of the major plant nutrients which imparts green colour to plants and encourages vegetative growth. The continuous application of inorganic fertilizers and organic manures has significantly influenced the available N status in the soil under different management treatments and the available N content in the Experimental soil was found to be medium and ranged from 260 to 408 kg ha⁻¹ (Table 3). The maximum available N content was recorded in (100% NPK + FYM @ 10 t ha⁻¹) plot of 408 kg ha⁻¹ followed by 150% NPK plot of 378kg ha⁻¹ (Bairwa *et al.*, 2021). However, the treatments like 100% NPK > 100% NPK +Zn >100% NPK +HW are on par with each other. The 100% NPK plots showed a that declined availability of N content than to other treatments. The minimum available N content was recorded in control plots of 260kg ha⁻¹. The combined application of inorganic nitrogenous fertilizer along with FYM has increased the mineralization of the organic form of N to Inorganic forms.

3.3.3 Soil Available Phosphorous (kg ha⁻¹)

Phosphorous is also a primary macronutrient which plays an important role in energy storage and transfer. The available P content in the research area was found to be medium to high under different treatments and the values ranged between 16.13 to 29.32 kg ha⁻¹ (Table 3). The maximum available P content was recorded in (100% NPK + FYM @ 10 t ha⁻¹) plot of 29.32 kg ha⁻¹ followed by 150% NPK plot of 28.27 kg ha⁻¹ (Kalaiselvi *et al.*, 2021) and (Mitran *et al.*, 2016). Other treatments like 100% NP > 100% NPK >100% NPK +Zn are on par with each other. The minimum available P content was recorded in control plots of 16.13 kg ha⁻¹. The soil pH is a the most important factor which influences the P fixation in the soil. Under acidic conditions (pH < 6), P is fixed as AlPO₄ and FePO₄ and in alkaline (pH > 8) conditions fixed as a CaPO₄. The greater P content in the INM treatment plot was due to the presence of Organic Matter that leads to the formation of organophosphate compounds that are more soluble and also increase the quantity of organic P to inorganic P.

3.3.4 Soil Available Potassium (kg ha⁻¹)

Potassium is the third most important primary nutrient absorbed by the plants in larger amounts than the other nutrients. The available K content in the research soil was found to be high since from the start of the experiment in (1972). The use of high-high-yield cultivars and inorganic fertilizers has significantly exhausted the soil K under different treatments and ranges between 236 to 642 kg ha⁻¹ (Table 3). The maximum available K content was recorded in (100% NPK + FYM @ 10 t ha⁻¹) plot of 642 kg ha⁻¹ followed by 150% NPK plot of 564 kg ha⁻¹ (Mazumdar *et al.*, 2014). Other treatments like 100% NPK > 100% NPK +Zn >100% NPK +HW are observed to be at on par with each other. The minimum available P content was recorded in control plots of 236 kg ha⁻¹. The application of NPK with FYM has increased the K availability in soil due to the release of fixed K ions in the exchangeable sites (Muneshwar *et al.*, 2014).

Comment [PP42]: Provide reference

Comment [PP43]: How large and any reference to this fact?

Comment [PP44]: Provide the value

Table 2. Long term fertilization effects on soil Physico- chemical and chemical properties of Maize (kg ha⁻¹) in an Inceptisol (Initial samples of 115th crop)

Treatments	pH	EC dSm ⁻¹	SOC g kg ⁻¹	N Kg ha ⁻¹	P Kg ha ⁻¹	K Kg ha ⁻¹
Control	8.09	0.52	0.40	227	11.09	259
50 % NPK	8.21	0.58	0.54	276	16.76	318
100% N	8.25	0.62	0.51	285	12.98	295
100% NP	8.46	0.58	0.60	289	18.94	307
100% NPK	8.23	0.57	0.63	321	19.08	476
100% NPK+ FYM	8.30	0.65	0.78	357	21.99	538
100% NPK +Zn	8.22	0.59	0.66	309	18.67	482
100% NPK +HW	8.15	0.56	0.64	302	18.54	475
100% NPK- S	8.36	0.63	0.61	293	17.91	434

Comment [PP45]: Consider rephrasing the titles of Tables 2 and 3 as they are confusing. Consider the main issues in the Tables I guess is to do with Initial samples of 115th Crop and post harvest samples. Try to think in those lines and consider revisit the titles as they are long and confusing

150% NPK	8.18	0.62	0.71	336	20.76	502
SEd	1.222	0.0097	0.0148	7.3533	0.4369	10.2525
CD (P= 0.05)	2.5673	0.0205	0.0311	15.449	0.918	21.54

Table 3. Long term fertilization effects on soil Physico–chemical and chemical properties of Maize (kg ha⁻¹) in an *Inceptisol* (post-harvest samples of 115th crop)

Treatments	pH	EC	N Kg ha ⁻¹	P Kg ha ⁻¹	K Kg ha ⁻¹
control	8.07	0.51	260.3	16.13	236
50 % NPK	8.20	0.57	301.3	21.27	376
100% N	8.25	0.63	314.3	19.53	312
100% NP	8.40	0.59	339.7	24.77	327
100% NPK	8.23	0.57	363.7	25.30	558
100% NPK+ FYM	8.27	0.67	408.7	29.32	642
100% NPK +Zn	8.21	0.60	366.7	25.50	497
100% NPK +HW	8.16	0.53	356.0	24.43	508
100% NPK- S	8.34	0.62	338.3	24.25	478
150% NPK	8.15	0.63	378.0	28.27	564
SEd	0.1977	0.0154	6.6191	0.4255	11.4556
CD (P=0.05)	0.4154	0.03	13.095	0.8939	24.068

3.4 Long term fertilization effects on Kernel and Stover yields of Maize (kg ha⁻¹) in an *Inceptisol*

The continuous application of inorganic fertilizers and organic manures caused significant variation in kernel and straw yield of Maize in an *inceptisol* (Fig.1). The maximum kernel and Stover yield of Maize was recorded in the 100%NPK + FYM at 10 t ha⁻¹ plot of 6898 kg ha⁻¹ and 9356 kg ha⁻¹, respectively followed by 150% NPK plot 5643 kg ha⁻¹ and 8978 kg ha⁻¹ (Choudhary *et al.*, 2021). The minimum kernel and straw yield of Maize was recorded in the control plot of 2872 kg ha⁻¹ and 4317 kg ha⁻¹, respectively. The yield increase in 100% NPK + FYM at 10 t ha⁻¹ plot obtained was 18.1% and 20.3% higher than 150% NPK and 100% NPK plots, respectively. This might have been due to the conjoint application of NPK + FYM that has significantly improved the soil physical, chemical and biological properties of soil leading to the increase in the yield and productivity of the soil in a sustained manner (Majhi *et al.*, 2019) and (Sharma *et al.*, 2018).




Fig.1 Long term fertilization effects on kernel and Straw yields of Maize (kg ha⁻¹) in an Inceptisol

4. CONCLUSION

The present study revealed that the application of a recommended dose of fertilizer along with the organic manure (i.e. 100% NPK + FYM at 10 ha⁻¹) was considered to be effective among different nutrient management treatments under the sandy clay loam of the Maize crop. Balanced application of nutrients would improve the physical, chemical and biological properties of the soil as well as maintaining the soil fertility and also enhance the productivity of Maize crops under irrigated conditions. With this result, our hypothesis revealed that, the application of inorganic fertilizer along with FYM at 10 t ha⁻¹ achieved the a greater yield and better soil fertility in a sustainable manner.

Comment [PP46]: Not sure if this experiment was on irrigation/irrigated condition

Comment [PP47]: Please refer to comment under PP10

REFERENCE

- Sharma, K. L., Mandal, U. K., Srinivas, K., Vittal, K. P. R., Mandal, B., Grace, J. K., & Ramesh, V. (2005). Long-term soil management effects on crop yields and soil quality in a dryland Alfisol. *Soil and Tillage Research*, 83(2), 246-259.
- Velu, V., & Subramaniam, P. (2012). Soil health and its sustenance through enriched compost. In *National seminar on recycling of solid wastes through composting held during March 8th and 9th at the Department of Soil and Environment, Agricultural College and Research Institute, Madurai* (pp. 5-11).
- Haribhushan, A., Shabir, H. W., David, K., Herojit, S. A., Jyotsna, N., Deepak, K., ... & Lamalakshmi, D. (2013). Soil macro-and micro-nutrient status of Senapati district, Manipur (India). *African Journal of Agricultural Research*, 8(39), 4932-4936.
- Tandon, H. L. S. (1995). Micronutrients in soils, crops and fertilisers-a sourcebook-cum-directory (1995)-. *Micronutrients in soils, crops and fertilisers-a sourcebook-cum-directory (1995)*.
- Swarup, A., & Ganeshamurthy, A. N. (1998). Emerging Nutrient Deficiencies under Intensive Cropping Systems and Remedial Measures for Sustainable High Productivity. *Fertiliser news*, 43(7).
- Manimaran, G., Jayanthi, D., Janaki, P., Amirtham, D., & Gokila, B. (2022). Long Term Impact of Fertilization and Intensive Cropping on Maize Yield and Soil Nutrient Availability under Sandy Clay Loam Soil (Inceptisol). *International Journal of Plant & Soil Science*, 34(20), 795-801.
- Gul, H., Rahman, S., Shahzad, A., Gul, S., Qian, M., Xiao, Q., & Liu, Z. (2021). Maize (*Zea mays* L.) productivity in response to nitrogen management in Pakistan. *American Journal of Plant Sciences*, 12(8), 1173-1179.

Comment [PP48]: Be specific in the year of reference as its 1999 in the document. Second paragraph of the Introduction section

8. Malarkodi, M., Elayarajan, M., Arulmozhiselvan, K., & Gokila, B. (2019). Long-term impact of fertilizers and manures on crop productivity and soil fertility in an alfisol. *The Pharma Innovation Journal*, 8(7), 252-256.
9. Prabakaran, S., Kaleeswari, R. K., Backiyavathy, M. R., Jagadeeswaran, R., Selvi, R. G., & Bama, K. S. (2023). Estimation of soil carbon pools under major cropping systems of Mayiladuthurai district of Cauvery Delta Zone, Tamil Nadu, India. *Journal of Applied and Natural Science*, 15(2), 802-810.
10. Nayak, A. K., Gangwar, B., Shukla, A. K., Mazumdar, S. P., Kumar, A., Raja, R., ... & Mohan, U. (2012). Long-term effect of different integrated nutrient management on soil organic carbon and its fractions and sustainability of rice-wheat system in Indo Gangetic Plains of India. *Field Crops Research*, 127, 129-139.
11. Bhattacharyya, R., Kundu, S., Srivastva, A. K., Gupta, H. S., Prakash, V., & Bhatt, J. C. (2011). Long term fertilization effects on soil organic carbon pools in a sandy loam soil of the Indian sub-Himalayas. *Plant and soil*, 341, 109-124.
12. Srinivasarao, C. H., Venkateswarlu, B., Lal, R., Singh, A. K., Kundu, S., Vittal, K. P. R & Patel, M. M. (2014). Long-term manuring and fertilizer effects on depletion of soil organic carbon stocks under pearl millet-cluster bean-castor rotation in Western India. *Land Degradation & Development*, 25(2), 173-183.
13. Bairwa, J., Dwivedi, B. S., Rawat, A., Thakur, R. K., & Mahawar, N. (2021). Long-term effect of nutrient management on soil microbial properties and nitrogen fixation in a vertisol under soybean-wheat cropping sequence. *Journal of the Indian Society of Soil Science*, 69(2), 171-178.
14. Kalaiselvi, K., Jayanthi, D., Santhy, P., Gnanachitra, M., & Gokila, B. (2021). Effect of long term fertilization on phosphorus dynamics in root zone environment under finger millet-Maize cropping sequence. *Journal of Applied and Natural Science*, 13(4), 1383-1389.
15. Mitran, T., Mani, P. K., Basak, N., Mazumder, D., & Roy, M. (2016). Long-term manuring and fertilization influence soil inorganic phosphorus transformation vis-a-vis rice yield in a rice-wheat cropping system. *Archives of Agronomy and Soil Science*, 62(1), 1-18.
16. Mazumdar, S. P., Kundu, D. K., Ghosh, D., Saha, A. R., Majumdar, B., & Ghorai, A. K. (2014). Effect of long-term application of inorganic fertilizers and organic manure on yield, potassium uptake and distribution of potassium fractions in the new Gangetic alluvial soil under jute-rice-wheat cropping system. *International Journal of Agriculture and Food Science Technology*, 5(4), 297-306.
17. Singh, M., Ram, S., Wanjari, R. H., & Sharma, P. (2014). Balance and forms of potassium under rice-wheat system in a 40-year-old long-term experiment on Mollisols of Pantnagar. *Journal of the Indian Society of Soil Science*, 62(1), 38-44.
18. Choudhary, M., Meena, V. S., Panday, S. C., Mondal, T., Yadav, R. P., Mishra, P. K. & Pattanayak, A. (2021). Long-term effects of organic manure and inorganic fertilization on biological soil quality indicators of soybean-wheat rotation in the Indian mid-Himalaya. *Applied Soil Ecology*, 157, 103754.
19. Majhi, P., Rout, K. K., Nanda, G., & Singh, M. (2021). Long term effects of fertilizer and manure application on productivity, sustainability and soil properties in a rice-rice system on Inceptisols of Eastern India. *Communications in Soil Science and Plant Analysis*, 52(14), 1631-1644.
20. Sharma, K. L., Reddy, K. S., Chary, G. R., Rao, C. S., Singh, D. N., Lal, M & Haindavi, P. (2018). Effect of Long Term Conjunctive Nutrient Management Practices on Soil Quality Indicators and Indices in Oxisol Soils under Rice (*Oryza sativa* L.)-Black gram (*Vigna radiata* L.) Cropping System in Ranchi Region. *Indian Journal of Dryland Agricultural Research and Development*, 33(2), 1-9.

Comment [PP49]: Review the year as it is 2011 in the document

Comment [PP50]: Review the year as it is 2019 in the document

Comment [PP51]: Be specific in the year of reference as in the document is indicated 2005. First paragraph in the Introduction section