

EFFECT OF VARIOUS NATURAL PRESERVATIVES & NATURAL COLOURING MATERIALS ON THE QUALITY OF BER (*Ziziphus mauritiana*) CANDY

ABSTRACT

An experiment was carried out in the Post Harvest Technology laboratory, Department of Horticulture, Sam Higginbottom University of Agriculture, Technology and Sciences, Allahabad (Uttar Pradesh) during the year 2022-2023. The experiment consisted of 10 treatments and 1 control. The investigation laid out in Completely Randomized Design with three replications. Ber candy was stored for 80 days at ambient temperature. From storage studies it was revealed that T3 (jaggery + ginger) performed best in terms of quality (taste, flavour, overall acceptability) and storage. With a fairly high Benefit-cost ratio of 2.34. On the basis of results it is concluded that treatment T3 be used for commercialization. The results indicated that the quality observations and sensory evaluation are affected by various treatments.

Key words :- Ber candy, quality, commercialization.

INTRODUCTION

Ber /Indian jujube /Indian plum / Chinese date /Chinese apple /dunks

Scientific name : *Ziziphus mauritiana*

Family : Rhamnaceae

Promising varieties : Umran, Kathapal

After harvest, fruits are liable to accelerated physiological, chemical, and microbial processes that invariably lead to deterioration and loss of wholesomeness.

It is then necessary to institute some measure of processing such as reduction in moisture content, denaturation of endogenous enzymes and microorganisms, or packaging in order to curtail perishability.

Value addition to food products has assumed vital importance in our country due to diversity in socio-economic conditions, industrial growth, urbanization and globalization.

It is not merely to satisfy producers and processors by way of higher monetary return but also with better taste and nutrition.

Value is added by changing their form, colour and other such methods to increase the shelf life of perishables.

Importance of Post-harvest technology lies in the fact that it has the capability to meet food requirement of growing population by eliminating losses making more nutritive food items from raw commodities by proper processing and fortification.

India is the world's second largest producer of fruits.

It has potential to grow all types of temperate, sub-tropical and tropical fruits because of varied agro-climatic diversity.

The total production of fruits is over 45 million tones.

The losses are estimated to the extent of 20 -30 per cent due to lack of proper harvesting, processing and storage facilities, which is valued at Rs. 230 billion.

This study aimed to evaluate the effects of different food additives on the quality of ber candy and to estimate the economics of various treatments.

MATERIALS AND METHODS

The present experiment was conducted with the objective of assessing the flavour, taste, texture, shelf life and economics for checking "Effect of various natural preservatives and natural colouring material on the quality of ber candy". Different treatment combinations were used (Table1). The details of the

materials and methods used are given below.

Prerequisites :-

Extraction of colour from beetroot, carrot and spinach.

Add 2 cups of grated/cut vegetable to 2 cups of water. Mix it well, close and cook for 10 minutes. Once done let it cool. Blend it into a smooth paste. Filter the paste and your colour is ready.

Ginger extract

The ginger is grated and the pulp is squeezed to extract the juice.

Procedure for preparation of ber candy

The sound fruits are washed under tap water to remove dirt or any foreign particle on the skin. It is then put in salt water for about 15 minutes and washed again with clean water. The stem of ber fruits is removed manually by using a sharp stainless steel knife. The edible fruit portion is cut in two halves and the seed is removed. The cut fruit is boiled for 2-5 minutes or till slightly soft. The sugar/jaggery syrup (30 degree brix) as per treatment + lemon juice is then poured on the fruits which are kept submerged in it for 4 days. The density of the syrup is gradually raised to 75 degree brix in a course of 10-12 days.

| | |
|----------------------|-------|
| 5 th day | 50 °B |
| 10 th day | 75 °B |

The spinach extract, beetroot extract, ginger extract, carrot extract, cardamom are added according to the treatments on the 10th day and the fruit is kept in it for 24 hours. The syrup is drained. The fruits are dried in shade till the moisture content in the fruit is 20 %. Another test to check if the fruit is dry enough is that there should be no visible moisture when the candied fruit is cut open or you should not be able to squeeze any moisture from the fruit.

Statistical Analysis

The data recorded during the course of investigation were subjected to statistical analysis as per method of analysis of variance (ANOVA) for Completely Randomized Design (CRD) by Fisher and Yates (1963). The significance and non-significance of the treatment effect were judged with the help of

'f' value (variance ratio) that was compared with the table value at 0.05% level of significance.

Table 1: Treatment combinations

| Treatments | Combinations | | | |
|------------|--------------|---------|------------|----------|
| T0 | - | - | - | - |
| T1 | ber | jaggery | lime juice | Spinach |
| T2 | ber | jaggery | lime juice | Beetroot |
| T3 | ber | jaggery | lime juice | Ginger |
| T4 | ber | jaggery | lime juice | Carrot |
| T5 | ber | jaggery | lime juice | cardamom |
| T6 | ber | sugar | lime juice | Spinach |
| T7 | ber | sugar | lime juice | Beetroot |
| T8 | ber | sugar | lime juice | Ginger |
| T9 | ber | sugar | lime juice | Carrot |
| T10 | ber | sugar | lime juice | cardamom |

Chemical analysis methods

(1) Total soluble solids (°B)

The percentage of total soluble solids was determined by using ERMA hand refractometer by placing a drop of the filtered juice on the prism of the refractometer and observing the coincidence of shadow of the sample with the reading on the scale and expressed as °B.

(2) Acidity (%)

"10 ml of homogenized sample was taken and made up to 100 ml volume in a volumetric flask. The contents were filtered through Whatman No.1 filter paper. An aliquot of 10 ml was taken for titration against 0.1N NaOH using phenolphthalein as an indicator. The turn of aliquot to light pink colour which persists for 15 seconds was considered as an end point. The titratable acidity was estimated in terms of percent citric acid" (Ranganna, 1986).

Factor for acidity=1ml of 0.1N NaOH = 0.0064g of citric acid.

Total acidity % = $\text{Titrate} \times \text{normality} \times 0.06404 \times 100$

Wt. Or volume of sample

(3) Moisture content

"Moisture content was determined by using hot air oven drying method 10gm of sample was taken in pre-weighed empty Petri plate and dried hot air oven at 100 °C till constant weights were obtained. Plates were cooled in desiccator. The moisture content was calculated by using formula". (Ranganna 1986).

$$\% \text{ Moisture} = \frac{\text{Initial weight} - \text{Final weight}}{\text{Weight of sample}} \times 100$$

(4) pH Value:

The sample was soaked in distill water till it soften and ground along with the little amount of distilled water. Then the pH was noted with the help of an electronic pH meter.

(5) Organoleptic evaluation:

The organoleptic evaluation for accessing the taste, color, flavor and texture of the samples were conducted by panel of 10 judges who scored on a 9 points hedonic (Srilakshmi 2007) scale as mentioned below.

RESULT AND DISCUSSION

According to the results recorded in Tables 3 & 4

Total soluble solids (°B) of ber candy

Total soluble solids of Ber candy were observed to be increased gradually up till the end of experiment under ambient storage conditions. At the beginning of storage there was significant difference amongst various treatments. At 80 days the minimum TSS was recorded in T0 (control) and maximum in T5 (jaggery + cardamom). Similar results was reported by Tandon *et al.*, (2008).

Acidity (%) of ber candy

The acidity(%) of Ber candy was observed to decrease upto the end of the experiment under ambient storage co re to be decreased gradually up to the end of the experiment under ambient storage conditions. At the beginning of storage there was significant difference amongst various treatments. At 80 days after storage the minimum acidity was recorded in T4 (jaggery + carrot) and maximum in T0 (control). Similar results was reported by Neelesh (2014) in papaya candy.

Physiological loss in weight (%) of ber candy

The moisture content (%) of ber candy was observed to gradually decrease up to the end of the experiment under ambient storage conditions. At 80 days the minimum physiological loss in weight was recorded in T0 (control) and maximum in T2 (jaggery + beetroot). Similar results was reported by **Daisy and Gehlot (2006)** in Aonla preserve.

pH of ber candy

The pH of ber candy was observed to decrease gradually up to the end of the experiment under ambient storage conditions. At the beginning of storage there was significant difference amongst various treatments maximum pH 5.03 was observed in T3. At 80 days after storage. The minimum pH was recorded in T0 (control) and maximum in T4 (jaggery + carrot) and T9 (sugar + carrot). Similar results was reported by **Jain et al. (2004)** in papaya cubes.

Ascorbic acid (mg/100 gms) of ber candy

Ascorbic acid of Ber candy were observed to be decreased gradually up till the end of experiment under ambient storage conditions. At the beginning of storage there was significant difference amongst various treatments. At 80 days after the minimum ascorbic acid was recorded in T9 (sugar + carrot) and maximum in T0 (control).

Reducing sugars (%) of ber candy

Reducing sugars of Ber candy were observed to be increased gradually up till the end of experiment under ambient storage conditions. At the beginning of storage there was significant difference amongst various treatments. At 80 days after the minimum reducing sugars were recorded in T6 (control) and maximum in T2 (jaggery + beetroot).

Total sugars (%) of ber candy

Total sugars of Ber candy were observed to be increased gradually up till the end of experiment under ambient storage conditions. At the beginning of storage there was significant difference amongst various treatments. At 80 days after The minimum total sugars were recorded in T0 (control) and maximum in T4 (jaggery + carrot).

Non-reducing sugars (%) of ber candy

Non-reducing sugars of Ber candy were observed to be decreased gradually up till the end of experiment under ambient storage conditions. At the beginning of storage there was significant difference amongst various treatments. At 80 days

the minimum non-reducing sugars were recorded in T9 (control) and maximum in T1 (jaggery + spinach) and T6 (sugar + spinach).

Colour (sensory score) of ber candy

The data are showing significant difference at 80 days after storage organoleptic score for colour of the ber candy showed a gradual decrease in all the treatments up to the end of experiment. A minimum appealing colour was recorded in T0 (control) and maximum appealing colour was recorded in T7 (sugar + beetroot) at 80 days after storage and findings agreed with the **singh et al., (2012)** in ber candy.

Taste of ber candy

By the result of sensory score for texture the minimum taste was recorded in T0 (control) and maximum in T3 (jaggery + ginger), T5 (jaggery + cardamom). The decreasing trend was observed for taste with increase storage period. This might be due to degradations of volatile substance and flavour constituent. Similar results were found by **Kannan and Susbeela (2002)** in ber candy.

Flavour of ber candy

By the result of sensory score for taste, the minimum flavour was recorded in T0 (control) and maximum in T3 (jaggery + ginger). The decreasing trend was observed for flavour with increase storage period. This might be due to degradation of volatile substance and flavour constituents. Score of flavour of candy was observed to be decreased continuously up to the end of experiment under ambient storage conditions. Similar results reported by **Hasanuzzaman (2014)** in tomato candy.

Overall acceptability of ber candy The result of sensory score for overall acceptability of ber candy, The minimum overall acceptability was recorded in T0 (control) and maximum in T3 (jaggery + ginger). There was significant differences between the treatments for taste, colour, and texture. Similar results observed by **Hiremanth and Rokhade (2006)** in sapota candy. The overall acceptability of mixed fruit leather is dependent on colour, texture, favour and taste rating of the product. The decrease in overall acceptability score may be due to absorption of atmospheric moisture, dilution of sugars and changes in acidity, oxidation of ascorbic acid as well as changes in biochemical constituents of candy. The similar observations were recorded by **Sharma (2013)** in apple candy.

Economics of preparation of different treatments

It is evident from the treatment details that the highest benefit cost ratio was recorded in T0 (control). Finding detail (benefit cost ratio in descending order) T0(2.65) > T2 & T3(2.34) > T4(2.33) > T1 & T7(2.31) > T8(2.29) > T9(2.27) > T5(2.20) > T6 & T10(2.18) as shown in **Table 2**.

Table 2. Economics of different treatments of ber candy.

| Treatments | Gross return(Rs/kg) | Net profit (Rs/kg) | Benefit Cost ratio |
|------------|---------------------|--------------------|--------------------|
| T0 | 900 | 160.63 | 2.65 |
| T1 | 1130 | 160.26 | 2.31 |
| T2 | 1190 | 171.75 | 2.34 |
| T3 | 1170 | 171 | 2.34 |
| T4 | 1150 | 166.56 | 2.33 |
| T5 | 1200 | 154.7 | 2.20 |
| T6 | 1000 | 141.37 | 2.18 |
| T7 | 1100 | 152.85 | 2.31 |
| T8 | 1070 | 152.11 | 2.29 |
| T9 | 1050 | 147.67 | 2.27 |
| T10 | 1120 | 135.81 | 2.18 |

Table 3. Means of readings taken at initial, 20 days, 40 days, 60 days and 80 days of the following parameters during storage at ambient conditions.

| Treatments | TSS °B | Acidity % | Physiological loss in weight % | pH | Ascorbic acid mg/100gms | Reducing sugars % | Total sugars % | Non- reducing sugars % |
|------------|-----------|--------------|---|------|-------------------------------|-------------------------|----------------------|---------------------------------|
| T0 | 25.56 | 1.34 | 17.16 | 3.62 | 64.2 | 16.02 | 21.92 | 5.9 |
| T1 | 70.21 | 0.82 | 21.11 | 4.75 | 19.78 | 37.82 | 83.58 | 45.76 |
| T2 | 70.55 | 0.69 | 21.15 | 4.95 | 19.36 | 38.46 | 83.92 | 45.46 |
| T3 | 69.5 | 1.12 | 21.13 | 4.09 | 19.64 | 37.98 | 83.52 | 45.54 |
| T4 | 70.36 | 0.57 | 20.97 | 4.96 | 19.38 | 38.14 | 83.4 | 45.26 |
| T5 | 71.18 | 0.84 | 20.99 | 4.41 | 19.46 | 38.1 | 83.68 | 45.58 |
| T6 | 70.48 | 0.81 | 20.26 | 4.74 | 19.74 | 33.42 | 79.18 | 45.76 |
| T7 | 69.22 | 0.68 | 20.31 | 4.95 | 19.3 | 34.06 | 79.52 | 45.46 |
| T8 | 69.01 | 1.06 | 20.28 | 4.10 | 19.4 | 33.58 | 79.12 | 45.54 |
| T9 | 69.81 | 0.59 | 20.36 | 4.96 | 19 | 33.74 | 79 | 45.26 |
| T10 | 70.89 | 0.82 | 20.40 | 4.40 | 19.34 | 33.7 | 79.22 | 45.52 |

Table 4. Means of readings taken at initial, 20 days, 40 days, 60 days and 80 days of the following parameters during storage at ambient conditions.

| Treatments | Colour | Taste | Flavour | Overall acceptability |
|------------|--------|-------|---------|--------------------------|
| T0 | 6.2 | 5.6 | 5.6 | 4.6 |
| T1 | 8.2 | 7.6 | 7.6 | 7.6 |
| T2 | 8.2 | 7.6 | 7.6 | 7.6 |
| T3 | 7.2 | 8.6 | 8.8 | 8.8 |
| T4 | 8.2 | 7.6 | 7.6 | 7.6 |
| T5 | 7.2 | 8.6 | 8.6 | 8.6 |
| T6 | 8.2 | 6.6 | 7.6 | 6.6 |
| T7 | 8.6 | 6.6 | 7.6 | 6.8 |
| T8 | 7.2 | 7.6 | 8.6 | 7.6 |
| T9 | 8.2 | 6.6 | 7.6 | 6.6 |
| T10 | 7.2 | 6.6 | 8.6 | 6.6 |

Conclusion

From the present investigation, it has been concluded that T3 (jaggery + ginger) performed best in terms of quality with a TSS of 69.5, the highest overall acceptability of 8.8 and a shelf life of more than 6 months. The highest Benefit-cost ratio was found in T0 (control) i.e. 2.65.

References

- Abbas M.F., A-Niami JH. and A-Ami R.F. (1988).** Some physiological characteristics of fruits of jujube (*Ziziphus spina-christi* L Willd.) at different stages of maturity. *Journal of Horticultural Science* 63:337-339.
- Agarwal, P. and Sandhu, K.S. (2006).** "Utilization of kinnow waste in value added products", *Beverage and Food World.*, 33 (5): 28
- Alhsan, H, Rehman, W., Wani, S. M., Dar, B. N., Dalal, M. R. and Malik, A.R. (2008).** Influence of potassium metabisulphite pre-treatment, osmotic dip and packaging materials on dehydration and some chemical properties of apple rings. *Applied Biological Research*, 10: 31-35.
- Alam S. and Singh (2005).** A Process for dehydration aonla powder. Abstract presented in Convention of Indian Society of Agriculture Engineers. 39: 222.
- Ambri -Rodriguez, S.L., Islas-Hernandez, J.J., Agama-Acevedo, E., Tover, J. and Bello-Perez, LIA., (2008).** "Characterization of a fibre-rich powder prepared by liquefaction of unripe banana flour" *Food Chemistry*, 107 (4): 1515-1521.
- Ames, J.M. (2003).** Browning. *Encyclopedia of Food Science and Nutrition* (2nd).
- Elsevier Science Ltd. UK. pp. 665-672.**
- Anonymous (2015-16).** Annual Report, Directorate of Economics and Statistics. New Delhi.
- Anusuya, A. G., Aswathy, P. S., Kousika, S., Alagammai, S. and Priyadarshini, P. (2006).** Study of osmotic dehydration of grapes. Paper presented in 18th Indian Convention of Food Scientists and Technologists, held at Hyderabad in November, pp-67.
- Ari S., Anju K., Dhiman, Rakesh Sharma and Manisha Kaushal (2015).** Standardization of Pre-treatments for the Development of Intermediate moisture Food Products from Papaya (*Carica papaya* L.). *Intl. J. Food. Ferment. Technol.* 6(1): 143-149.
- Ashuqullah (2017).** Evaluation of sweet potato (*pomoea balatas* (L.) Lam) varieties and pre-treatments for dehydration into flour. M.Sc. Thesis, pp:42 Naseri Agricultural University, Navsari, Gujarat, India.
- Bal, J. S. (1982).** A study on biochemical changes during room and refrigerator storage of ber. *Progressive Horticulture*. 14:
- Bhuiyan, M. H. R., Shams-Ud-Din, M. and Isam, 158-161.**
- M, N. (2012).** Development of Functional Beverage Based on Taste Preference. *Journal of Environmental Science and Natural Resources*, 5(1): 83-87.
- Bose, TK., Mitra, S.K. and Sanyal, D. (2001).** *Fruits: tropical and subtropical traits*, 3rd Revised Edition, Naya Udyog Publication, Calcutta.
- Chandu and Prasad M. (2006).** Development of guava candies, *Journal of Food Science and Technology* 43: 210- 212.
- Dar B. N., Ahsan H., Wani S. M. and Dalal M. R. (2011).** Effect of CaCl₂, Citric Acid and Storage Period on Physico-Chemical Characteristics of Cherry Candy. *Journal of Food Science and Engineering* 154-160.
- Neelesh C., Singh P.S., Singh B.R., Singh S. (2014).** Effect of different citric acids levels on quality papaya candy prepared by various methods. *paripex Indian journal of research* 3(9): 2250-1991.
- Chavan, U.D., Prabhukhanolkar and panwar, V.D. (2010).** Preparation of osmotic Dehydration ripe banana slices. *Journal of food sciences and technology*. 47(4):380-386.
- Divya A.R., Jayashree S. and Basavarajapa B. (2012).** Effect of storage methods on the nutritional quality of sapota candy. *Asian J. Dairy & Food Res.* 33(2): 104-108.
- Daisy and Gehlot, R. (2006).** Physical and biochemical differences in fresh aonla fruits and preserve of cvs. Banarasi and Chakaiya. *Haryana Journal of Horticultural Sciences* 35 (1): 57-59.
- Devaraju, K.R., Rokhade, A.K., Patil, P.B. and Jagadeesh, S.I. (2003).** Influence of various dehydration treatments on ber slices. *Indian Food Packer* 57(5): 63-65.
- Dhingra, D. Kadam, D. M. Singh, J. and Patil, R. T. (2013).** Osmotic dehydration of pineapple with sucrose: mass transfer kinetics. *J. Agril. Eng.*, 50 (1): 14-18.
- Goyal R.K., Patil R. T., Kingsly A, R. P., Walia H. and Kumar P. (2008).** Status of post-harvest technology of aonla in india - A Review. *American journal of food technology*. 3(1): 13-23.
- Guiamba LR., Svanberg U. (2016).** Effects of blanching, acidification or addition of EDTA on

vitamin C and B-carotene stability during mango puree preparation. *Food Sci. Nutr.*, 4, 706-715.

Gupta O. P. (1983). Delicious candy from ber fruits. *Indian Horticulture*, April- June 25-27.

Gupta O.P., Kainsa R.L., and Chauhan K.s, (1980). Postharvest studies on ber fruits (*Ziziphus mauritiana* Lamk.). Preparation of candy. Haryana Agricultural University. *Journal of Research* 10:163.

Hasanuzzaman, M., Kamruzzaman, M., Islam M.M., Khanom S.A.A., Rahman

M.M., Lisa, L.A. and Paul D.K. (2014). A Study on Tomato Candy prepared by Dehydration Technique Using Different Sugar Solutions. *Food Nutr Sci.*, 5: 1261-1271.

Heredia-Leon J.C. Talamas-Abbud R, Mendoza-Guzman V., SolisMartinez F. CastroJ.J., Barnard J., Quintero-Ramos A. (2004). Structural and physical properties of dried Anaheim chilli peppers modified by low temperature blanching. *J. Sci. Food Agric.*, 84: 5965.

Jain, S.K., Verma, R.C, Mathur, A.N. and Murdia, L.K. (2004). Studies on osmotic dehydration of papaya cubes. *Journal of interacademia*. 8(2): 221-229.

Jothi J.S., Monirul Islam, M.D., Serajul Islam, M.D., Rahman R.T. and ShireenAkther (2014). Development and Shelf-Life Prediction of Pineapple (*Ananas comosus*) Preserve and Candy. *International Journal of Innovation and Scientific Research* 10(1): 77-85.

Kaikadi, M.A, Chavan, U.D. and Adsule, R.N. (2016). *Studies on preparation and shelf-life of ber candy", 33 (8): 49-50.

Kakali B., Chakraborty C., Bhattacharyya S. (2014). Fortification of mango peel and kernel powder in cookies formulation. *J. Academia Industria! Research.* 2:161-164. naleemullah, S., Kailappan, R. and Varadharaju, N. (2002). Studies on the Osmotic air drying characteristics of papaya cubes. *J. Food Sci. Technol.* 39(1): 82-84.

Kamble A.K., Kalbande S.R. and Gadge S.R. (2011). Solar drying system for energy conservation applied solar energy. *Applied Solar Energy (Springer)*, 47(2): 124-133.

Kannan, S. and Susheela, T.A. (2002). Studies on the syruling and drying methods of ber (*Zizyphus Mauritiana* Lamk) candy. *Beverage & Food World*, 29(2): 39-41.

Karunaratna, E.J.C.N. and Rathnayaka, R.M.U.S.K. (2012). Influence of the calcium on microbial stability and texture of osmotic dehydrated pineapple slices. *The journal of Agricultural sciences.* 7(1): 1-14.

Khurdiya, D.S. and Singh, R.N. (1975). Ber and its products. *Indian Horticulture* 20: 5 &25.

Konopacka, D., Jesionkowska, K., Klewicki, R., and Bonazzi, C. (2009). The effect of different osmotic agents on the sensory dried fruit. *J. Hort. Sci. Biotech.* 84, 80-84. perception of osmo-treated

Krishnaveni, A., Manimegalai, G. and Saravanakumar, R., (2001). Storage stability of jack fruit (*Artocarpus heterophyllus*) RTS beverage. *J. Fd. Sci. Technol.*, 38(6): 601-602.

Kumar, S. P. and Devi, P. (2011). Optimization of some process variables in mass Transfer kinetics of osmotic dehydration of pineapple slices. *Inter. Food. Res. J.* 18: 221-238.

Kumar, S. P. and Sagar, V. R. (2009). Effect of osmosis on chemical parameters and sensory attributes of mango, guava slices and aonla segments. *Indian J. Hort.*, 66 (1): 53-57.

Kustagi K.U. (2002). Studies on processing of aonla (*Embllicaofficinalis* L.) fruits. M.Sc. (Hort.) Thesis, University of Agricultural Sciences, Dharwad, pp: 44.

Lakkond, B.R., (2002). Studies on processing of sapota (*Manilkaraachras* (Mil) Fosberg.). Fruits. M.Sc. (Hort) Thesis, University of Agricultural Sciences, Dharwad, pp: 44.

Lande R., Kale P.B., and Taley S.M. (1999). Studies on dehydration of ber fruits. 4th Agricultural Science Congress, Jaipur, India. February, 21-24: 214. atapi G. and Barret D. (2005). Influence of pre-drying treatments on quality and Safety of sun-dried tomatoes. Part 1: Use of steam blanching, boiling brine blanching and dips in salt or sodium metabisulphite. *J. Food Sci.*: 71(1):S24-S31.

Lewicki,P.P. and Lukaszuk, A. (2000). Effect of osmotic dewatering on rheological Properties of apple subjected to convective drying. *Journal of Food Engineering* 45(3): 119-126.

Lombard, G.E., Oliveira, J.C, Fito, P. and Andres, , (2008). Osmotic dehydration of pineapple as a pretreatment for further drying. *J. Food. Eng.*, 85 (2): 277-284.

- Madan S. and Dhawan S.S. (2005).** Development of value added product 'candy' from carrots. *Process Food Ind* 8(3):26-29
- Manivasagan, S. (2004).** Evaluation of karonda (*Carissa cardandas* Linn) cultivars for processing, CCS Haryana Agricultural University, Hisar, India, M.Sc. Thesis pp.:22.
- Mehta, A., Ranote, P.S. and Bawa, A.S. (2005).** "Processing of kandi lemon (*Galgal*) peel waste for candy making processing", CCS Haryana Agricultural University, Hisar, India, M.Sc. Thesis *Indian Food Packer*, 54: 67-74.
- Narayana, C.K., Shivasankar, S., Mustafa, M.M. and Sathiamoorthy, S. (2003).** Effect of different treatments on the quality of dehydrated banana (banana figs). *Indian Food Packer* 57(5): 66-68.
- Nath, A., Yadav, D.S., Pranabjyoti S. and Dey, B., (2005).** Standardization of ginger-kinnow squash and its storage. *J. Food Sci. Technol*, 42(6): 520-522.
- Nayak P., Bhatt D.K., Shukla D.K., Tandon D.K. (2011).** Evaluation of aonla (*Emblica Officinalis* G.) segments-in-syrup prepared from stored fruits. *Res. J. Agric. Sci*, 43(2).
- Nowakunda K., Andres. A, and Fito, P. (2004).** Osmotic dehydration of banana slices as a pre-treatment for drying process. *Proceedings of the 14th International drying symposium*. 20-25:2077-2083.
- Oyeyinka S.A., Ade-Omowave B.I.O., Ngoddy P.O., Karim O.R. (2011).** Selected quality attributes of jam produced from osmo-dehydrated cashew apple. *Journal of Food Technology*. 9(1):27-31.
- Panse V. G. and Sukhatme P. V. (1985).** Statistical methods for agricultural workers, ICAR, New Delhi.
- Panwar, J. K. (1981).** Postharvest physiology and storage behaviour of ber fruit (*Ziziphus mauritiana* Lamk.) in relation to temperature and various treatments. *Thesis Abstracts*. Haryana Agricultural University, Hisar. 7: 64-65.
- Parcek O.P. (2001).** Fruits for the future-2: Ber, International Centre for Underutilized Crops, University of Southampton, UK, 2001.
- Parcek, O. P. (2001).** Ber. International Centre for Underutilized Crops, Southampton, UK: 13.
- Patil, D.D.M., Katecha P.M. and Kadanm S.S. (1999).** Drying of ber preparation of shreds and powder. *Processed Food Industry Prajapati V.K., Nenma P.K. and Rathore S.S. (2010).* August Effect of pre-treatment and drying method on quality of value added dried aonla (*Emblica officinalis* Caeru) shreds. *Journal of food science and technology*, 48(1): 45-52.
- Priya, M.D. and Khatkar, B.S. (2013).** Effect of processing methods on keeping quality of aonla (*Emblica officinalis* Gaertn.) preserve *International Food Research Journal* 20(2): 617-622.
- Danoanna, S. (1980).** Hand book of Analysis and quality control for fruits and vegetable products. Tata McGraw Hill Publishing Company Limited, New Delhi.
- Rashmi. H.B., Doreyappa, G.I.N. and Mukunda, G.K., (2005).** Studies on osmo air dehydration of pineapple fruits. *J. Food Sei, Technol*, 42: 64-67.
- Rokhade, A. K., Chandan, K., Patil, P. B. and Patil, C. P. (2006).** Changes in physico-chemical characteristics of sweetened; dehydrated aonla slices during storage. Paper presented in National Seminar on production and processing of aonla (*Emblica officinalis* G.) pp: 38.
- Sharma S. K., Chaudhary S. P., Rao V. K., Yadav V. K. and Bisht T. S. (2013).** "Standardization of technology for preparation and storage of wild apricot fruit bar," *Journal of Food Science and Technology*, 50: 784-790. Babalola S. O. Ashaye O. A. Babalola A. O. and Aina J. O. (2002). "Effect of cold temperature storage on the quality attributes of pawpaw and guava leathers," *African Journal of Biotechnology*, 2: 57-60.
- Sagar, V.R. and Suresh. K.P. (2010).** "Recent advances in drying and dehydration of fruits and vegetables: A review", *J. Food Sci. Technol.*, 47 (1): 15-26.
- Saputra, D. (2001),** Osmotic dehydration of pineapple. *Drying Technol.*, 19 (2): +13 425.
- Sawate, A.R., Patil, V.P., Ghatge, P.U., Kshirsagar, R.B. and Tapre, A.R. (2005).** Studies on effect of syruing and drying methods on quality of papaya Candy. *J. Soils & Crops*, 15(1): 105-110.
- Siddiqui A.A., Bhuiyan M.H.R. and Easdani M. (2012).** Ginger (*Zingiber officinale*) preserve and candy development Bangladesh research publications journal 7(3): 283-290.

Shankersuwan Singh and AyushccGautam (2010). Effect of prctrcatment methods research an the qualitative and organoleptic attributes of pincapple candy during storage. Food science research journal 1(2): 76-85.

Sharma S.R., Alam S. and Gupta S. (2002).Storage study on dehydrated aonla powder. Annual Convention of Indian Society of Agricultural Engineers. 35: 155-1 56.

Shobha D., Sreedevi M.S. and Puttaramanaik (2018). Baby Corn Candy: Development and Assessment of Nutritional, Sensory and Storage Quality. It. J.Curr. Microbiol. App. Sci 7(3): 2261-2272. 17

Sharma, H. R., Pooja and Ranjana, V. (2006). Organoleptic and chemical evaluation of osmotically processed apricot wholes and halves.Natural Product Radiance, 5 (5): 350-356.

Singh, S.K., Narain, M. and Kumbhar, B.K., (2001). Effect of drying air temperatures and standard pre-treatments on the quality of fluidized bed dried button mushroom. Indian Food Packer, 55(5): 82-85.

Singh, B., Parmjit P. S. and Nanda, V. (2007). Rehydration Kinetics of Un Osomosed and Pre-Osomosed Carrot Cubes. World J Dairy Food Sei.2(1):10

Singh, O.M., Pathak, S. and Singh, R. (2012). Evaluation of sugar and stevia ratio for the preparation of aonla based low calorie quality beverages. Beverage & Food World, 39(4): 57-59.

Srilakshmi B. (2007). "Sensorv evaluation" Food science 4th Ed pp 286-297,246-256.

Sultana A.A., Khanom, M.D., MasbiarRahman and BurhanUddin M. (2015). Preparation of Pineapple (Ananas comosus) Candy Using Osmotic Dehydration Combined With Solar Drying. The Agriculturists a Scientific Journal of Krishi Foundation 13(1): 87-93.

Tandon D.K., Yadav R.C., Sood S., Kumar S. and Dikshit (2003). Effect of Blanching andlye

peeling on the quality of aonla candy. Indian food packer. 57(6): 147-152.

Unde, P.A., More, H.G., Jorwar, M.M., Sonar, D.J, and Sonawane, A.L. (2001). Effect of pre-treatments on drying characteristics and quality of banana slices, Journal of Maharashtra Agricultural University. 26 (3): 310-313.

Zeeshan, Salcem S.A., Ayub, S. nd Jan (2017). Physicochemical and Sensory Evaluation of Dhakki Dates Candy. Jornal of food processing and technology 8(3): 2157-7110.

Arghya mani, VSSV Prasanna, Shuvadeep Halder and J Praveena (2018) Efficacy of edible coatings blended with aloe vera in retaining post-harvest quality and improving storage attributes in Ber (*Ziziphus mauritiana* Lamk.) International Journal of Chemical Studies 6(6): 1727-1733.

Madhu Sangwan, V. M. Prasad (2023) Development and Evaluation of Natural Flavored Ber Candy. Journal of Experimental Agriculture International 45(8):72-78.

