

Original Research Article

Epidemiology of *Candida* species colonizing mucosae of HIV-infected patients in Cameroon

ABSTRACT

Aims: This study aimed to investigate *Candida* species colonizing HIV-infected patients in Bafoussam and Yaounde in Cameroon.

Study design. cross-sectional study.

Place and Duration of Study: Yaoundé Central Hospital (YCH) and Bafoussam Regional Hospital (BRH) between October 2018 and December 2020.

Methodology: We included 804 HIV-infected (681 women; 123 men; age range 21-81 years). Mucosal samples were collected, and mycological diagnosis was performed. Yeast isolates were identified using matrix-assisted laser desorption ionization-time-of-flight mass spectrometry (MALDI-TOF MS).

Results: Eight hundred and four patients were recruited, and 2754 samples were collected. The colonisation frequency was 17.35%, and 513 yeasts were isolated. Overall, *Candida albicans* 251 (48.92%) was the most frequently isolated. Non-*albicans Candida* (NAC) isolates have been classified into 16 species, including *Candida krusei* (14.23%), *Candida glabrata* (9.94%), *Candida parapsilosis* (8.18%), and *Candida tropicalis* (7.99%) as the major ones. There was a relationship (P -value= 0.00) between antiretroviral therapy and *Candida* species colonisation.

Conclusion: The results provide information on the epidemiology of *Candida* species in HIV-infected patients in Cameroon.

Keywords: *Candida* species, mucosae, HIV, epidemiology

1. INTRODUCTION

Candida species are the most common opportunistic pathogens in human immunodeficiency virus (HIV) infection [1]. Although *Candida albicans* remains the most common pathogen overall, non-*albicans Candida* species (NAC) has emerged as a potent pathogenic yeast in HIV-infected patients [2]. The most common types include *Candida glabrata*, *Candida tropicalis*, *Candida parapsilosis*, and *Candida krusei* [3,4]. The rarer ones are *Candida guilliermondii*, *Candida lusitanae*, *Candida kefyr*, *Candida famata*, *Candida inconspicua*, *Candida rugosa* and *Candida norvegensis* [3 -5]. The relative frequency of *Candida* species other than *Candida albicans* also varies with region [6]. *Candida albicans* was most common in Europe (52.5%) and least common in North America (42.7%), whereas *Candida glabrata* was most common in North America (24.3%) and least common in Latin America (7.1%). *Candida parapsilosis* and *Candida tropicalis* were more common than *Candida glabrata* in Latin America (24.3% and 17.0% vs 7.1%, respectively). *Candida tropicalis* was most common in the Asia-Pacific region (14.1%). *Candida krusei* was more common in North America (2.9%) and Europe (3.0%), and other miscellaneous species of *Candida* were common in Asia-Pacific region (7.3%) and North America (7.3%) [7]. Some epidemiological

studies have been carried out in Cameroon and have shown that the prevalence of *Candida* species in people living with HIV remains high [2, 8]. However, an important fact observed during the study of *Candida* species in Cameroon is that the identification of *Candida* species is generally limited on the basis of their phenotypic and biochemical characteristics. These classic techniques for identifying pathogenic yeasts are sometimes inadequate and have limitations such as their lack of sensitivity, specificity, and rapidity. In mycology, rapid detection and accurate, reliable identification are needed to confirm a diagnosis and implement appropriate antifungal treatment. Conventional identification techniques must therefore be accompanied by extremely precise molecular biology techniques to confirm a diagnosis. The present study was conducted to determine the *Candida* species distribution isolated from Cameroon HIV-infected patients by MALDITOF MS.

2. METHODOLOGY

2.1 Ethical considerations and enrollment of participants

The survey was carried out at Yaoundé Central Hospital (YCH) and Bafoussam Regional Hospital (BRH). The study involved HIV-infected patients presenting or not presenting clinical signs of any mucosal candidiasis and was approved by the Cameroonian National Ethical Committee. The survey took place from October 2018 to December 2020. Patients enrolled for this study were HIV-infected individuals of both genders, between 21 and 81 years old, who did not receive any antifungal treatment during the last 3 months preceding their enrollment.

2.2 Determination of the clinical status of patients and collection of study samples

Prior to sample collection, the patient's information was registered, including age and HIV status (CD4+ count, type of HIV, stage of the HIV infection, and antiretroviral therapy). Samples collected included vaginal discharge, oropharyngeal swab, stools, and urine.

2.3 Mycological diagnosis

Samples were submitted to direct macroscopic and microscopic analyses using routine laboratory protocols prior to culture on Sabouraud chloramphenicol medium for 24 to 48 hours at 37 °C. Primary identification was assessed by combination of culture on chromogenic medium (Chromagar Candida from Biomerieux, Marcy l'Etoile, France), germ tube test, evidence of chlamydospores production, and biochemical analysis (ID32 C kit from Biomerieux, Marcy l'Etoile, France). The reference *Candida krusei* ATCC 6258 strain was used throughout the mycological diagnosis as a quality control strain.

2.4 Identification by Mass Spectrometry

The second identification was performed by MALDI-TOF MS achieving as described by Cassagne et al. (2016) [9]. Briefly, analyses were performed on a MicroflexLT (Bruker Daltonics GmbH, Germany) equipped with a nitrogen laser (337nm). The mass range from 2000 to 20000 Da was recorded by using the linear mode. *C. krusei* ATCC 6258 was used for quality control. The spectra were compared to the MALDI Biotyper v3.0 software (Bruker Daltonics GmbH) containing the Bruker Daltonics database supplemented with our in-house yeast reference spectra for identification [9].

3. RESULTS

3.1 Patients characteristics

From October 2018 to December 2020, 804 HIV-infected patients from the two hospitals agreed to participate in the study by signing an informed consent form. At BRH, of the 322 participants recruited, 273 (84.78%) were women and 49 (15.21%) were men. At the YCH, of the 482 participants, 408 (84.64%) were women and 74 (13.35%) were men. The mean age of the general study population in the two hospitals was 42.39 ± 7.5 years, with extremes ranging from 21 to 81 years. 99.87% (n = 803/804) of patients in the study population were infected with HIV 1-M. 0.12% (n = 1/804) of HCY were coinfecting with types 1-M and 1-0.

3.2 Mycological diagnosis and yeast identification

3.2.1 Distribution of mucosal samples by sites sampled for the detection of *Candida* in the study population

A total of 2754 samples (732 stool samples, 617 urine samples, 601 vaginal swabs, and 804 oral swabs) were collected from the entire study population (Table 1). Cases with positive cultures for germs other than *Candida* species were excluded from the study. In the two hospitals, 31.42% (n = 230/732) of positive cultures were found in stools, 16.47% (n = 99/601) in vaginal swabs, 10.69% (n = 66/617) in urine, and 10.32% (83/804) in oral swabs. When considering the collected samples, the frequency of colonisation in the study population is 17.35%.

Table 1. Distribution of mucosal samples by sites sampled in the study population

		oral swabs	Vaginal swabs	stool	urine	Total (%)
BRH	Number of collected samples	322	231	279	205	1037
	Number of colonised samples	30	61	118	29	238 (22.95)
YCH	Number of collected samples	482	370	453	412	1717
	Number of colonised samples	53	38	112	37	240 (13.97)

*BRH : Bafoussam Regional Hospital; YCH : Yaounde Central Hospital

3.2.2 Distribution of *Candida* species isolated within the study population of the 02 hospitals

At BRH, the 5 most represented *Candida* species were in order *Candida albicans* 52.75% (n= 134/254), *Candida parapsilosis* 11.02% (n=28/254), *Candida krusei* 8.66% (n= 22/254), *Candida glabrata* 8.66% (n= 22/254) and *Candida dubliniensis* 7.87% (n=20/254). At the YCH, the 5 most common *Candida* species were *Candida albicans* 45.17% (n= 117/259), *Candida krusei* 19.69% (51/259), *Candida glabrata* 11.19% (29/259), *Candida tropicalis* 8.49% (22/259) and *Candida parapsilosis* 5.40% (14/259). In the 02 hospitals, *Candida*

albicans was the germ isolated in the majority of cases, whatever the sample. Table 2 shows the distribution of *Candida* species isolated and identified within the study population of the 02 hospitals.

Table 2. Distribution of *Candida* species isolated in the study population of the 02 hospitals

<i>Candida</i> species	BRH n(%)	YCH n(%)
<i>Candida albicans</i>	134(52.75)	117 (45.17)
<i>Candida parapsilosis</i>	28(11.02)	14 (5.40)
<i>Candida krusei</i>	22(8.66)	51 (19.69)
<i>Candida glabrata</i>	22(8.66)	29 (11.19)
<i>Candida dubliniensis</i>	20(7.87)	0
<i>Candida tropicalis</i>	19(7.48)	22 (8.49)
<i>Candida guilliermondi</i>	7 (2.75)	1(0.38)
<i>Candida catenulata</i>	1(0.39)	0
<i>Candida rugosa</i>	1(0.39)	11(4.24)
<i>Candida kefyr</i>	0	4(1.54)
<i>Candida intermedia</i>	0	1(0.38)
<i>Candida famata</i>	0	1(0.38)
<i>Candida sake</i>	0	3(1.15)
<i>Candida lusitanae</i>	0	2(0.77)
<i>Candida norvegensis</i>	0	2(0.77)
<i>Candida valida</i>	0	1(0.38)
Total	254 (49.51)	259 (50.48)

*BRH : Bafoussam Regional Hospital; YCH : Yaounde Central Hospital

Table 3 illustrates distribution of *Candida* species isolates according to gender, age, sample type, CD4 count, ARV status, and locality. Although *Candida* species were predominantly isolated from stools in the 02 hospital study population, there was no relationship (P -value= 0.39) between *Candida* colonisation and the different specimens collected. Within the study population of the 02 hospitals, the majority of *Candida* isolates were in the 31-40 age group. There was no relationship (P -value= 0.09) between the *Candida* spp. colonisation and age. *Candida* species were predominantly isolated from patients who were not on antiretroviral therapy compared to patients on antiretroviral therapy. There was a relationship (P -value= 0.00) between antiretroviral therapy and *Candida* spp. Colonization.

Table 3. Distribution of *Candida* species isolates according to gender, age, sample type, CD4 count, ARV status, and hospitals

Parameter	Category	<i>Candida</i>	Frequency	Chi-Square	P -value
-----------	----------	----------------	-----------	------------	------------

		species (N= 513)	(%)	(X ²)	
Gender	Male	98	19.10	0.057	0.80
	Female	415	80.89		
Age range (years)	21-30	84	16.37	6.428	0.09
	31-40	274	53.41		
	41-50	64	12.47		
	51-60	70	13.64		
	≥61	21	4.09		
Sample Type	Oral swab	70	13.64	4.066	0.39
	Vaginal swab	64	12.47		
	Stool	344	67.05		
	Urine	35	6.82		
CD4 count (CD4/mm ³)	<200	84	16.44	1.688	0.68
	200-349	204	39.73		
	350-499	120	23.29		
	≥500	105	20.54		
Antiretroviral therapy status	ARV (+)	14	2.74	14.261	0.00
	ARV (-)	499	97.26		
Locality of sample collection	BRH	254	31.51	7.524	0.11
	YCH	259	68.49		

*BRH: Bafoussam Regional Hospital; YCH: Yaounde Central Hospital; ARV: Antiretroviral therapy

4. DISCUSSION

HIV infection facilitates various opportunistic infections [10], including fungal infections [11], in particular candidiasis. Candidiasis is the most common opportunistic fungal infection in HIV-infected patients [11]. It is responsible for the high morbidity and mortality rates in these patients [12]. Mucosal candidiasis occurs in healthy individuals under the influence of a number of favourable factors, but is most common in patients with impaired cellular immunity, such as HIV-infected patients [13]. The study of mucosal candidiasis in HIV-infected patients is very often limited to the oropharyngeal mucosa, as this is an important indicator of the state of HIV infection [14-19]. However, the results obtained by this study reveal that all the human mucosae can be affected by *Candida* species. This result is similar to that obtained by Ngouana et al. [2] in Cameroon.

Nine and fourteen yeasts of the *Candida* genus were isolated at the BRH and YCH respectively. *Candida albicans* was the most species frequently isolated in the 02 hospitals. Whatever the type of sample analysed, *Candida albicans* was the species isolated in the majority. These results are in agreement with the results of studies carried out on mucosal candidiasis by Esebelahie et al. [11] in Nigeria, Maheshwari et al. [12] in India, and Ngouana et al. [2] in Cameroon.

Although *Candida albicans* remains the most frequently isolated agent in candidiasis studies, non-*albicans Candida* now account for a significant proportion of clinical isolates collected worldwide [2, 20]. More and more non-*albicans Candida* species are being isolated during studies of candidiasis in people living with HIV worldwide, and the prevalence of species varies from study to study, and from region to region. This increased emergence of non-*albicans Candida* could be attributed to the increase in the number of immunocompromised patients, particularly HIV-infected patients, the use of narrow-spectrum antifungal agents that act essentially only on *Candida albicans* [21], increased resistance of species other than

Candida albicans to antifungal agents, improvements in diagnostic techniques, such as the use of MALDI-TOF MS and routine diagnosis with molecular techniques [22, 23].

Candida species colonisation was predominantly observed in HIV-positive patients not receiving antiretroviral therapy compared with patients receiving antiretroviral therapy. In fact,

Candida species colonizing mucosae is common in HIV-positive patients who are naive to antiretroviral therapy [19]. Furthermore, there has been a significant decrease in the incidence of *Candida* spp. colonisation in HIV-infected patients since the discovery of antiretroviral therapy [24]. The intestinal tract was the most colonized site. This result is in agreement with that obtained by a previous study conducted in Cameroon on patients living with HIV [2].

Candida spp. colonisation in the study population was predominantly observed in female patients compared to male patients. This result is similar to that obtained in South Korea by Yoon et al. [25], Chile by Cruz et al. [26] and Cameroon by Ngouana [27]. This difference may be explained by differences in activities, personal hygiene and exposure to contamination between the female and male populations [28, 29].

4. CONCLUSION

A total of 513 *Candida* species were isolated. The frequency of colonisation in the study population was 17.35%. *Candida albicans* was the most common species isolated. *Candida krusei* was the most frequently isolated NAC species.

REFERENCES

1. Hosain PA, Salari S, Ghasemi NA. Oropharyngeal candidiasis in HIV/AIDS patients and non-HIV subjects in the Southeast of Iran. *Curr Med Mycol*. 2018 ; 4(4): 1-6. DOI: 10.18502/cmm.4.4.379
2. Ngouana TK., Toghueo RM, Kenfack KI, Lachaud L, Tadjou AK, Kouanfack C, Boyom FF, & Bertout S. "Epidemiology and antifungal susceptibility testing of non-*albicans* *Candida* species colonizing mucosae of HIV-infected patients in Yaounde (Cameroon)," *Journal de Mycologie Médicale*. 2019 ; 29 : 233-238.
3. Silva S, Negri M, Mariana HM, et al. *Candida glabrata*. *Candida parapsilosis* and *Candida tropicalis* : biology, epidemiology, pathogenicity and antifungal resistance. *FEMS Microbiol Rev*. 2012 ; 36 : 288–305.
4. Pfaller MA, Diekema DJ, Gibbs DL, et al. Results from the ARTEMIS DISK Global Antifungal Surveillance Study, 1997 to 2007 : a 10.5-year analysis of susceptibilities of *Candida* Species to fluconazole and voriconazole as determined by CLSI standardized disk diffusion. *J Clin Microbiol*. 2010 ;48 :1366–77. [http :// dx.doi.org/10.1128/JCM.02117-09](http://dx.doi.org/10.1128/JCM.02117-09).
5. Pfaller MA, Moet GJ, Messer SA, Jones RN, Castanheira M. Geographic variations in species distribution and echinocandin and azole antifungal resistance rates among *Candida* bloodstream infection isolates: report from the SENTRY anti- microbial surveillance program (2008 to 2009). *J Clin Microbiol*. 2011 ; 49 :396–9.
6. Xiao M, Xin F, Sharon CA, Chen HW, Zi-Yong S, Kang L, Shu-Lan C, Yan Y, Mei K, Zhi-Dong Hu, Yun-Zhuo C, Tie-Shi H, Yu-Xing N, Gui-Ling Z, Fanrong K, Ying-Chun X. Antifungal susceptibilities of *Candida glabrata* species complex, *Candida krusei*, *Candida parapsilosis* species complex and *Candida tropicalis* causing invasive candidiasis in

- China : 3 year national surveillance. *J Antimicrob Chemother* 2015 ; 70 : 802–810. doi: 10.1093/jac/dku460.
7. Pfaller MA, Diekema DJ, Turnidge JD, Castabheira M, Jones RN. Twenty years of the SENTRY antifungal surveillance program : results for *Candida* species from 1997–2016. *Open Forum Infect Dis.* 2019 ; 6 : S79–S94.
 8. Mandengue CE, Denning DW. The Burden of Serious Fungal Infections in Cameroon. *Journal Of Fungi* 2018 ; 4 : 44. doi :10.3390/jof4020044.
 9. Cassagne C, *et al.*, “Routine identification and mixed species detection in 6,192 clinical yeast isolates,” *Medical Mycology.* 54(3): 256–265, 2016, doi: 10.1093/mmy/myv095.
 10. Okolie MN, Eghafona NO, Omoregie R. Antihuman immunodeficiency virus agents. *J Med Lab Sci.* 2003 ; 12 : 1_14.
 11. Esebelahie NO, Enweani IB, Richard, OR. *Candida* colonisation in asymptomatic HIV patients attending a tertiary hospital in Benin City, Nigeria. *Libyan J Med.* 2013 ; 8 : 20322. <http://dx.doi.org/10.3402/ljm.v8i0.20322>.
 12. Maheshwari M, Kaur R, Sanjim C. *Candida* Species Prevalence Profile in HIV Seropositive Patients from a Major Tertiary Care Hospital in New Delhi, India. *Journal of Pathogens.* 2016 ; 2016 (1) : 1_8. <http://dx.doi.org/10.1155/2016/6204804>.
 13. Bloch M. Les candidoses chez l'immunodéprimé d'ordre secondaire. [Thèse de doctorat, Université de lorraine]. HAL. 2013 ; <https://hal.univ-lorraine.fr/hal-01733557/document>.
 14. Vazquez JA. Optimal management of oropharyngeal and oesophageal candidiasis in patients living with HIV infection HIV/AIDS. *Res Pal Care.* 2010 ; 2 : 89–101.
 15. Yang YL, Hung CC, Wang AH, Tseng FC, Leaw SN, Tseng YT, Su CL, Chen HT, Lauderdale TL, Lo HJ. Oropharyngeal colonization of HIV infected outpatients in Taiwan by Yeast Pathogens. *J Clin Microbiol.* 2010 ; 48(7) : 2609–12. <http://dx.doi.org/10.1128/JCM.00500-10>.
 16. Kalpanadevi V, Geethalakshmi S, Sumathi, GA. study on speciation and antifungal susceptibility pattern of *Candida* isolates from HIV patients with oropharyngeal candidiasis and correlation with CD4 count. *BMC Infect Dis.* 2012 ; 12 : 19.
 17. Lortholary O, Petrikos G, Akova M, Arendrup MC, Arikian-Akdagli S, Bassetti, M, BJ, Calandra T, Castagnola E, Cornely OA, Cuenca-Estrella M, Donnelly JP, Garbino J., Groll AH, Herbrecht R, Hope WW, Jensen HE, Kullberg, BJ, C Lass-Flörl, C, et al. ESCMID guideline for the diagnosis and management of *Candida* diseases 2012 : patients with HIV infection or AIDS. *Clin Microbiol Infect.* 2012 ; 18 : 68–77. <http://dx.doi.org/10.1111/1469-0691.12042>
 18. Merenstein D, Hu H, Wang C, Hamilton P, Blackmon M, Chen H, Calderone R, Dongmei L. Colonization by *Candida* species of the oral and vaginal Mucosa in HIV-Infected and non-infected women. *Aids Res Hum Retroviro.* 2013 ; 29 (1) : 30-4. <http://dx.doi.org/10.1089/aid.2012.0269>.
 19. Ngwa FA, Longdoh NA, Tebid P, Bobga TP, Nkfusai CN, Ngwa SB, Nsai FS, Nambile CS. The prevalence, risk factors and antifungal sensitivity pattern of oral candidiasis in HIV/AIDS patients in Kumba District Hospital, South West Region, Cameroon. *Pan African Medical Journal* 2020 ; 36 : 1-14. <http://www.panafrican-med-journal.com/content/article/36/23/full/>.
 20. Papon N, Courdavault V, Clastre M, Bennett RJ. Emerging and Emerged Pathogenic *Candida* Species : Beyond the *Candida albicans* Paradigm. *PLoS Pathog* 2013 ; 9(9) : 1003550.
 21. Sullivan DJ, Westerneng TJ, Haynes KA, Bennett DE, Coleman DC. *Candida dubliniensis* sp. nov. : phenotypic and molecular characterization of a novel species associated with oral candidosis in HIV-infected individuals. *Microbiol.* 1995 ; 141 : 1507–1521.
 22. Quindos G, Marcos-Arias C, San-Millan R, Mateo E, Eraso E. The continuous changes in the aetiology and epidemiology of invasive candidiasis: from familiar *Candida albicans* to multiresistant *Candida auris*. *Int Microbiol* 2018 ; 21 :107-119.

23. Sadeghi G, Ebrahimi-Rad M, Mousavi SF, Shams-Ghahfarokhi M, Razzaghi-Abyaneh M. Emergence of non-*Candida albicans* species : epidemiology, phylogeny and fluconazole susceptibility profile. *J Mycol Med.* 2018, 28 : 51–58.
24. Taverne-Ghadwal L, Kuhns M, Buhl T, Schulze MH, Mbaitolum WJ, Kersch L, Weig M, Bader O, Groß U. Epidemiology and Prevalence of Oral Candidiasis in HIV Patients From Chad in the Post-HAART Era. *Frontiers in microbiology.* 2022 ; 13 : 408. <https://doi.org/10.3389/fmicb.2022.844069>.
25. Yoon HJ, Choi HY, Kim YK, Song YJ, Ki M. Prevalence of fungal infections using National Health Insurance Data From 2009-2013. *South Korea Epidemiol Health.* 2014 ; 36 : E2014017. doi: 10.4178/epih/e2014017
26. Cruz CR, Ponce EE, Calderón RL, Delgado VN, Vieille OP, Piontelli LE. Superficial mycoses in the city of Valparaiso, Chile : period 2007-2009. *Rev. Chil. Infectol,* 2011 ; 28 : 404–409. doi: 10.4067/S0716-10182011000600002.
27. Ngouana KT. Diversité génétique d'isolats de *Cryptococcus* et *Candida* issus des patients VIH positifs à Yaoundé et étude de leur sensibilité aux antifongiques et aux extraits de plantes. [Thèse de doctorat, Université de Montpellier I]. HAL. 2014 ; <https://tel.archives-ouvertes.fr/tel-01132146>.
28. Albuquerque MF, Oliveira-Júnior JB, dos Santos MA, Silva DM, “Mycoses in northeastern Brazil: epidemiology and prevalence of fungal species in 8 years of retrospective analysis in Alagoas.” *Brazilian Journal of Microbiology* 2019; 50(4): 969–978, doi: 10.1007/s42770-019-00096-0.
29. Bilal H, Hou B, Shafiq M, Chen X, Shahid MA, Zeng Y. “Antifungal susceptibility pattern of *Candida* isolated from cutaneous candidiasis patients in eastern Guangdong region: A retrospective study of the past 10 years.” *Frontiers in Microbiology.* 2022; 13: 1-9. doi: 10.3389/fmicb.2022.981181.