

# Original Research Article Knowledge, Attitude, and Perception of Nigerians Towards Herbal Contraception: A Questionnaire-Based Study

---

## ABSTRACT

**Aims:** With a population of about 222 million, which is expected to rise to 377 million making Nigeria the 3<sup>rd</sup> largest country in the world by 2050, a well-planned family will improve the quality of life of mother and child.

**Methodology:** A non-probability online survey was conducted (October 2021 and November 2021), recruiting 1113 participants across Nigeria. Socio-demographic data and information on contraceptives were obtained, and Pearson's chi-squared test of independence at a 95% confidence interval (CI) was computed using R 4.2.1.

**Results:** Significant associations ( $P < .001$ ) were found between age, marital status, education, and profession of respondents. About 44.65% were 15-29 years, while 41.2% were 30-34 years. No significant association was observed between the age group, gender, and the need for population control. Females (50.04%), 15-29 years old (66.48%) are more likely to approve of the need for population control. Likewise, females (22.01%), with tertiary education (18.78%) are well-informed about contraceptive use ( $P < .001$ ). Only 19.38% ( $n = 184$ ;  $P < .001$ ) of participants used herbal contraceptives, in the form of liquid mixtures (55.98%), teas (17.93%), pills (12.50%), and powders (10.87%), with 26.63% > 7.61% > 6.52% > 5.98% = 5.98% = 5.98% > 4.89% > 3.80% who took Moringa, Alligator Pepper, Neem Plant, Castor Beans, African Star apple Momordica, Rosary Pea, and Horse-eye Bean respectively as contraceptives. Aged parents (32.07%), friends (31.52%), doctors (15.76%), herbalists (5.98%), pharmacists (5.43%), and nurses (3.80%) played a significant ( $P < .001$ ) role in recommending these herbs. Furthermore, 76.37% recommended the development of indigenous herbal contraceptives in Nigeria, while 23.63% did not.

**Conclusion:** Although participants have expressed a preference for herbal contraceptives, knowledge of them is lacking, which calls for more awareness campaigns and policy formulations for their promotion, development, and adoption in Nigeria.

*Keywords:* Nigeria; Family planning; conventional contraceptives; herbal contraceptives; population control

## 1. INTRODUCTION

According to the United Nations, Nigeria has the largest population in Africa, with 222,266,236 people (as of 3/17/2023), and is expected to grow to more than 377 million people by the year 2050 [1,2]. Managing the population is an essential part of the Sustainable Development Goals (SDGs). The negative impacts of rapid growth on a country are manifested in employment, species extinction, deforestation, desertification, climate change, destruction of ecosystems, and pressure on housing [3]. Hence, accessible and cost-effective family planning (FP) programs are needed to boost Nigeria's economic development [4].

Family planning programs play an important role in reducing the number of children born and thus improving their quality of life [5]. Couples who choose FP freely space, and time, and limit the number of children they want [6]. It has been reported [7] that about 50 % of pregnancies are unintended, with 50 % of undesirable outcomes such as risky abortions, deaths, and unwanted children who bear the burden of poverty and neglect. Nigeria utilizes only 16 % of contraceptives when compared to what is obtainable globally [7]. Even though contraceptives provide many benefits, some contraceptive methods are associated with undesirable side effects, including heart attacks, hypertension, migraines, strokes, blood clots, headaches, depression, obesity, increased risk of breast cancer, cervical cancers, and suicide attempts [8]. It is against this backdrop that, many countries are turning to the use of herbs to control fertility [9], resulting in natural product researchers re-focusing their interest on the use of herbs with potential contraceptive effects [10]. In Nigeria, herbs have been used as contraceptives due to their availability, cost-effectiveness, minimal side effects, and popularity [11]. According to the World Health Organization (WHO), about 80 % of the population relies on herbal medicines (HMs) for their primary healthcare needs [12]. About 133 plants have been identified to possess contraceptive ability [13], these possess sterilizing, contraceptive, or abortifacient properties by either inhibiting the implantation of the ova, influencing sperm count, motility/viability, or causing early expulsion of the fetus, and termed as antifertility, anti-ovulatory, anti-implantation, or abortifacient herbs [14,15]. A recent survey conducted by members of our group [11] confirmed that the preference for HMs has continued to increase, however, our study did not look at the specific disease areas, neither are there substantial studies in recorded literature in this area. It is therefore of high critical priority to assess the knowledge, attitude, and perception of Nigerians about herbal contraceptives (HCs) in order to not only inform evidence-based regulatory policies, but also to direct efforts towards Research and Development (R&D) of therapeutics for population control towards national socio-economic development.

## **2. METHODOLOGY**

### **2.1 Study Setting**

Nigeria is situated in the West Africa region, and lies between longitudes 3° and 14° and latitudes 4° and 14°, between the Sahel to the North and the Gulf of Guinea to the South in the Atlantic Ocean and has a landmass of 923,769 km<sup>2</sup> [16].

### **2.2 Sampling Method**

To recruit participants in Nigeria, an online survey (October 2021 and November 2021) was conducted by sharing the link to the questionnaire on social media platforms, using the snowball sampling technique. Participation was completely consensual, anonymous, and voluntary. Socio-demographic data and information on contraceptives were obtained. A descriptive statistical analysis was conducted on the data obtained, and Pearson's chi-squared test of independence at 95 % confidence interval (CI) was computed using R 4.2.1 statistical package.

## **3. RESULTS**

### **3.1 State of Origin of Respondents**

A total of 1113 participants were recruited from the six (6) geographical zones of Nigeria (Fig. 1.). Results show that 27.67 % (n = 308) of the respondents were from North-Central

Nigeria and 8.63 % (n = 96) from the Northeast. About 6.38 % of respondents live in Kogi state; 6.20 % in Benue, and 5.57 % live in Delta state.




Fig. 1. Respondents according to geopolitical zones in Nigeria

### 3.2 Socio-demographic characteristics of respondents

The results of the demographic information (Age group, gender, marital status, educational qualification, and profession) of the respondents are shown in Table 1. It was observed that there is a significant difference ( $P < .001$ ) in the age, marital status, educational qualification, and profession of the respondents.

### 3.3 Need for Population Control

Details of the demography of respondents and the need for population control are shown in Table 2. No significant association was observed between the age group and gender of the respondents to their perception of the need for population control. However, there was a significant association ( $P < .001$ ) between the need for population control and marital status, educational qualification, and profession. These parameters showed that 41.78 % of participants who are married and have obtained their tertiary education (43.86 %) indicated the need for population control. However, only 26.06 % of the population are healthcare workers.

Table 1: Demographic Information of Respondents

Variables	Demographic Information (N = 1113)	
	Responses n (%)	Chi-Squared Test (95 % CI)
<b>Age Group</b>		
15-29	497 (44.65)	$\chi^2 = 1027$ df = 4 p-value < .001
30-44	459 (41.24)	
45-59	135 (12.13)	
60-74	21 (1.89)	
75-89	1 (0.09)	

<b>Gender</b>		
Female	668 (60.02)	$\chi^2 = 616.37$
Male	442 (39.71)	df = 2
Others	3 (0.27)	p-value < .001
<b>Marital Status</b>		
Single	475 (42.68)	$\chi^2 = 1344.30$
Married	565 (50.76)	df = 4
Divorced	37 (3.32)	p-value < .001
Separated	26 (2.34)	
Others	10 (0.90)	
<b>Education</b>		
Primary	25 (2.25)	$\chi^2 = 731.45$
Secondary	119 (10.69)	df = 4
Tertiary	534 (47.98)	p-value < .001
Postgraduate	304 (27.58)	
Others	128 (11.50)	
<b>Profession</b>		
Healthcare	314 (28.21)	$\chi^2 = 221.34$
Others	799 (71.79)	df = 1
		p-value < .001

N = Number of observations; n = number of respondents; % = Percentage response; df = Degree of freedom; CI = Confidence Interval

**Table 2: Perception of Respondents on the Need for Population Control According to Demographic Information**

Demographic Information	Need for population control (N = 1113)		
	No n (%)	Yes n (%)	Chi-Squared Test (95 % CI)
<b>Age Group</b>			
15-29	74 (6.65)	423 (66.48)	$\chi^2 = 8.88$
30-44	83 (74.57)	376 (38.01)	df = 4
45-59	22 (1.98)	113 (10.15)	P-value = .06
60-74	8 (0.72)	13 (1.17)	
75-89	0 (0.00)	1 (0.08)	
<b>Gender</b>			
Female	111 (9.97)	557 (50.04)	$\chi^2 = 5.35$
Male	74 (6.65)	368 (33.06)	df = 2
Others	2 (0.18)	1 (0.09)	P-value = .07
<b>Marital Status</b>			
Single	60 (5.39)	415 (37.29)	$\chi^2 = 30.64$
Married	100 (8.98)	465 (41.78)	df = 4
Divorced	11 (0.99)	26 (2.34)	P-value < .001
Separated	11 (0.99)	15 (1.35)	
Others	5 (0.45)	5 (0.45)	
<b>Education</b>			
Primary	13 (1.17)	12 (1.08)	$\chi^2 = 68.82$
Secondary	36 (3.23)	38 (7.46)	df = 4
Tertiary	57 (5.12)	477 (42.86)	P-value < .001

Postgraduate	43 (3.86)	264 (23.72)	
Others	38 (3.41)	90 (8.09)	
<b>Profession</b>			
Healthcare	24 (2.16)	290 (26.06)	$\chi^2 = 25.34$
Others	163 (14.65)	636 (57.14)	df = 1
			P-value < .001

N = Number of observations; n = number of respondents; % = Percentage response; df = Degree of freedom; CI = Confidence Interval

### 3.4 Awareness of Respondents on the Use of Contraceptives

Our study found a significant association between the age group ( $P = .003$ ), gender, marital status, educational qualification, and profession ( $P < .001$ ) of the participants, and how informed they are on the use of contraceptives (Table 3). The data presented shows that 22.01 % of females and 18.78 % of those who have obtained their tertiary education are well-informed about the use of contraceptives.

**Table 3: Awareness of Respondents on the Use of Contraceptive**

Demographic Information	Information on contraceptive use (N = 1113)					Chi-Squared Test (95 % CI)
	Very well informed n (%)	Well informed n (%)	Little informed n (%)	Not informed n (%)	Others n (%)	
<b>Age</b>						
15-29	72 (6.47)	154 (13.83)	187 (16.80)	80 (7.18)	4 (0.36)	$\chi^2 = 35.48$
30-44	82 (7.37)	152 (13.66)	152 (13.65)	65 (5.84)	8 (0.72)	df = 16
45-59	22 (1.98)	58 (5.21)	39 (3.50)	14 (1.26)	2 (1.80)	P-value = .003
60-74	0 (0.00)	4 (0.36)	11 (0.99)	4 (0.36)	2 (0.18)	
75-89	0 (0.00)	0 (0.00)	0 (0.00)	1 (0.09)	0 (0.00)	
<b>Gender</b>						
Female	104 (9.34)	245 (22.01)	228 (20.46)	82 (7.37)	9 (0.81)	$\chi^2 = 104.86$
Male	72 (6.47)	123 (11.05)	160 (14.38)	82 (7.37)	5 (0.45)	df = 8
Others	0 (0.00)	0 (0.00)	1 (0.09)	0 (0.00)	0 (0.00)	P-value < .001
<b>Marital Status</b>						
Single	71 (6.38)	137 (12.31)	192 (17.25)	71 (6.38)	4 (0.37)	$\chi^2 = 82.88$
Married	98 (8.81)	215 (19.32)	175 (15.73)	71 (6.38)	6 (0.54)	df = 16
Divorced	6 (0.54)	10 (0.89)	11 (0.99)	7 (0.63)	3 (0.27)	P-value < .001
Separated	1 (0.09)	3 (0.27)	10 (0.89)	11 (0.99)	1 (0.09)	
Others	0 (0.00)	3 (0.27)	1 (0.09)	4 (0.36)	2 (0.18)	
<b>Education</b>						
Primary	0 (0.00)	3 (0.27)	8 (0.72)	14 (1.23)	0 (0.00)	$\chi^2 = 147.46$
Secondary	10 (0.90)	20 (1.80)	61 (5.48)	26 (2.34)	2 (0.18)	df = 16
Tertiary	79 (7.10)	209 (18.78)	198 (17.79)	38 (3.41)	10 (0.90)	P-value < .001
Postgraduate	70 (6.29)	110 (9.88)	82 (7.37)	44 (3.95)	1 (0.09)	
Others	17 (1.53)	26 (2.34)	40 (3.59)	42 (3.77)	3 (0.27)	
<b>Profession</b>						
Healthcare	79 (6.83)	146 (1.08)	76 (0.09)	12 (7.10)	1 (0.09)	$\chi^2 = 101.65$
Others	97 (28.12)	222 (13.66)	313 (1.35)	152 (8.72)	15 (1.35)	df = 4
						P-value < .001

N = Number of observations; n = number of respondents; % = Percentage response; df = Degree of freedom; CI = Confidence Interval

### 3.5 Knowledge of Respondents on Types of Contraceptives

Table 4 shows a significant association ( $P < .001$ ) between the demographic information of the participants and the knowledge of the types of contraceptives. Most of the respondents are familiar with conventional contraceptives, compared to HCs.

**Table 4: Knowledge of Respondents on Types of Contraceptives**


Demographic Information	Knowledge of Types of Contraceptives (N = 949)				Chi-Squared Test (95 % CI)
	Conventional Contraceptive n (%)	Herbal Contraceptive n (%)	Both n (%)	None n (%)	
<b>Age</b>					
15-29	293 (30.87)	67 (7.06)	50 (5.27)	7 (0.74)	$x^2 = \text{NaN}$
30-44	279 (29.40)	51 (5.37)	52 (5.48)	12 (1.26)	df = 12
45-59	87 (9.17)	14 (1.48)	18 (1.90)	2 (0.21)	$P\text{-value} = \text{NA}$
60-74	8 (0.84)	3 (0.32)	4 (0.42)	2 (0.21)	
75-89	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)	
<b>Gender</b>					
Female	447 (47.10)	62 (6.53)	66 (6.95)	11 (1.16)	$x^2 = 35.88$
Male	220 (23.18)	71 (7.59)	56 (5.90)	12 (1.26)	df = 6
Others	0 (0.00)	1 (0.11)	2 (0.21)	0 (0.00)	$P\text{-value} < .001$
<b>Marital Status</b>					
Single	383 (29.80)	64 (6.74)	51 (5.37)	6 (0.63)	$x^2 = 66.1$
Married	366 (38.57)	52 (5.48)	62 (6.53)	14 (1.48)	df = 12
Divorced	11 (1.16)	11 (1.16)	5 (0.53)	3 (0.32)	$P\text{-value} < .001$
Separated	5 (0.53)	8 (0.84)	2 (5.37)	0 (0.00)	
Others	2 (0.21)	0 (0.00)	4 (0.42)	0 (0.00)	
<b>Education</b>					
Primary	8 (0.84)	3 (0.32)	0 (0.00)	0 (0.00)	$x^2 = 119.47$
Secondary	63 (6.64)	27 (2.85)	2 (0.21)	1 (0.11)	df = 12
Tertiary	376 (39.62)	54 (5.69)	57 (6.00)	9 (0.95)	$P\text{-value} < .001$
Postgraduate	193 (20.34)	18 (1.90)	46 (4.85)	6 (0.63)	
Others	27 (2.84)	33 (3.48)	19 (2.00)	7 (0.74)	
<b>Profession</b>					
Healthcare	238 (25.08)	12 (1.26)	49 (5.16)	3 (0.32)	$x^2 = 44.43$
Others	429 (45.21)	123 (12.96)	75 (7.90)	20 (2.11)	df = 9
					$P\text{-value} < .001$

N = Number of observations; NaN = Not a Number; NA = Not applicable; n = number of respondents; (%) = Percentage response; df = Degree of freedom; CI = Confidence Interval

### 3.6 Respondents who have Used Herbal Contraceptives

#### 3.6.1 The proportion of respondents who used HCs and the dosage form used

Fig. 2. presents data on respondents who have used one form of HCs or the other. Only 19.38 % ( $n = 184$ ;  $P < .001$ ) of the respondents have used HCs, in the form of herbal liquid mixtures: 55.98 % ( $n = 103$ ); herbal contraceptive teas: 17.93 % ( $n = 33$ ); herbal contraceptive pills: 12.50 % ( $n = 23$ ); and herbal contraceptive powders: 10.87 % ( $n = 20$ ).



**Fig. 2. (a) Proportion of respondents who used herbal contraceptives; (b) Dosage form of herbal contraceptives used by respondents**

### 3.6.2 Types of herbal contraceptives used

Interestingly, there was a significant association ( $P < .005$ ) between male participation and females who took HCs, results show that more males took *Chrysophyllum albidum*, *Aframomum melegueta*, *Moringa oleifera*, and *Azadirachta indica* than females (Table 5).


**Table 5: Proportion of males and females who took herbal contraceptives**

Herbs used as contraceptives	Scientific Name	Number of Observations (N = 184)			Chi-Squared Test (95 % CI)
		Females n (%)	Males n (%)	Others n (%)	
African Star apple	<i>Chrysophyllum albidum</i>	3 (1.63)	8 (4.35)	0 (0.00)	$\chi^2 = 37.14$ df = 18 P-value = .01
Alligator Pepper	<i>Aframomum melegueta</i>	6 (3.26)	8 (4.35)	0 (0.00)	
Castor Beans	<i>Ricinus communis</i>	8 (4.35)	3 (1.63)	0 (0.00)	
Horse-eye Bean	<i>Mucuna urens</i>	7 (3.80)	0 (0.00)	0 (0.00)	
Momordica	<i>Momordica charantia</i>	6 (3.26)	4 (2.17)	1 (0.54)	
Moringa	<i>Moringa oleifera</i>	23 (12.50)	26 (14.13)	0 (0.00)	
Neem Plant	<i>Azadirachta indica</i>	3 (1.63)	9 (4.89)	0 (0.00)	
Rosary Pea	<i>Abrus precatorius</i>	5 (2.71)	4 (2.17)	0 (0.00)	
Other	-	16 (8.70)	17 (9.24)	0 (0.00)	
None	-	20 (10.87)	7 (3.80)	0 (0.00)	

N = Number of observations; n = number of respondents; % = Percentage response; df = Degree of freedom; CI = Confidence Interval

### 3.6.3 Who recommended the use of HCs

Furthermore, Fig. 3 shows that aged parents, friends, doctors, herbalists, pharmacists, and nurses played a significant ( $P < .001$ ) role in recommending these herbs to the respondents.



**Fig. 3. Proportion of those who were prescribed herbal contraceptives**

### 3.7 Recommendation for the development and use of an indigenous herbal contraceptive

#### 3.7.1 Recommendation for the development of HCs

Table 6 shows that 76.37 % (n = 850) of the respondents recommended the development, promotion, adoption, and use of indigenous HCs. No significant association between age group, gender, marital status, and profession was observed. Only educational qualification has a significant association ( $P < .001$ ) with the recommendation for the development and use of HCs.

**Table 6: Recommendation for the Development and Use of HCs**


Demographic information	Recommendation for use of herbal contraceptives (N = 1113)		Chi-Squared Test (95 % CI)
	No (%)	Yes (%)	
<b>Age</b>			
15-29	119 (10.91)	378 (33.96)	$\chi^2 = 7.28$
30-44	111 (9.73)	348 (31.26)	df = 4
45-59	24 (21.56)	111 (9.97)	P-value = .12
60-74	9 (0.81)	12 (1.08)	
75-89	0 (0.00)	1 (0.09)	
<b>Gender</b>			
Female	152 (13.66)	516 (46.36)	$\chi^2 = 3.623$
Male	109 (9.79)	333 (29.92)	df = 2
Others	2 (0.18)	1 (0.09)	P-value = .16
<b>Marital Status</b>			
Single	125 (11.23)	350 (31.45)	$\chi^2 = 7.36$
Married	116 (10.42)	449 (40.34)	df = 4
Divorced	10 (0.90)	27 (2.43)	P-value = .12
Separated	8 (0.72)	18 (1.62)	
Others	4 (0.36)	6 (0.54)	

<b>Education</b>			
Primary	13 (1.17)	12 (1.08)	$\chi^2 = 19.74$
Secondary	39 (3.50)	80 (7.19)	df = 4
Tertiary	109 (9.79)	425 (38.19)	P-value < .001
Postgraduate	72 (6.47)	235 (21.11)	
Others	30 (2.70)	98 (8.80)	
<b>Profession</b>			
Healthcare	66 (5.93)	248 (22.28)	$\chi^2 = 1.4566$
Others	197 (17.70)	602 (54.09)	df = 1
			P-value = .23

N = Number of observations; n = number of respondents; % = Percentage response; df = Degree of freedom; CI = Confidence Interval

### 3.7.2 Recommended dosage form and frequency of HCs

Figure 4 shows that 48.12 % (n = 409) of respondents preferred the development of HC teas; 18.35 % (n = 156%) preferred chocolate-based HCs; 22.47 % (n = 22.47%) preferred HC pills; 9.65 % (n = 82) preferred herbal vaginal tablets; while 1.41 % (n = 12) preferred herbal vaginal suppositories (Figure 4a). Regardless of the dosage form specified, 29.88 % (n = 254), 25.53 % (n = 217), 17.65 % (n = 150), 14.00 % (n = 119), and 9.29 % (n = 79) of the respondents indicated that the HCs should be taken monthly, every 3 months, 6 months, weekly, and yearly basis respectively (Fig. 4b).



**Fig. 4. Development of herbal contraceptives: (a) Dosage form for the development of herbal contraceptives; (b) Duration of herbal contraceptive use**

### 3.7.3 Who should HCs be developed for?

Table 7 shows respondents' perception of who HCs should be developed for. Although this is not significantly different within gender ( $P = .05$ ), and age group ( $P = .11$ ), more participants think HCs be developed for both males and females (Table 7).

**Table 7: Perception of Respondents on Whom Herbal Contraceptives Should be Developed**

Demographic Information	Who should herbal contraceptives be developed for (N = 850)			Chi-Squared Test (95 % CI)
	Both	Men	Women	
<b>Age</b>				
15-29	186 (21.88)	26 (3.06)	166 (19.53)	$x^2 = 13.22$ df = 8 P-value = .11
30-44	185 (21.76)	35 (4.12)	128 (15.06)	
45-59	61 (7.18)	3 (0.35)	47 (5.53)	
60-74	5 (0.59)	0 (0.00)	7 (0.82)	
75-89	0 (0.00)	0 (0.00)	1 (0.12)	
<b>Gender</b>				
Female	260 (30.59)	30 (35.29)	226 (26.59)	$x^2 = 9.4517$ df = 4 P-value = .05
Male	177 (20.82)	34 (4.00)	122 (14.35)	
Others	0 (0.00)	0 (0.00)	1 (0.12)	
<b>Marital Status</b>				
Single	197 (23.18)	29 (3.41)	124 (14.59)	$x^2 = 21.64$ df = 8 P-value = .01
Married	227 (26.71)	29 (3.41)	193 (22.71)	
Divorced	8 (0.94)	2 (0.24)	17 (2.00)	
Separated	3 (0.35)	3 (0.35)	12 (1.41)	
Others	2 (0.24)	1 (0.12)	3 (0.35)	
<b>Education</b>				
Primary	4 (0.47)	0 (0.00)	8 (0.94)	$x^2 = 80.655$ df = 8 P-value < .001
Secondary	35 (4.12)	5 (0.59)	40 (4.71)	
Tertiary	237 (27.88)	21 (2.47)	169 (19.47)	
Postgraduate	146 (17.18)	20 (2.35)	69 (8.12)	
Others	15 (1.76)	18 (2.12)	65 (7.65)	
<b>Profession</b>				
Healthcare	170 (20.00)	14 (1.64)	64 (7.53)	$x^2 = 41.493$ df = 2 P-value < .001
Others	267 (31.41)	50 (5.88)	285 (33.53)	

N = Number of observations; n = number of respondents; % = Percentage response; df = Degree of freedom; CI = Confidence Interval

### 3.7.4 Preference for the type of contraceptives

Table 8 shows that there was a significant association between age ( $P = .01$ ), gender ( $P = .01$ ), marital status ( $P = .003$ ), and educational qualification ( $P < .001$ ) on the preference for HCs. More respondents who preferred the HCs are within the age groups of 15-29 and 30-44 years as seen in the table.

**Table 8: Responses on the Preference for Type of Contraceptive**


Demographic Information	Preference for Contraceptive (N = 1113)			Chi-Squared
	Conventional	Herbal	None	

	Contraceptive	Contraceptive		Test (95 % CI)
<b>Age</b>				
15-29	228 (20.49)	231 (20.75)	38 (3.41)	$\chi^2 = 19.27$
30-44	213 (19.14)	213 (19.14)	33 (2.96)	df = 8
45-59	52 (4.67)	77 (0.99)	6 (0.18)	<i>P</i> -value = .01
60-74	8 (0.72)	11 (0.99)	2 (0.18)	
75-89	0 (0.00)	0 (0.00)	1 (0.09)	
<b>Gender</b>				
Female	279 (25.070)	332 (29.83)	57 (5.12)	$\chi^2 = 13.08$
Male	222 (19.94)	197 (17.70)	23 (2.07)	df = 4
Others	0 (0.00)	3 (0.00)	0 (0.27)	<i>P</i> -value = .01
<b>Marital Status</b>				
Single	249 (22.37)	192 (17.25)	34 (3.05)	$\chi^2 = 23.76$
Married	229 (20.58)	294 (26.42)	42 (3.77)	df = 8
Divorced	11 (0.99)	23 (2.07)	3 (0.27)	<i>P</i> -value = .003
Separated	9 (0.81)	16 (1.44)	1 (0.09)	
Others	3 (0.27)	16 (0.63)	0 (0.00)	
<b>Education</b>				
Primary	13 (1.17)	12 (1.08)	0 (0.00)	$\chi^2 = 30.66$
Secondary	45 (4.04)	69 (6.20)	5 (0.45)	df = 8
Tertiary	241 (21.65)	256 (23.00)	37 (3.32)	<i>P</i> -value < .001
Postgraduate	155 (13.93)	118 (10.60)	34 (3.05)	
Others	47 (4.22)	77 (6.92)	4 (0.36)	
<b>Profession</b>				
Healthcare	171 (15.36)	111 (9.97)	32 (2.87)	$\chi^2 = 28.34$
Others	330 (29.65)	421 (37.82)	48 (4.31)	df = 2
				<i>P</i> -value < .001

N = Number of observations; n = number of respondents; % = Percentage response; df = Degree of freedom; CI = Confidence Interval

### 3.7.5 Reasons for preference of HCs

Figure 5 shows respondents who prefer the HCs with reasons such as they are indigenous (9.59 %), they have few side effects (38.16%), they are organic (31.77 %), and are affordable (18.23 %) as shown in figure 5a. On the other hand, they do not prefer conventional contraceptives because they think they have so many side effects (70.86 %), they are expensive (11.84 %), have a high failure rate (8.08 %), and are not readily available (5.64 %) as shown in Fig. 5b.



**Fig. 5. Respondents who preferred HCs: (a) Reasons for preference for herbal contraceptives; (b) Reasons for non-preference for conventional contraceptives**

#### 4. DISCUSSION

Respondents' state of origin reveals a good representation of the Nigerian populace. It is possible, however, that the low responses recorded in other geographical zones are a result of bias in the sampling method employed, attitudes/perceptions to social media, lack of interest/knowledge of the topic, or reluctance to participate in the survey. While women in Northeastern Nigeria are unlikely to use contraceptives due to religious beliefs, spousal consent, knowledge of contraceptives, and age [17], obstacles to contraception in the Southeast include a lack of knowledge about contraceptives, side effects, barriers to sex education, and lack of access to contraceptives [18]. In contrast, contraceptives are widely known and accepted in North and Northcentral Nigeria, but actual use has not been demonstrated [19,20]. Furthermore, there was a low rate of contraceptive use in Southern Nigeria [21], and Adebisi, and Bello [22] mentioned male participation in South West Nigeria.

A positive correlation existed between age group and sexual activity. A total of 85.89 % of our participants were within the reproductive age group (15-49 years), with 15-29 years being the most sexually active. This age group is particularly knowledgeable due to their disposition towards social media, and have been reported to use one or more forms of contraceptives globally [23]. Moreover, respondents with postgraduate education (75.56 %) showed a high degree of association with information on contraceptives, which demonstrates their ability to make decisions on FP. This study showed a higher percentage compared to that obtained by Odusina et al. [24] with 60 % of participants.

Healthcare providers play a significant role in the acceptance and uptake of contraceptives. Although, highly intellectual individuals participated in this study, only a handful (28.21 %) of them are healthcare professionals. Several reasons such as hesitancy, lack of interest in the topic, and the desire to stay focused on what they believe in could be postulated. There is therefore a need for a mind-shift in these individuals by embracing the current trend and

direction of contraceptives, understanding their roles/involvement in healthcare delivery and the urgent need for more contraceptive options in Nigeria, in addition to formulating and implementing policies for adequate inclusion of FP in healthcare delivery system in Nigeria. Studies conducted by Saka et al. [25] also showed this trend.

The need for population control was supported by 83.11 % of our respondents. Nigeria is currently (3/17/2023) the 6<sup>th</sup> largest population in the world, with a total fertility rate of 5.3 live births per woman [26]. This pre-empted the establishment of the National Policy on Population (NPP) in 1988, which was later revised in 2004 to curb the rise in population through FP [27]. Despite this, Nigeria has witnessed a population growth of over 59 % due to a lack of political will in implementing the policy, in addition to multi-ethnicity, and fear of side effects of conventional contraceptives among others.

Most of our respondents used herbal liquid mixtures, popularly called “*agbo*” in Nigeria. Traditionally, herbal medicines are prepared as decoctions, infusions, tinctures, or dried herbs formulated as powders. These forms of preparations are easily prepared and administered, moreover, the traditional herbal practitioners (TMPs) lack the expertise and skills to prepare capsules or tablets, including the lack of standardization required for dosage form design. It is therefore not surprising to see that 55.98 % of our respondents took herbal liquid mixtures, supporting the study conducted by Okoh et al. [11] who reported that Nigerians prefer taking herbal liquid mixtures. Despite the popularity of HMs in Nigeria, knowledge about HCs (19.39 %) is still lacking.

Of the herbs listed in Table 5, *Moringa oleifera* Lam is the most used by both males and females (26.63 %). *M. oleifera* is regarded as a multipurpose plant, cultivated in over 80 countries because of its nutritional properties. It has been used in several cultures and communities for its anti-inflammatory, anti-oxidant, anti-spasmodic, anti-hypertensive, and anti-diuretic properties, and its use dates back to 150 B.C. [28]. Like our study, Balamurugan et al. [29] reported the high use of *M. oleifera* for gynecological disorders, while it was used as an abortifacient in other instances [30].

According to WHO [31], 80 % of Africans rely on herbal medicines. This has been the practice by TMPs, especially in rural areas due to inaccessible healthcare facilities, limited healthcare professionals, and the high cost of medicines. Therefore, they resort to plants for their healthcare needs. As a result, information is passed down from one generation to another especially within families and associates, demonstrating the recommendations made by aged parents (32.07 %), and friends (31.52 %) recorded in our study. Conversely, physicians are more knowledgeable about conventional than HMs. However, because of their perceptions about HMs containing impurities, not being prepared by professionals, and the lack of standardization, their recommendation has been limited. Awodele et al. [32] stated that physicians discourage the use of HMs because of their divergent beliefs from those of TMPs. Our results show this low (15.76 %) level of approval by doctors, hence, the need for R&D into these HCs, while encouraging the formulation of policies for their inclusion in the healthcare delivery system.

There is a sense of male inclusion in decision-making regarding the use of contraceptives. Our study has shown that more males (33.69 %) took HCs than females (33.14 %), with the majority of them within 15-29 years. This shows a willingness of male participation in FP, however, most available contraceptive options are centered on women, with few options (condoms and vasectomy) for the male that may not be ideal in some cases, especially the irreversibility of vasectomy [33]. Hence, these men resort to taking these HCs as they have few side effects and are readily available and recommended by families or friends. This prompts the need for more awareness, staging enlightenment campaigns on male inclusion

in FP, providing more contraceptive options for male contraception through R&D, and the willingness of pharmaceutical industries to invest in the development of these contraceptives.

While our study showed that 70.28 % of respondents are familiar with conventional contraceptives and 14.23 % with HCs, 47.80 % prefer HCs because they are organic, indigenous, affordable, and have low side effects. This shows consistency with previous studies and the fact that 80 % of Africans rely on HMs [11]. Since HMs are prone to adulterations, with crude methods of preparation, limited knowledge of the source of raw materials, and unskilled TMPs, HCs must be developed following guidelines provided by NAFDAC (National Agency for Food Administration and Control) and they must maintain proper surveillance of these vendors, and ensure proper standardization of the products for safety, efficacy, and quality [34], while also encouraging the TMPs to follow the guidelines on good agricultural and collection practices [34,35].

## 5. CONCLUSION

Nigeria is currently the 6<sup>th</sup> largest population in the world and it is expected to reach a population of 377 million people by 2050. This study found that 83.11 % of participants felt the need for population control. A majority (70.28%) are familiar with conventional contraceptives, and few (14.12%) are familiar with HCs, with *M. oleifera* being mostly used. Also, 76.37 % recommended that HCs should be developed, promoted, and used. Therefore, HC awareness campaigns and policy formulations are necessary to ensure its inclusion in the healthcare delivery system.

## CONSENT

Participation in this survey was completely consensual, voluntary, and anonymous.

## ETHICAL APPROVAL

It is not applicable

## CONFLICT OF INTEREST – ALL AUTHORS SHOULD MAKE A CONFLICT-OF-INTEREST STATEMENT PLEASE

ACKNOWLEDGEMENT – ACKNOWLEDGE ANY INDIVIDUAL OR GROUP THAT SUPPORTED YOU IN YOUR STUDY

DECLARATION OF ROLE PLAYED BY EACH CO-AUTHOR SHOULD BE MADE PLEASE

## REFERENCES

1. Gideon HE. Impact of Population Growth on Unemployment in Nigeria. American University of Nigeria, Department of Economics, Senior Research Project. 2017.
2. UN. World Population Prospects. Department of Economic and Social Affairs. 2015. p. 1–66.
3. Messerli P, Murniningtyas E. The future is now. Global Sustainable Development Report. 2019. p. 163–4.
4. Ibidunni SO, Thelma P. Population (Birth) Control: Virtue for Satisfying Classes of Consumers in Nigeria. *Int Bus Manag.* 2017;15(2):25–32.
5. Dare AA, Daniel OG, Emmanuel A, Ari E, Wina F. Survey of traditional family planning methods used in Kogi state, Nigeria. *Afr J Midwifery Womens Health.* 2012;6(4):199–204.

6. Alade G, Oladele A, Okpako E, Ajibesin K, Awotona O. A survey of plants used for family planning in Bayelsa State, southern Nigeria. *J Intercult Ethnopharmacol.* 2018;7(1):1-20.
7. Takyi A, Sato M, Adjabeng M, Smith C. Factors that Influence Modern Contraceptive use Among Women Aged 35-49 Years and their Male Partners in Gomoa West District, Ghana: A Qualitative Study. *Res Sq.* 2021;1-15.
8. Ukwuani-kwaja AN, Nwaogu J, Salako MB. Contraceptive Effect of Methanol Leaves Extract of *Dalbergia saxatilis* on Female Albino Rats. *J Infertil Reprod.* 2021;9(1):35-42.
9. Horibe M, Hane Y, Abe J, Matsui T, Kato Y, Ueda N, et al. Contraceptives as possible risk factors for postpartum depression: A retrospective study of the food and drug administration adverse event reporting system, 2004 – 2015. *Wiley.* 2018;5:131-8.
10. Quijano N. Herbal contraceptives: exploring indigenous methods of family planning. *Initiatives Popul.* 1986;8(222-35):31-2.
11. Ekhaton CN, Osifo UC. Abortifacient Efficacy of *Moringa oleifera* Leave: An Experimental Study on Adult Female Wistar Rats. *Asia Pacific Bus Rev.* 2015;3(6):269-72.
12. Okoh J, Ekere K, Isimi C, Olayemi O, et al. Popularity and Customer Preferences for Herbal Medicines in Nigeria: A Questionnaire Based Survey. *Humanit Soc Sci Lett.* 2016;4(3):69-76.
13. World Health Organization (WHO). WHO guidelines on safety monitoring of herbal medicines in pharmacovigilance systems. World Health Organisation, Geneva. 2004. p. 1-82.
14. Pradhan DK, Mishra MR, Mishra A, Panda AK, Behera RK, Jha S, et al. A comprehensive review of plants used as contraceptives. *Int J Pharm Sci Res.* 2012;4(1):148-55.
15. Igwillio UC, Nwokeke CC, Nwachukwu VC, Inchikida BM, Kure TK, Danjuma N, et al. Production and Development of Herbal Contraceptives in Nigeria – A Necessity. *Int J Recent Res Life Sci.* 2020;7(4):33-43.
16. Adedeji O, Olusola A, James G, Shaba HA, Orimoloye IR, Singh SK, et al. Early warning systems development for agricultural drought assessment in Nigeria. *Environ Monit Assess.* 2020;192(12):1-21.
17. Kana MA, Tagurum YO, Hassan ZI, Afolanranmi TO, Ogbeyi GO, Difa JA, et al. Prevalence and determinants of contraceptive use in rural Northeastern Nigeria: Results of a mixed qualitative and quantitative assessment. *Ann Niger Med.* 2016;10(1):3-10.
18. Ezenwaka U, Mbachu C, Ezumah N, Eze I, Agu C, Agu I, et al. Exploring factors constraining utilization of contraceptive services among adolescents in Southeast Nigeria: An application of the socio-ecological model. *BMC Public Health.* 2020;20(1):1-11.
19. Avidime S, Aku-akai L, Mohammed AZ, Adaji S, Ejembi SC. Fertility Intentions, Contraceptive Awareness and Contraceptive Use among Women in Three Communities in Northern Nigeria. *African J Reprod Heal Sept.* 2010;14(3):65-70.
20. Utoo BT, Mutahir TJ, Utoo PM. Knowledge, attitude and practice of family planning methods among women attending antenatal clinic in Jos, North-central Nigeria. *Niger J Med.* 2010;19(2):214-8.
21. Eko EJ, Osonwa K, Osuchukwu CN, Offiong AD. Prevalence of Contraceptive use among women of reproductive. *Int J Humanit Soc Interv.* 2013;2(6):27-34.
22. Adebisi IM, Bello SO. An ethnobotanical survey of herbal male contraceptives used in South-West Nigeria. *African J Pharm Pharmacol.* 2011;5(2):289-91.
23. WHO. Contraception. *Br Med J.* 2019;2(3892):265-7.
24. Odusina EK, Ugal DB, OO. Socio-Economic Status, Contraceptive Knowledge And Use Among Rural Women In Ikeji Aakeji, Osun State, Nigeria. *Afro Asian J Soc Sci.* 2012;3(3):1-10.

25. Saka MJ, Saidu R, Balogun OR, Raji HO, Ijaiya MA, Saka AO, et al. Contraceptive Use Among Adolescent In University Of Ilorin Teaching Hospital.pdf. 2011. p. 50–2.
26. NPP. National Population Policy & Family Planning. 2022. p. 1–4.
27. Turnwait O, Mayowa A. Nigeria's Population Policies: Issues, Challenges and Prospects. Ibadan J Soc Sci. 2017;15(1):1–13.
28. Razis AFA, Ibrahim MD, Kntayya SB. Health benefits of Moringa oleifera. Asian Pacific J Cancer Prev. 2014;15:8571–6.
29. Balamurugan S, Vijayakumar S, Prabhu S, Morvin Yabesh JE. Traditional plants used for the treatment of gynaecological disorders in Vedaranyam taluk, South India - An ethnomedicinal survey. J Tradit Complement Med. 2018;8(2):308–23.
30. Magtalas MC, Balbin PT, Cruz EC, Guevarra RC, Cruz ARDP, Silverio CE, et al. A systematic review of ethnomedicinal plants used for pregnancy, childbirth, and postpartum care in the Philippines. Phytomedicine Plus. 2023;3(1):1–15.
31. WHO. The WHO strategy for traditional medicine: Review of the global situation and strategy implementation in the Eastern Mediterranean Region. World Health Organization. 2002. p. 1–14.
32. Awodele O, Agbaje EO, Abiola OO, Awodele DF, Dolapo DC. Doctors' attitudes towards the use of herbal medicine in Lagos, Nigeria. J Herb Med. 2012;2(1):16–22.
33. Amory JK. Development of Novel Male Contraceptives. Clin Transl Sci. 2020;13:228–37.
34. NAFDAC. Herbal Medicines and Related Products (Registration) Regulations. Fed Repub Niger Off Gaz. 2021;108(109):1–7.
35. FOA. Guidelines "Good Agricultural Practices for Family Agriculture". Departmental Program on Food and Nutritional Security. 2007. 1–60 p.