

# Effect of Weed Management Practices on Growth, Yield Attributes and Yield of Clusterbean [*Cyamopsistetragonoloba* (L.)Taub]

## ABSTRACT

The growth, yield attributes and yield of clusterbean were influenced by various weed management treatments. The objective of the current study was to evaluate the impact of different weed management practices on growth, yield attributes and overall yield. An agronomic investigation was conducted at the Agricultural Research Station, Mandor, affiliated with the Agriculture University, Jodhpur, Rajasthan, during the *Kharif* season of 2018. The experimental design utilized a randomized block design (RBD) with three replications. The experiment encompassed ten treatments: W<sub>1</sub> - weedy check, W<sub>2</sub> - weed-free, W<sub>3</sub> - pendimethalin at 750 g/ha (pre-emergence), W<sub>4</sub> - imazethapyr at 40 g/ha (early post-emergence), W<sub>5</sub> - pendimethalin + imazethapyr at 750 g/ha (pre-emergence), W<sub>6</sub> - imazethapyr + imazamox at 50 g/ha (early post-emergence), W<sub>7</sub> - pendimethalin at 750 g/ha (pre-emergence) + 1 hand weeding (25 days after sowing), W<sub>8</sub> - imazethapyr at 40 g/ha (early post-emergence) + 1 hand weeding (35 days after sowing), W<sub>9</sub> - pendimethalin + imazethapyr at 750 g/ha (pre-emergence) + 1 hand weeding (25 days after sowing), and W<sub>10</sub> - imazethapyr + imazamox at 50 g/ha (early post-emergence) + 1 hand weeding (35 days after sowing). The early post-emergence application of imazethapyr + imazamox at 50 g/ha along with one hand weeding at 35 days after sowing resulted in significantly enhanced growth, yield attributes and overall yield. This treatment exhibited higher values for parameters such as dry matter per plant (g), plant height (cm), number of branches per plant, number of pods per plant, number of seeds per pod, 1000-seed weight (g), yields of both seed and stover and harvest index. Additionally, this treatment was comparable in effectiveness to the application of imazethapyr at 40 g/ha (early post-emergence) + 1 hand weeding at 35 days after sowing and the weed-free condition. The application of imazethapyr + imazamox at 50 g/ha + 1 hand weeding at 35 days after sowing emerged as a promising strategy during the *Kharif* season for achieving the highest yield and net monetary returns in clusterbean cultivation.

*Keywords:* Clusterbean, growth attributes, yield attributes and yield

## 1. INTRODUCTION

Clusterbean (*Cyamopsistetragonoloba* (L.)Taub), commonly referred to as "Guar," is a significant crop native to India and belongs to the family **Fabaceae**, subfamily **Papilionaceae**. "It thrives in rainfed regions, particularly the semi-arid and arid zones of India. This crop is recognized for its flower and pod clusters, earning it the name "clusterbean." Its primary origin can be traced back to India" (Vavilov, 1951). "In the **State** of Rajasthan, where conditions are well-suited, clusterbean cultivation is prominent, covering approximately 2.39 million hectares and yielding around 1.0 million **tones**, with a productivity of 419 kg/ha" (Anon., 2022). "Clusterbean serves various purposes such as vegetable consumption, green fodder, green manure, and seed production. The seeds contain a significant gum content (28-33**percent**), finding utility in various industries including textiles, paper, petroleum, pharmaceuticals, food processing, cosmetics, mining explosives, and oil drilling, thus contributing to foreign exchange earnings" (Kumawat *et al.*, 2017). "In the context of Indian farming, pulses play a crucial role, serving as the main crop, catch crop, cover crop, green manure, and intercrop. However, low

yields are often attributed to factors such as poor cultivation practices, cultivation on marginal lands, insufficient fertilization, monsoon-dependence, and lack of suitable varieties. Among these challenges, weed infestation stands out as a significant hurdle in clusterbean cultivation. The crop faces intense competition from weeds due to its slow initial growth and the rapid growth of weeds in the rainy season. Extended weed presence beyond the critical crop-weed competition phase leads to substantial yield reductions, ranging from 40-45percent” (Sangwan *et al.*, 2016). Studies have shown that “crop-weed competition contributed to a 53.7percent reduction in seed yield” (Singh *et al.*, 2008). “While hand weeding remains a traditional and effective option for weed management, challenges such as labor availability and rising labor costs limit its economic feasibility” (Singh and Godara, 2015). “A combination of chemical and cultural weed control methods can mitigate herbicide expenses while ensuring adequate aeration and moisture conservation for the crop” (Prakash *et al.*, 1991). While pre-emergence application of pendimethalin is recommended, the study also explores post-emergence herbicide application for effective control of mixed weed flora. The investigation aims to identify an appropriate weed management practice to control weeds in clusterbean cultivation.

## 2. MATERIALS AND METHODS

**Soil Characterization:** The soil's key elements (N, P, K) were analyzed using the alkali potassium permanganate method (Subbiah and Asija, 1956), Olsen method (Olsen, 1954), and flame photometer method (Jackson, 1973) for nitrogen, phosphorus and potassium, respectively. pH was determined using the method outlined by Richards (1954) and organic carbon was assessed following the method by Walkley and Black (1934). Homogeneous composite soil samples were collected for the respective analyses. The soil exhibited a loamy sand texture, a slightly alkaline pH (8.2), low organic carbon content (1.3 g/kg), medium available nitrogen (174.0 kg/ha), medium available phosphorus (22.0 kg/ha), high availability of potash (325.0 kg/ha) and an electrical conductivity of 0.12 ds/m.

**Field Investigation:** During the *Kharif* season of 2018, a field investigation was conducted at the Agricultural Research Station, Mandor (Agriculture University, Jodhpur), situated between 26<sup>0</sup>15' N to 26<sup>0</sup>45' north latitude and 73<sup>0</sup>00' E to 73<sup>0</sup>29 East longitude at an altitude of 231 m above mean sea level within the western dry region agro-climatic zone. The weather conditions during the crop season are illustrated in Fig. 1. The experiment employed a randomized block design (RBD) with three replications. It included ten treatments: W<sub>1</sub> - weedy check, W<sub>2</sub> - weed-free, W<sub>3</sub> - pendimethalin at 750 g/ha (pre-emergence), W<sub>4</sub> - imazethapyr at 40 g/ha (early post-emergence), W<sub>5</sub> - pendimethalin + imazethapyr at 750 g/ha (pre-emergence), W<sub>6</sub> - imazethapyr + imazamox at 50 g/ha (early post-emergence), W<sub>7</sub> - pendimethalin at 750 g/ha (pre-emergence) + 1 hand weeding (25 days after sowing), W<sub>8</sub> - imazethapyr at 40 g/ha (early post-emergence) + 1 hand weeding (35 days after sowing), W<sub>9</sub> - pendimethalin + imazethapyr at 750 g/ha (pre-emergence) + 1 hand weeding (25 days after sowing) and W<sub>10</sub> - imazethapyr + imazamox at 50 g/ha (early post-emergence) + 1 hand weeding (35 days after sowing). The test crop used was clusterbean variety RGC-1017, sown in rows spaced 30 cm apart at a seed rate of 15 kg/ha. The recommended nutrient dose included 10 kg N and 40 kg P<sub>2</sub>O<sub>5</sub>/ha applied using urea and DAP. Basal application covered the entire dose of nitrogen and phosphorus. While adhering to the recommended practices of the zone, the cultivation practices were followed, with the exception of weed management.

**Plant Growth and Yield Analysis:** Plant growth parameters such as dry matter per plant (g), plant height (cm), number of branches per plant, number of pods per plant, and number of seeds per pod were manually recorded by selecting five representative plants from each plot in each replication. Yield attributes and actual yields were also recorded, with seed and stover yields measured from the net plot area of each treatment. The collected data underwent comprehensive statistical analysis, including treatment means, standard error means, critical differences, and ranges of variation. The F-test for significance was also conducted for each parameter. The entire dataset was compiled and statistically analyzed using the established procedures for a randomized block design as suggested by Gomez and Gomez (1984).

**Application of treatments :** Pendimethalin and its combination with imazethapyr were applied as pre-emergence treatments within 2 DAS, while imazethapyr and its combination with imazamox were applied as early post-emergence treatments at 20 DAS. Herbicides were sprayed using a knapsack sprayer at a rate of 800 liters/ha. Hand weeding was conducted in designated plots at 25 and 35 DAS according to treatment specifications.

### 3. RESULTS AND DISCUSSION

The major weed flora founded throughout crop season revealed that weedy check plots were heavily invaded by broad leaved weeds chiefly *Digeraarvensis*, *Tribulusterristris*, *Corchorustrilocularis*, *Phyllanthusniruri*, *Amaranthusviridis*, *Celosia argentea* and *Portulacaoleracea* and *Eragrostis minor*, *Cyperusrotundus*, *Cynodondactylon* and *Dactylocteniumaegyptium* among narrow leaved weeds. *Digeraarvensis* and *Tribulusterristris* were found to be the prominent broad leaved weeds.

#### 3.1 Effect on growth attributes:

In the present study, the impact of different weed management treatments on the growth attributes and yield of clusterbean was investigated. The findings revealed a significant enhancement in the growth parameters, such as dry matter per plant, plant height, and number of branches per plant, due to the application of weed management treatments during the crop season. Specifically, the dry matter per plant increased significantly with the application of the herbicide imazethapyr + imazamox at a rate of 50 g/ha. Interestingly, when this treatment was combined with one hand weeding at 35 days after sowing (DAS), the plant's dry matter further improved by 28.8%. Moreover, the plant height was significantly taller (65.1 cm) with the application of imazethapyr + imazamox at 50 g/ha along with one hand weeding at 35 DAS. This was on par with the combination of imazethapyr at 40 g/ha + one hand weeding at 35 DAS (63.6 cm) and weed-free conditions (65.8 cm). The treatment of imazethapyr + imazamox at 50 g/ha + one hand weeding at 35 DAS recorded a remarkable increase in plant height by 22.8% and 73.6% compared to the sole application of imazethapyr + imazamox at 50 g/ha and the weedy check, respectively. Further, treatments like imazethapyr + imazamox at 50 g/ha + one hand weeding at 35 DAS and imazethapyr at 40 g/ha (early post-emergence) + one hand weeding at 35 DAS exhibited promising results in enhancing the growth attributes of clusterbean, with 82.5% and 81.3% higher branches per plant at the harvest stage compared to weedy check plots. The presence of less crop-weed competition, as evidenced by the lowest population of narrow and broad leaf weed species, along with lower weed dry weight, led to improved growth attributes. This allowed the crop to efficiently utilize nutrients, moisture, light, and space, contributing to better growth and development. Previous studies by Yadav et al., (1999) and Sharma and Singh (2003) also supported the idea that "effective weed management practices can significantly increase plant height and the number of branches per plant. Furthermore, weed control was linked to improved moisture, nutrient availability, and solar radiation for the crops. This resulted in

higher chlorophyll content, photosynthetic rates, and nitrate reductase activity, ultimately promoting greater carbohydrate production and growth". Similar improvements in growth characteristics due to imazethapyr + imazamox at 40 g/ha were noted by Singh *et al.*, (2016) and Sharma *et al.*, (2017) in clusterbean.

### 3.2 Effect on yield attributes and yield:

Moving on to yield attributes, the treatment involving early post-emergence application of imazethapyr + imazamox at 50 g/ha along with one hand weeding at 35 DAS recorded the highest number of pods per plant (45.3). This was significantly superior to other treatments except for imazethapyr at 40 g/ha + one hand weeding at 35 DAS. The same treatment also led to the highest number of seeds per pod (7.5), showing a 44.2% increase over the control. Additionally, this treatment resulted in the highest 1000-seed weight (33.5 g), which was on par with imazethapyr at 40 g/ha + one hand weeding at 35 DAS. The yield was significantly improved with various weed management treatments compared to the control, with the highest seed yield (1180 kg/ha) achieved by the imazethapyr + imazamox at 50 g/ha + one hand weeding at 35 DAS treatment. Alone application of imazethapyr + imazamox 50 g/ha recorded stover yield of 2670 kg/ha which was significantly higher over weedy season long plots by 316.5 per cent. When this treatment was super imposed with one hand weeding at 35 DAS, the stover yield was significantly further improved by 25.1 percent over application of imazethapyr + imazamox 50 g/ha. These results highlight that effective weed management strategies can substantially impact the growth attributes and yield of clusterbean. The improvement in the yield components of the clusterbean could be attributed to increase in the growth characters of the crop. The combination of herbicidal treatments with one hand weeding at the appropriate stage appears to be a promising approach in enhancing both growth and yield. The findings align with previous research by Dungarwal *et al.*, (2002), Sharma and Singh *et al.*, (2003), Kumar *et al.*, (2014), Singh *et al.*, (2015), Singh *et al.*, (2016), Sharma *et al.*, (2017) and Yadav and Mundra (2017) in the context of clusterbean cultivation.

### 4. CONCLUSION

Based on one year experiment, it may concluded that early post-emergence application of imazethapyr + imazamox at 50 g/ha+ 1 HW at 35 DAS recorded higher growth, yield attributes and seed yield of clusterbean but equally effective with imazethapyr at 40 g/ha+ 1 HW at 35 DAS. Keeping all views in mind, imazethapyr + imazamox at 50 g/ha+ 1 HW at 35 DAS may be viable and taken for further research.

### REFERENCES

- Anonymous 2022, Rajasthan. Directorate of Economics & Statistics. Final Estimates of Area, Production and Yield of Crops in Respect of Rajasthan State/U.T. for the Year 2021-22.
- Dhaker H, Mundra SL, Jain NK. Weed management in clusterbean [*Cyamopsistetragonoloba* (L.) Taub.]. *Indian Journal of Weed Science*. 2009;41(3and4):224-7.
- Dungarwal HS, Chaplot PC, Nagda BL. Chemical weed control in clusterbean (*Cyamopsistetragonoloba* L.). *Indian Journal of weed Science*. 2002;34(3&4):208-212.
- Gomez KA, Gomez AA. *Statistical Procedure for Agricultural Research*. John Willey and Sons, 2nd edition p329. 1984.
- Jackson ML. *Soil Chemical Analysis (II Edition)*. Prentice Hall of India Private Limited. New Delhi, India. 1973.

- Kumar N, Hazra KK, Nadarajan. Weed management strategies through post-emergence herbicides in pulses. Biennial Conference of Indian society of Weed Science on Emerging Challenges of Weed Management , DWSR, Jabalpur, February 15-16, 2014. pp 28.
- Kumawat P, Kaushik MK, Meena VK, Chouhan BS, Meena RK, Kumar R. Effect of weed management and fertility levels on productivity of clusterbean [*Cyamopsistetragonoloba* (L.) Taub]. Legume Research. 2017;40(5):884-889.
- Olsen SR. Estimation of available phosphorus in soils by extraction with sodium bicarbonate. USDA: Washington. Circular, 939. 1954.
- Prakash, Prasad K, Singh P. Chemical weed control in soybean. Indian Journal of Weed Science. 1991; 23(1&2):29-31.
- Richards LA. Diagnosis and improvement of saline and alkali soil. USDA Hand Book No. 60 Oxford & IBH Publication Company, New Delhi. 1954.
- Sangwan M, Singh S, Satyavan. Efficacy of sequential application of imazethapyr + imazamox and propaquizafop in clusterbean (*Cyamopsistetragonoloba*) in two texturally different soils. Indian Journal of Agronomy. 2016;61:519-52.
- Sharma K, Rawat GS, Gaur D, Sharma A. Effect of post-emergence herbicides on weed control, growth and yield of clusterbean [*Cyamopsistetragonoloba* (L.) Taub] in M.P. Agricultural Science Digest. 2017;37(3):179-184.
- Sharma RP, Singh P. Effect of weed management and phosphorus levels on yield and quality of clusterbean (*Cyamopsistetragonoloba* L.). Annals of Agricultural Research. 2003;24(3):605-609.
- Singh R, Dhillon BS. Evaluation of different weed management strategies in clusterbean [*Cyamopsistetragonoloba* (L.) taub] under rainfed conditions of Punjab. Legume Research-An International Journal. 2023;46(1):112-8.
- Singh R, Godara AS. Weed control efficiency of post-emergence herbicides and their effect on productivity of clusterbean [*Cyamopsistetragonoloba* (L.) Taub]. Legume Research. 2015; 38(3):415-418.
- Singh SK, Jain AK, Puniya BL. Integrated weed management in clusterbean [*Cyamopsistetragonoloba* (L.)]. Indian Journal of Agricultural Sciences. 2008;70(2):850-852.
- Singh SP, Yadav RS, Gupta V. Yield performance and nutrient uptake as influenced by integrated weed management in clusterbean. Indian Journal of Weed Science. 2015;47(1):82-84.
- Singh SP, Yadav RS, Sharma V. Weed control in clusterbean through post-emergence herbicides. Indian Journal of Weed Science. 2016;48(2):202-205.
- Singh SP, Yadav RS, Sharma V. Weed control in clusterbean through post-emergence herbicides. Indian Journal of Weed Science. 2016;48(2):202-205.
- Subbiah B, Asija G. 1956. A rapid procedure for the estimation of available nitrogen in soils. Current Science. 1956;25:259-60.
- Vavilov NI. The origin, variation, immunity and breeding of cultivated plants. Chronic Botanica 1951;13:1-54.
- Walkley A, Black IA. An examination of the Degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method. Soil science. 1934;37(1):29-38.

Yadav RK, Mundra SL. Weed management and sulphur nutrition in clusterbean for higher productivity and profitability. *Journal of Pharmacognosy and Phytochemistry*. 2017;6(3): 06-08.

Yadav RP, Shrivastava UK, Dwivedi SC. Comparative efficiency of herbicides in controlling *Cyperusrotundus* and other weeds in clusterbean (*Cyamopsistetragonoloba* (L.)). *Indian Journal of Agronomy*. 1999;44(1):151-155.




Fig 1: Meteorological data recorded at ARS, Mandor-Jodhpur during crop season

Table 1. Effect of different weed management treatments on growth attributes of clusterbean

Treatments	Plant population (Lakh/ha)		Dry matter/plant (g)	Plant height (cm)	Number of branches (/plant)
	Initial	Final			
Weedy check	3.12	1.91	12.2	37.5	3.60
Weed free	3.20	3.18	35.2	65.8	7.67
Pendimethalin 750 g/ha (pre-em.)	3.18	2.25	23.3	44.3	4.47
Imazethapyr 40 g/ha(early post-em.)	3.11	2.27	23.6	45.6	4.53
Pendimethalin + imazethapyr 750 g/ha (pre-em.)	3.17	2.31	24.7	46.0	5.33
Imazethapyr + imazamox 50 g/ha (early post-em.)	3.15	2.69	27.0	53.0	5.40
Pendimethalin 750 g/ha (Pre-em.) + 1 HW at 25 DAS	3.19	2.77	27.4	53.4	5.47
Imazethapyr 40 g/ha (early post-em.) + 1 HW at 35 DAS	3.16	3.07	32.5	63.6	6.53
Pendimethalin + imazethapyr 750g/ha (pre-em.)+ 1 HW at 25 DAS	3.19	2.78	28.7	55.5	6.47
Imazethapyr + imazamox 50 g/ha (early post-em.) + 1 HW at 35 DAS	3.19	3.11	34.8	65.1	6.57
S Em ±	0.12	0.11	0.78	2.07	0.28
CD (P=0.05)	NS	0.34	2.33	6.16	0.86

Table 2. Effect of different weed management treatments on yield attributes, yield and harvest index of clusterbean

Treatments	Pods/plant (No.)	Seeds/pod (No.)	1000-seed weight (g)	Seed yield (kg/ha)	Stover yield (kg/ha)	Harvest Index (%)
Weedy check	22.0	5.2	26.4	210.0	641.0	24.67
Weed free	48.2	7.7	34.0	1237.0	3503.0	26.09
Pendimethalin 750 g/ha (pre-em.)	29.4	5.8	29.5	694.0	2097.0	24.86
Imazethapyr 40 g/ha (early post-em.)	31.0	5.8	29.7	713.0	2140.0	24.99
Pendimethalin + imazethapyr 750 g/ha (pre-em.)	31.5	5.9	29.8	752.0	2257.0	24.99
Imazethapyr + imazamox 50 g/ha (early post-em.)	35.9	6.7	30.6	893.0	2670.0	25.06

Pendimethalin 750 g/ha (pre-em.) + 1 HW at 25 DAS	38.6	6.7	30.8	903.0	2711.0	25.01
Imazethapyr40 g/ha (early post-em.) + 1 HW at 35 DAS	43.4	7.4	33.2	1120.0	3240.0	25.68
Pendimethalin + imazethapyr750g/ha (pre-em.)+ 1 HW at 25 DAS	38.6	6.8	31.0	975.0	2923.0	25.01
Imazethapyr + imazamox 50 g/ha (early post-em.) + 1 HW at 35 DAS	45.3	7.5	33.5	1180.0	3340.0	26.10
S Em ±	1.39	0.19	0.96	44.23	106.02	0.96
CD (P=0.05)	4.16	0.59	2.88	131.44	315.03	NS