

Short Research Article **Enersol promotes growth and yield performance of pechay (*Brassica rapa* L. var. *chinensis* (L.) Hanelt)**

ABSTRACT

This study presents the results of a field bio-efficacy evaluation to assess the impact of Enersol® LDG, a mineral fertilizer containing leonardite and humic acid, on the growth and yield of pechay (*Brassica rapa*) in Puypuy, Bay, Laguna, Philippines. Pechay is a common leaf vegetable in the Philippines, known for its mild flavor and short duration crop cycle. In organic pechay production, there is a need for a plant food supplement that facilitates immediate nutrient absorption. An experiment in a randomized complete block design with three replications was conducted, with different combinations of Enersol® LDG and inorganic fertilizer applied basally at seed sowing. Parameters, including total fresh yield, marketable yield, non-marketable yield, plant height, number of leaves, and leaf width, were recorded and analyzed statistically. Results showed that the combination of Enersol® LDG and inorganic fertilizer significantly increased the total fresh yield of pechay compared to other treatments. However, reducing the amount of inorganic fertilizer resulted in lower yields, regardless of the Enersol® LDG application rate. The marketable yield and percent marketable yield were also significantly higher in plants treated with Enersol® LDG and inorganic fertilizer combinations compared to the control and Enersol® LDG alone. The percentage of non-marketable yield was highest in plants treated with Enersol® LDG alone. Based on these findings, it can be concluded that Enersol® LDG, when combined with inorganic fertilizer, significantly enhanced the growth and yield of pechay. Further research is recommended to explore optimal dosage and application methods for maximizing pechay production using Enersol® LDG.

Keywords: Enersol, humic acid, leonardite, low dust granules, pechay

1. INTRODUCTION

Pechay (*Brassica rapa* L. var. *chinensis* (L.) Hanelt) group cultivars, is an erect, biennial herb, cultivated as an annual about 15–30 cm tall in vegetative stage. Leaves are ovate arranged spirally and spreading, while the petioles are enlarged and grow upright forming a near-cylindrical bundle (Jimenez et al., 2021). The inflorescence is cone-shaped with pale yellow flowers. The seeds are 1 mm in diameter and are somewhat maroon in color (Gawryś, 2008). Pechay is an important component of Filipino food, such as “puchero”, “ginisa” and “nilaga.” It is a green leafy vegetable rich in calcium and other important nutrients. Its nutritional values are made up of 93.0 g water, 1.7 g protein, 0.2 g fat, 3.1 g carbohydrates, 0.7 g fiber, 0.8 g ash, 2.3 g β-carotene, 53.0 mg vitamin C, 102.0 mg calcium, 46.0 mg phosphorous, 2.6 mg iron, and an energy of 86.0 kJ. Pechay is used for its immature, but fully expanded tender leaves (Knot, Deanon, 1967). The succulent petioles

are used as main ingredient for soup and stir-fried dishes, while its green petioles and leaves are also used as garnish in Chinese cuisine (Jim, Tony, 2006; Jimenez et al., 2021).

Pechay is one of the top twenty vegetables and an important source of food globally, according to the Food and Agricultural Organization (FAO). The country's Chinese pechay production slightly decreased by 0.95% in 2017. Approximately 86.3% of the country's total *B. chinensis* production comes from the Cordillera Administrative Region, while Central Visayas comes next with a 7.0% share. The remaining 6.7% comes from a combined share from Northern Mindanao, Davao Region, and the rest of the country (Philippine Statistics Authority, 2019). Therefore, the root of this type of changes are being investigated.

Several studies in using various fertilizers and foliar supplements have been conducted to determine their bio-efficiency in maximizing the growth and yield of various crops. In pummelo plants (*Citrus maxima* Merr.) the use of fertilizer from fermented banana peel, nutrition by foliar or soil fertilization with potassium was investigated by (Magbalot-Fernandez, De Guzman, 2019; 2021). The effect of hormones and Bio-Forge fertilizer were analyzed (Fernandez, Sabay, 2016; Magbalot-Fernandez et al., 2020) in abaca (*Musa textilis* Nee) and banana (*M. paradisiaca* L.) plants. For rice (*Oryza* L.) the effect of organic-based foliar fertilizer was checked (Montifalcon, Fernandez, 2017). In pechay (*B. chinensis*) the effect of organic foliar fertilizer or the stimulating effect of hormones was examined by (Fernandez, Miñoza, 2015; Fernandez, Andigan, 2017). Just recently, Adorada et al. (2023) demonstrated the field efficacy of a seaweed-based foliar fertilizer in enhancing the growth and yield of pechay.

This study aimed to evaluate the bio-efficacy of Enersol Low Dust Granules (LDG) on the growth and yield of pechay under farmer's field conditions and being a newly introduced product.

2. MATERIAL AND METHODS

The trial was conducted for one season at Puypuy, Bay, Laguna Philippines from September-October 2019. An open-pollinated pechay cultivar, Pavito was used as the test crop/cultivar. Prior to preparing the seedbeds, surface (0-15 cm) soil samples were randomly collected from five different points on the trial site for physical and chemical analysis. Complete soil physical and chemical analyses were requested from an accredited soil testing laboratory. The field for the trial was plowed and harrowed twice in different directions. Seedbeds were prepared manually, and eighteen plots measuring 2m x 5m were made.

2.1 Soil Analysis. Before bed preparation, surface (0-15 cm) soil samples were randomly collected from five different points on the experimental site for physical and chemical analysis. A complete soil physical and chemical analysis was requested from UPLB ASC Soils Laboratory. Organic matter, nitrogen, phosphorus, and potassium levels were determined while moisture and pH were also measured.

2.2 Land Preparation. The experimental field was plowed and harrowed twice in different directions. Bed preparations were done manually using spades. Plots measuring 2x5 m² were made. A total of 18 plots were established.

2.3 Experimental Design and Treatment Application. The experiment followed a randomized complete block design with three replications (Figure 1). The inorganic fertilizer and Enersol® LDG, based on the treatment combinations, were applied basally during the seed sowing. Table 1 shows the different treatments used in the bioefficacy trial.

Table 1. Treatment assignments for the bioefficacy trial

Treatments	Granular Fertilization	Enersol LDG
1	Inorganic Fertilizer alone @ 8.5 bags 14-14-14 & 4 bags urea/ha	
2		1rr Enersol® @ 50 kg/ha
3	Inorganic Fertilizer @ 8.5 bags 14-14-14 & 4 bags urea/ha	1rr Enersol® @ 50 kg/ha
4	½ Inorganic Fertilizer @ 4.25 bags 14-14-14 & 2 bags urea/ha	1rr Enersol® @ 50 kg/ha
5	½ Inorganic Fertilizer @ 4.25 bags 14-14-14 & 2 bags urea/ha	2rr Enersol® @ 100 kg/ha
6	Control	




Figure 1. Field layout using a randomized complete block design with three replications.

2.4 Seed Sowing. Five rows per meter were line drilled covering the 5-meter stretch of the plot. The distance between rows is 20cm. Pechay seeds were directly sown into the soil using drill method at 1 gram per 2m². The sown seeds were covered with soil. The plots were dusted with carbaryl to prevent ant foraging.

2.5 Cultural Management. The following cultural practices were implemented as needed:

2.5.1 Irrigation. Watering was done every day depending on soil moisture content and weather condition.

2.5.2 Weeding. Weeds were removed at weekly intervals to avoid competition with the crop.

2.5.3 Cultivation. Tilling the soil was done to aerate the soil and to control some weed pests.

2.5.4 Disease Control. No disease was observed in the experimental plants.

2.5.5 Insect Pest Control. Insect infestation was managed by applying insecticides appropriate to the target insect pest.

2.6 Harvesting. The pechay plants were harvested ~28 days after planting. The harvested plants were sorted into either marketable or non-marketable. Non-marketable yield was classified based on plants whose leaves had holes ≥ 1 cm in diameter.

2.7 Data Gathering. Fresh weight yield of the above ground biomass plants was expressed in kg/ha and recorded. In addition, the non-marketable yield was also expressed in percent (%) of the total (marketable + non-marketable) yield. Growth parameters such as plant height, number of leaves and leaf width were also gathered from a sample of thirty plants per treatment (10/rep).

Data gathering procedure:

2.7.1 Fresh weight yield. Total yield was derived from matured plants harvested per treatment per replication. The plants for each plot were weighed using a digital weighing scale.

2.7.2 Percent marketable and non-marketable yield. This was computed after separating the marketable from non-marketable plants. Non-marketable plants are plants which are deformed, small and damaged by insects or other organisms (≥ 1 cm holes).

2.7.3. Plant height. The distance between the base of the plant and the longest/highest leaf were measured using a ruler after harvesting. A sample of ten plants per replication was measured.

2.7.4. Number of leaves. The number of mature leaves not smaller than one-inch diameter was counted. A sample of ten plants per replication was measured.

2.7.5 Leaf width. This was done by getting ten sample plants per replication and measuring the biggest leaf for each plant.

2.8 Data Analysis and Interpretation. The generated data were subjected to statistical analysis using SAS portable version. Analysis of variance and treatment mean comparison using Tukey's Test were done at 5% probability level. Two-way tables were prepared to show the means and their corresponding alphabet notations. Interpretations were done by comparing each treatment, especially with the control. Conclusions and recommendations were formulated based on the results generated.

3. RESULTS AND DISCUSSION

3.1. Total Fresh Yield of Pechay

The total yield was derived from the 20 m² plot area and converted to fresh yield per hectare. The highest fresh yield was obtained from the plants treated with the 1rr Enersol® and inorganic fertilizer. However, this is not significantly different with the plants treated with other combinations of Enersol® and inorganic fertilizer at any rate. All Enersol® and inorganic fertilizer combinations are significantly different with the plants treated with Enersol® alone and the control plants. On the other hand, the lowest fresh yield was obtained from the control plants but is not significantly different from the plants treated with Enersol® alone (Table 2).

Table 2. Total fresh yield of pechay treated with Enersol® and inorganic fertilizer combinations.

TREATMENT	MEAN YIELD (tons/ha)			MEAN
	1	2	3	
1 Inorganic fertilizer alone	89.0	76.4	68.4	77.93 A
2 1rr Enersol® alone	40.0	33.4	28.5	33.97 B
3 1rr Enersol® + inorganic fertilizer	91.0	95.0	66.6	84.20 A
4 1rr Enersol® + ½ inorganic fertilizer	70.0	87.8	51.4	69.73 A
5 2rr Enersol® + ½ inorganic fertilizer	84.6	72.4	66.2	74.40 A
6 Control	21.0	29.9	28.1	26.33 B

cv 19.11816, p=.0002 *Means with the same letter are not significantly different at $\alpha = 0.05$

The results revealed that pechay are responsive to inorganic fertilization in combination with Enersol®, which gave a yield increase of 6.27 and 57.87 tons per hectare over the granular inorganic fertilization alone and the control, respectively. However, lowering the amounts of inorganic fertilizer in half would give lower yields regardless of the amount of Enersol® applied. This only supported the fact that Enersol® is only effective when combined with the full dose inorganic fertilizer in which the soil enhancer product promotes efficient nutrient absorption by improving soil condition.

3.2. Number and Percent Marketable Yield of Pechay

The marketable yield of pechay was gathered by selecting excellent quality undamaged whole plants of pechay wherein the percentage marketable yield was computed out of these values (Table 3). The highest marketable yield was obtained in the plants treated with 1rr Enersol® and inorganic fertilizer combination while the lowest marketable yield was derived from the control plots which conformed to the trend in the total fresh yield data. The results revealed that at harvest period (30 days after sowing), all Enersol® and inorganic fertilizer combination treated plants, either full or half dose inorganic when in combination with any rate of Enersol®, gave significant difference with the control plants. Only the plants treated with Enersol® alone are comparable with the control plants and the plants treated with 1rr Enersol® and ½ inorganic fertilizer combination. Both the plants treated with Enersol® alone and the control plants are distinctively light green in color and smaller, making them unacceptable in the market.

In terms of percentage, the highest marketable percentage is obtained in 1rr Enersol® + inorganic fertilizer, followed by inorganic fertilizer alone, 2rr Enersol® + ½ inorganic fertilizer, 1rr Enersol® + ½ inorganic fertilizer, 1rr Enersol® alone and finally the control plants.

Table 3. Marketable yield and percent marketable yield of pechay treated with Enersol® and inorganic fertilizer combinations.

TREATMENT	MARKETABLE YIELD (kg/ha, %)			MEAN
	1	2	3	
1 Inorganic fertilizer alone	84,550.00 (95.00)	76,400.00 (100.00)	66,587.40 (97.35)	75,845.80 A (97.35)
2 1rr Enersol® alone	37,396.00 (93.49)	24,408.70 (73.08)	24,467.30 (85.85)	28,757.32 BC (84.14)
3 1rr Enersol® + inorganic fertilizer	91,000.00 (100.00)	89,300.00 (94.00)	66,600.00 (100.00)	82,300.00 A (98.00)
4 1rr Enersol® + ½ inorganic fertilizer	59,521.00 (85.03)	87,800.00 (100.00)	42,831.60 (83.33)	63,384.21 AB (89.45)
5 2rr Enersol® + ½ inorganic fertilizer	81,782.80 (96.67)	59,563.50 (82.27)	61,109.20 (92.31)	67,485.17 A (90.42)
6 Control	17,850.00 (85.00)	27,131.30 (90.74)	24,447.00 (87.00)	23,142.75 C (87.58)

cv 22.90115, p=.0004 *Means with the same letter are not significantly different at $\alpha = 0.05$

Note: Values in parentheses are the percentages of marketable plants per total yield

3.3. Number and Percent Non-Marketable Yield of Pechay

The percent non-marketable yield was computed by determining the non-marketable yield over the total yield and presenting in the table below (Table 4). The highest percent of non-marketable plants were obtained in the plant treated with 1rr Enersol® alone while the lowest non-marketable plants were obtained in the plants treated with 1rr Enersol® and inorganic fertilizer combination. Based on the variability test, the comparability of the treatments is due to the variability among replications which does not consistently provide low or high numbers of non-marketable plants.

The deformities, damaged and unacceptable pechay plants are due to insect damage and non-application of fertilizer specifically in the control and Enersol® alone treated plants. During the trial, the application of insecticides was limited to only three times – during seed sowing and two during the vegetative stage. Thus, no insecticide application was done 10 days prior to harvesting which made them vulnerable to insect attack, particularly by aphids and larvae. Despite the infestation, control measures were not applied since the crop was already in its harvesting stage.

Table 4. Non-marketable yield and percent non-marketable yield of pechay treated with Enersol® and inorganic fertilizer combinations.

TREATMENT	NON-MARKETABLE YIELD (kg/ha, %)			MEAN*
	1	2	3	
1 Inorganic fertilizer alone	4,450.00 (5.00)	0.00 (0.00)	1,812.60 (2.65)	2,087.53 (2.55)
2 1rr Enersol® alone	2,604.00 (6.51)	8,991.28 (26.92)	4,032.75 (14.15)	5,209.34 (15.86)
3 1rr Enersol® + inorganic	0.00	5,700.00	0.00	1,900.00

fertilizer	(0.00)	(6.00)	(0.00)	(2.00)
4 1rr Enersol® + ½ inorganic fertilizer	10,479.00 (14.97)	0.00 (0.00)	8,568.38 (16.67)	6,349.13 (10.55)
5 2rr Enersol® + ½ inorganic fertilizer	2,817.18 (3.33)	12,836.52 (17.73)	5,090.78 (7.69)	6,914.83 (9.58)
6 Control	3,150.00 (15.00)	2,768.74 (9.26)	3,653.00 (13.00)	3,190.58 (12.42)

*Not significant

Note: Values in parentheses are the percentages of marketable plants per total yield

3.4. Plant Height of Pechay

The plant height was gathered (Table 5) during harvesting by measuring the length between the base of the plant and the tallest leaf. The data revealed that the tallest plants were obtained from the plants treated with 1rr Enersol® and inorganic fertilizer combination but were not significantly different from the rest of the treatments except with the plants treated with 1rr Enersol® alone and the control. On the other hand, the shortest plants were obtained from the control but were not significantly different from the plants treated with 1rr Enersol® alone but significantly different from the plants treated with inorganic fertilizer alone or in combination with Enersol® at any rate.

Table 5. Plant height of pechay treated with Enersol® and inorganic fertilizer combinations.

TREATMENT	PLANT HEIGHT (cm)			MEAN
	1	2	3	
1 Inorganic fertilizer alone	27.10	24.30	26.70	26.03 A
2 1rr Enersol® alone	20.40	21.20	19.80	20.47 B
3 1rr Enersol® + inorganic fertilizer	28.10	27.00	25.20	26.77 A
4 1rr Enersol® + ½ inorganic fertilizer	25.90	26.80	25.50	26.07 A
5 2rr Enersol® + ½ inorganic fertilizer	24.50	25.10	24.60	24.73 A
6 Control	19.10	20.70	22.60	20.80 B

cv 1.94217, p<0.0001

*Means with the same letter are not significantly different at $\alpha = 0.05$

3.5. Leaf Length of Pechay

The leaf width was gathered from the biggest leaf of each plant sample (Table 6). The data is remarkably similar to the plant height data in terms of the relationship of treatments among each other. The data revealed that the widest leaves were obtained from the plants treated with 1rr Enersol® and inorganic fertilizer combinations but were not significantly different from the rest of the treatments except with the plants treated with 1rr Enersol® alone and the control.

On the other hand, the narrowest leaves were obtained from the plants treated with 1rr Enersol® alone but were not significantly different with the control but significantly different with inorganic fertilizer alone or in combination with Enersol® at any rate.

Table 6. Leaf length of pechay with Enersol® and inorganic fertilizer combinations at harvest

TREATMENT	LEAF LENGTH (cm)			MEAN
	1	2	3	
1 Inorganic fertilizer alone	17.60	17.20	17.30	17.37 A
2 1rr Enersol [®] alone	13.10	13.50	12.50	13.03 B
3 1rr Enersol [®] + inorganic fertilizer	18.00	17.90	16.20	17.37 A
4 1rr Enersol [®] + ½ inorganic fertilizer	16.30	18.00	15.60	16.63 A
5 2rr Enersol [®] + ½ inorganic fertilizer	16.70	17.00	15.10	16.27 A
6 Control	12.75	13.20	14.00	13.32 B

cv 5.396682, $p < 0.0001$

*Means with the same letter are not significantly different at $\alpha = 0.05$

3.6. Leaf Width of Pechay

The leaf width was gathered from the biggest leaf of each plant sample (Table 7). The data is remarkably similar to the plant height data in terms of the relationship of treatments among each other. The data revealed that the widest leaves were obtained from the plants treated with 1rr Enersol[®] and inorganic fertilizer combination but were not significantly different with the rest of the treatments except the plants treated with 1rr Enersol[®] and control plants.

On the other hand, the narrowest leaves were obtained from the plants treated with Enersol[®] alone but were not significantly different from the control plants and significantly different with inorganic fertilizer alone or in combination with Enersol[®] at any rate.

Table 7. Leaf width of pechay with Enersol[®] and inorganic fertilizer combinations at harvest

TREATMENT	LEAF WIDTH (cm)			MEAN
	1	2	3	
1 Inorganic fertilizer alone	15.70	15.30	15.60	15.53 A
2 1rr Enersol [®] alone	11.30	12.60	10.80	11.57 B
3 1rr Enersol [®] + inorganic fertilizer	16.70	16.10	15.30	16.03 A
4 1rr Enersol [®] + ½ inorganic fertilizer	13.70	16.00	14.50	14.73 A
5 2rr Enersol [®] + ½ inorganic fertilizer	14.90	14.80	15.30	15.00 A
6 Control	10.80	12.10	12.90	11.93 B

cv 5.745721, $p < 0.0001$

*Means with the same letter are not significantly different at $\alpha = 0.05$

3.7. Number of Leaves of Pechay

The number of leaves data was gathered by counting all matured leaves bigger than one inch in size and summarized in the table below (Table 8). The data revealed that no significant difference was found among all the treatments evaluated, although the greatest number of leaves was obtained in the plants treated with inorganic fertilizer alone while the least number was obtained in the control.

Table 8. Number of leaves of pechay with Enersol® and inorganic fertilizer combinations at harvest

TREATMENT	NUMBER OF LEAVES			MEAN*
	1	2	3	
1 Inorganic fertilizer alone	7.70	4.60	4.00	5.43 A
2 1rr Enersol® alone	6.10	3.40	3.40	4.30 A
3 1rr Enersol® + inorganic fertilizer	6.50	5.10	4.60	5.40 A
4 1rr Enersol® + ½ inorganic fertilizer	6.30	4.40	3.70	4.80 A
5 2rr Enersol® + ½ inorganic fertilizer	5.30	4.10	4.20	4.53 A
6 Control	3.20	3.50	3.40	3.37 A

cv 27.29593, p=0.4026 *Not significant

Only a constant supply of nutrients will determine proper plant growth (The efficiency of fertilization is influenced by many factors, such as: soil quality (agronomic category, humus and nutrient content, pH values); the course of weather during the growing season (air temperature and amount of precipitation), and the type of fertilizers used (Magbalot-Fernandez and Agan, 2021). Substances contained in fertilizers improve the absorption of nutrients from the soil, increase resistance to low and high temperatures, and strengthen the plants' ability to defend themselves against pests and diseases (Olaniyi, Ojetayo, 2011). Plants develop properly when grown in fertile and regulated pH soil (Magbalot-Fernandez and Agan, 2021). The acidic nature of the soil limits the growth and development of plants by adversely affecting the soil structure and the biological activity of its microorganisms, and in conditions of low soil pH, humus loses its sticking properties, air-water relations deteriorate, and the activity of soil organisms is reduced (Maziarek, Krawczyk, 2015).

Enersol® Low Dust Granules (LDG), is a mineral consisting mainly of leonardite and humic acid (Enersol LDG, 2015). Both leonardite and humic products are recognized as valuable in improving crop growth. As a fertilizer additive, Leonardite is used as a soil additive to improve a crop's ability to absorb nutrients, resulting in faster growth, healthier plants, and higher yields. Commonly used as a soil amendment, humates add organic matter to marginal soils improving texture, moisture-holding capacity, and pH stability. Enersol® LDG is 63-68% humic acid and 2-4% fulvic acid that is naturally complex with multiple plant essential nutrients at an acid pH for efficient plant uptake. It also has 75-90% organic matter and traces of macro and micronutrients. The product can be applied anytime in the growing season and is excellent for blending and storage with dry granule fertilizers. Enersol® LDG improves plant health and crop yield, by itself or blended with fertilizers. It provides better nutrient uptake and better root zone soil quality. It helps plants against stress and has micronutrients such as calcium, sulfur, nitrogen, iron, magnesium, boron, manganese, zinc copper, molybdenum, and nickel. It easily breaks down in the soil, does not generate dust during handling and in the bag, does not clump urea after blending during storage, and has a high concentration of Gascoyne leonardite.

4. CONCLUSION

The product, Enersol® when in combination with inorganic fertilizer performed well in the trial by giving yield advantages over the control and to the inorganic fertilization alone. The study revealed that the test product was able to enhance pechay production in terms of total fresh yield, marketable yield, plant height, and leaf width with an estimated increase of more than 8%, 8%, 2%, and 3%, respectively relative to inorganic fertilizer application alone. The plants treated with Enersol® have significant improvement in yield and quality of pechay. Furthermore, the bio-efficacy data on total fresh yield, marketable yield, plant height, leaf height, and width showed significant differences with the control. The plants treated with

Enersol® alone and the control plants are inferior in quality in terms of size and color. Thus, Enersol® cannot be applied as a sole fertilizer but only as an amendment to inorganic fertilization to enhance pechay production.

Generally, in this experiment, pechay was very responsive to granular fertilization and in the application of Enersol®. At a very conservative amount of 50 kilograms/hectare, the combination gave a yield advantage of 6,266.67 and 57,866.67 kgs/ha over the full dose of inorganic fertilizer alone and the control, respectively. This could be translated to additional income of PhP 600,973.65 and 5,549,413.65 at PhP 95.90/kg based on the Philippines Retail Price at the National Capital Region for 15 June 2023 over the full dose of inorganic fertilizer alone and the control, respectively (<https://www.ceicdata.com/en/philippines/retail-price-selected-agricultural-commodities/retail-price-pechay-national-capital-region>) The data also revealed that the granular inorganic fertilization dictates the growth and performance of pechay as shown by the inferior performance of the plants treated with half dose of inorganic fertilizer even when supplemented with 2rr Enersol® as compared to the plants with full dose inorganic fertilizer application.

Proper fertilization consists in providing plants with nutrients in the right proportions and amounts, which will enable them to obtain maximum yields of the desired consumption quality, thus protecting the environment and natural resources as well. In the experiment carried out with the use of various combinations and rates of inorganic fertilizers and Enersol LDG, the most favorable effect in terms of its feasibility in pechay production profitability, a full dose of Enersol® in combination with one-half dosage recommended rate of inorganic fertilizer is optimal and highly recommended to obtain the highest possible growth and yield performance of pechay.

REFERENCES

Adorada JL, Adorada AL, Recuenco-Adorada J. Field efficacy of Alka Foliar 5-5-8 fertilizer on the growth and yield of *Brassica rapa* subsp. *chinensis* (Pechay). International Journal of Multidisciplinary Research and Publications, 2023; 5(12):132-137.

Enersol LDG. ENERSOL LDG (Low Dust Granules). Product Information 2015.

Fernandez AM, Agan MS. Bio-Forge promotes growth and yield performance of pechay (*Brassica rapa* L. var. *chinensis* (L.) Hanelt). Annales Universitatis Paedagogicae Cracoviensis, Studia Naturae, 2021; 6: 2021; ISSN2543-8832

Fernandez A, De Guzman C. Physico-chemical quality and sensory evaluation of pummelo fruit as influenced by potassium fertilization. Annals of Tropical Research, 2021; 43(1), 1–20. <https://doi.org/10.32945/atr4319.2021>

Fernandez AM, Andigan AM. Stimulate hormones for higher yield of pechay (*Brassica pekinensis*). Germany, Saarbrücken: Lambert Academic Publishing, 2017.

Fernandez AM, De Guzman CC. Quality and nutrition of Pummelo as influenced by potassium. Journal of Environmental Science and Engineering, 2013; 2(2A), 97–105. <http://doi.org/10.17265/2162-5298/2013.02.004>

Fernandez AM, Miñoza E. Growth and yield of pechay (*Brassica pekinensis*) as affected by green herds organic based foliar fertilizer. Special Issue: First International Conference on

Quality Management of Organic Horticultural Produce 2015. Book of Proceedings, Horticulturae. Basel, Switzerland. P.346.

Fernandez AM, Sabay JL. Growth of tissue-cultured abaca hybrid (*Musa textiles* var. 'seven') plantlets using bioforge supplement. Imperial Journal of Interdisciplinary Research, 2016; 2(8), 249–252.

Gawryś W. Latin-Polish dictionary of herbaceous plants (Słownik roślin zielnych łacińsko-polski). Kraków: Oficyna Botanica, 2008. [In Polish]

Jim M, Tony N. Cabbage growing Primefact 90 replaces AGFACT. H8. 1.27. NSW Department of primary Industries 2006; pp. 1–7.

Jimenez EF, Mariano JS, Ferrer MJ, De Leon V. Pechay production guide. http://bpi.da.gov.ph/bpi/images/Production_guide/pdf/PECHAY%20.pdf (Accessed on June 15, 2021).

Knot JE, Deanon JR. Vegetable production in Southeast Asia. UPCA, 1967.

Kruczek A. Effect of row fertilization with different kinds of fertilizers on the maize yield. Acta Scientiarum Polonorum Agricultura, 2005; 4(2), 37–46. [In Polish]

Magbalot-Fernandez A , Matuguinas JP , Basu SK . Growth performance of tissue - cultured 'Lakatan' Banana (*Musa acuminata*) plantlets using stimulant. International Journal of Agricultural Sciences, 2020; 12(2), 56–58.

Magbalot-Fernandez A, Montifalcon L. Effects of organic-based fortified foliar fertilizer on the growth and yield of 'Lakatan' Banana (*Musa acuminata*). Asian Journal of Research in Crop Science, 2019; 3(4), 1–9. <https://doi.org/10.9734/ajrcs/2019/v3i430053>.

Magbalot-Fernandez A, Agan MS. Bio-Forge promotes growth and yield performance of pechay (*Brassica rapa* L. var. *chinensis* (L.) Hanelt). Annales Universitatis Paedagogicae Cracoviensis Studia Naturae, 2021; 95-108.

Maziarek A, Krawczyk A. Soil as a plant nutrient environment (Gleba jako środowisko odżywczeroślin). Opole: Agenda, 2015. [In Polish]

Montifalcon JR, Fernandez AM. Enhanced growth and yield of lowland rice (*Oryza sativa* L.) with greenshield organic-based fortified foliar fertilizer. Asian Journal of Soil Science and Plant Nutrition, 2017; 1(1), 1–10.

Olaniyi JO, Ojetayo AE. Effect of fertilizer types on the growth and yield of two cabbage varieties. Journal of Animal and Plant Sciences, 2011; 12(2), 1573–1582.

Philippine Statistics Authority. 2017 CAR Crops Production Situationer: Broccoli, Cabbage, Carrots, Habichuelas, Chinese Pechay and White Potato, 2019. <http://rssocar.psa.gov.ph/sites/default/files/CAR-SR-2019-08-2017%20Crops%20Production%20Situationer.pdf> (Accessed on August 25, 2021)