

## Effect Assessment of weed management practices on growth and yield of zero till chickpea (*Cicer arietinum* L.)

### ABSTRACT

A field experiment was conducted to study to compare the "Effect of weed management practices on growth and yield of zero till chickpea (*Cicer arietinum* L.)" at Regional Agricultural Research Station, Polasa, Jagtial during rabi, 2022. The experiment was laid out in randomized block design with seven treatments by replicating thrice. The treatments includes such as application of  $T_1$ -Pendimethalin 38.7 % CS @ 700 g a.i.ha<sup>-1</sup> at 48 hours after sowing,  $T_2$ - Topramezone 33.6 % SC @ 20.6 g a.i. ha<sup>-1</sup> applied at 2-3 leaf stage of weed,  $T_3$ - Quizalofop ethyl 7.5 % + Imazethapyr 15 % EC @ 100 g a.i. ha<sup>-1</sup> applied at 2-3 leaf stage of weed,  $T_4$ - Propaquizafop 2.5 % + Imazethapyr 3.75 % (w/w) ME @ 125 g a.i. ha<sup>-1</sup> applied at 2-3 leaf stage of weed,  $T_5$ - Two hand weeding at 15 and 30 DAS,  $T_6$ - Weed free (Season long) and  $T_7$ - Un-weeded (Check). Obviously, weed free treatment and hand weeded plots showed highest growth and yield attributing characters because of season-long weed-free conditions and better microclimate during critical period of crop. Among different herbicides, Topramezone treated plot recorded higher growth parameters viz.; plant height (38.7 cm), dry matter accumulation (320.8 g m<sup>-2</sup>) and yield attributing characters viz.; total number of pods plant<sup>-1</sup> (30.97), with seed yield (1397 kg ha<sup>-1</sup>), haulm yield (2518 kg ha<sup>-1</sup>) and harvest index (35.7 %). Post emergence herbicide topramezone can be effectively adopted for getting higher yield of chickpea in rice fallow zero tillage conditions.

**Key words:** zero till, chickpea, weed management, herbicides, topramezone, toxicity, growth and yield parameters.

### INTRODUCTION

Pulses are the third-largest crop in India after next to cereals and oilseeds in terms of production, acreage and economic value (Choudhary, 2009). Chickpea (*Cicer arietinum* L.) is a legume of the family Fabaceae. Chickpea ranks as the second-most important pulse crop and the third-most important food legume (Niranjan *et al.*, 2020). Chickpea is valued for its nutritional quality in the vegetarian diet as it is an important source of proteins and minerals (Nair *et al.*, 2014).

In world, Worldwide chickpea is grown on 13.7 million ha of area and produces with a yield of 12.8 million tonnes annually (Afzal, 2021). In India, chickpea is grown on an area of 9.9 million ha with a production of 11.9 million tonnes and productivity is of 1192 kg ha<sup>-1</sup>. In Telangana, chickpea is grown on an area of 1.43 lakh ha with a production of 2.38 lakh tonnes and productivity is 1667 kg ha<sup>-1</sup> (Indiastat, 2021).

According to the ICMR, the daily recommended intake of pulses is 75 g, but only 47.2 g are now accessible. To close this gap, the chickpea crop production must be increased, but due to a variety of biotic and abiotic factors, chickpea productivity is relatively very poor. Due to the vulnerability of the chickpea to weeds, severe weed infestation is one of the significant factors limiting higher output (Nath *et al.*, 2018). Chaudhary *et al.* (2005) reported that weed infestation

**Formatted:** Font: Italic, English (United States)

**Formatted:** Font: Italic, English (United States)

**Formatted:** Font: Italic, English (United States)

**Formatted:** Font: Italic, English (United States)

**Formatted:** Font: (Default) Arial, 10 pt, Not Italic, Font color: Black, English (United States)

**Formatted:** Font: (Default) Arial, 10 pt, Not Italic

can cause up to 75 % yield losses in chickpea due to its shorter plant and poor early vigour. The first 30 days are too crucial for weed crop competitiveness in chickpea (Singh and Singh, 2000).

Recent years, conservation tillage techniques such as minimal or ~~zero-tillage~~ ~~tillage~~ systems have gained relevance. ~~On of the conservation tillage techniques is zero-tillage systems, where soil disturbance is limited to sowing activities. It keeps crop leftovers on the soil surface, which raises the amount of soil organic carbon and enhances soil quality and health.~~ In comparison to conventional tillage, zero tillage systems typically needed less maintenance and provided higher economic returns (Smart and Bradford, 1999) ~~and .This zero tillage reduces cost by 3.8 % to 13.7 % and frees up at least 8 days for the growth of succeeding crops. Zero tillage farming is an option for low-income farmers (Quddus et al., 2020).~~

~~No-tillage chickpea yield was significantly greater (24–57 %) than under reduced, minimum or conventional tillage and chickpea could be replaced by conservation tillage systems that increase yield and will likely improve soil properties in the long term (Hommat and Eskandari, 2004). But, in comparison to conventional systems, weed control is more difficult in conservation tillage systems (Rajkumar et al., 2014). When a conservation tillage system is used, the weed flora alters. According to Bergtold and Sailus (2020), perennial weeds are more likely to cause issues in conservation tillage systems.~~

Chemicals are anticipated to be more commonly used in zero till agriculture ~~as it continues to grow and replace traditional weed control methods where there is adue to~~ labour shortage and rising labour costs ~~make making~~ manual weeding problematic. ~~Pendimethalin~~ Though the pre-emergence herbicide pendimethalin as pre-emergence is the ruling herbicide is effective, however, post-emergence herbicides are limited in this crop particularly for managing broad-leaved weeds. ~~So, our aim is identification of to identify~~ a selective post-emergence herbicide is required to manage broad-leaved weeds in this crop. Hence this study was taken up

~~Therefore, this present investigation was conducted~~ to evaluate ~~the effect of~~ different weed management practices on zero- till chickpea, which can be cost effective and acceptable to the chickpea growers.

## **MATERIALS AND METHODS**

A field experiment was conducted at Regional Agricultural Research Station (RARS), Polasa, Jagtial, Telangana during *rabi*, 2022. The RARS was located at an altitude of 243.4 m above mean sea level (MSL) ~~on 18°84'26" N latitude and 78°95'24" E longitude and it is categorized under Northern Telangana Zone (NTZ).~~ The weekly mean maximum and minimum temperature during the crop growth period ranged from 30.4°C to 34.2°C ~~with an average of 31.34 °C~~ and 13.4 °C to 19.6 °C ~~with an average of 15.92°C~~ respectively. ~~The weekly mean relative humidity in the morning (7.14 AM) and afternoon (2.14 PM) during the crop season~~

Formatted: Space Before: 0 pt

ranged from 80.9 to 93.7 % with an average of 87.84 % and from 23 to 48.6 % with an average of 38.62 % respectively. The weekly mean wind velocity ranged from 0.2 to 41.9 km hr<sup>-1</sup> with an average of 4.95 km hr<sup>-1</sup>. The weekly mean bright sunshine hours day<sup>-1</sup> ranged from 5.1 to 9.1 hr day<sup>-1</sup> with an average of 7.35 hr day<sup>-1</sup>. The weekly mean evaporation (USWB Class A open pan evaporimeter) ranged from 2.2 to 2.9 mm day<sup>-1</sup> with an average of 2.55 mm day<sup>-1</sup>. The total rainfall received during the crop growth period was nil.

The experiment was laid out in randomized block design with seven treatments by replicating thrice. The treatments include T<sub>1</sub>- Pendimethalin 38.7 % CS @ 700 g a.i.ha<sup>-1</sup> at 48 hours after sowing, T<sub>2</sub>- Topramezone 33.6 % SC @ 20.6 g a.i. ha<sup>-1</sup> applied at 2-3 leaf stage of weeds, T<sub>3</sub>- Quizalofop ethyl 7.5 % + Imazethapyr 15 % EC @ 100 g a.i. ha<sup>-1</sup> applied at 2-3 leaf stage of weeds, T<sub>4</sub>- Propaquizafop 2.5 % + Imazethapyr 3.75 % (w/w) ME @ 125 g a.i. ha<sup>-1</sup> applied at 2-3 leaf stage of weeds, T<sub>5</sub>- ~~Two hand weedings~~Hand weeding done twice at 15 and 30 DAS, T<sub>6</sub>- Weed free (Season long) and T<sub>7</sub>- Un-weeded (Check). The soil of experimental field was a sandy clay loams in texture, soil pH (7.31), EC (0.30 dsm<sup>-1</sup>), low in available nitrogen (179.7 kg ha<sup>-1</sup>) and available phosphorus (10.9 kg ha<sup>-1</sup>) but high in available potash (358.4kg ha<sup>-1</sup>). After harvesting of rice crop, in *rabi* season within a week ~~duration~~ time paraquat was sprayed at 5 ml L<sup>-1</sup> to kill the rice stubbles and existing weeds. For ensuring good germination, healthy and good quality NBeG-3 variety chickpea seeds were used with at a seed rate of 65 kg ha<sup>-1</sup> with planting geometry of 30 cm x 10 cm. The recommended dose of fertilizer 20:50:20 kg ha<sup>-1</sup> of NPK was applied at the time of sowing. Three uniform irrigations were given to the crop at pre sowing, pre flowering and pod development stages. The pre-emergence herbicide was sprayed at 48 hours after sowing and post emergence herbicides were sprayed when the weeds are 2-3 leaf stage (18 DAS). The amount of various herbicides needed for the corresponding plots was calculated based on the amount of the active ingredient mentioned in the treatment. Herbicides were sprayed using a knapsack sprayer with a flat fan nozzle to the corresponding plots by mixing the exact amount of herbicides and measured water at a rate of 500 L ha<sup>-1</sup>. The data was analysed by the method of "Analysis of Variance" as described by Panse and Sukhatme (1985).

## RESULTS AND DISCUSSIONS

**GROWTH PARAMETERS:** All the growth parameters were significantly influenced by different weed management practices in zero-till chickpea and are presented in Table 1. The plant height, number of branches plant<sup>-1</sup>, leaf area, and dry matter accumulation were the highest in weed free plots as well as hand weeding plots followed by the topramexone plots.

### Plant height (cm)

Data on plant height from table depicts that weed free treatment produced the tallest plants (43.6 cm) and it was on par with hand weeding treatment (42.0 cm). Among herbicides,

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Comment [GN1]: How was this enabled? Explain.

Formatted: Highlight

Formatted: Highlight

Comment [GN2]: formulation, please

Formatted: Highlight

Comment [GN3]: Post paraquat application, what was the time gap followed before sowing chickpea?

Formatted: Highlight

Comment [GN4]: Pendimethalin? Mention so.

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Comment [GN5]: Were the data subjected to transformation?

Formatted: Font color: Auto

Comment [GN6]: Please mention the major species of weeds observed in unweeded plots to comprehend the efficacy of the treatments better.

treatment topramezonerecorded higher plantheight (38.7 cm) ~~due to maintaining weed free conditions during critical period of crop by applying herbicide that exerted less competition from weeds and produced taller plants.~~ Similar results were found with Quddus *et al.* (2020) and Ratnam *et al.* (2011).

#### **Number of branches plant<sup>-1</sup>**

~~The number of branches plant<sup>-1</sup> varied significantly due to different treatments. Data from table 1 shows that number of branches plant<sup>-1</sup> registered was the highest in weed free plots (30.51) followed by those in hand weeding plots (26.32). Among herbicides, higher number of branches plant<sup>-1</sup> registered in plots applied with topramezone (25.95), due to maintaining weed free conditions during critical period of crop that favours the crop to spread freely.~~ Similar results documented with Sanketh *et al.* (2021).

Formatted: Indent: First line: 0.5"

#### **Leaf area (cm<sup>2</sup>-plant<sup>-1</sup>)**

~~Leaf area was significantly affected by weed management treatments and Table 1 depicts data at 75 DAS, where higher leaf area plant<sup>-1</sup> recorded in weed free treatment (433.99 cm<sup>2</sup>) which was comparable with hand weeding treatment (410.48 cm<sup>2</sup>). Among herbicides, topramezonerecorded higher leaf area plant<sup>-1</sup> (403.75 cm<sup>2</sup>) due to weed free conditions at critical period of crop that results in taller plants coupled with higher number of branches that finally lead to higher leaf area plant<sup>-1</sup>. After 75 DAS leaf area plant<sup>-1</sup> decreases due to leaf senescence.~~ These results are in line with those of Thakur *et al.* (2019).

#### **Dry matter accumulation (g m<sup>-2</sup>)**

~~Dry matter accumulation was significantly influenced by different weed management treatments. An overview of the data from table 1 shows that Again, weed free plot accumulated maximum dry matter (383.5 g m<sup>-2</sup>) which was followed by hand weeded plot (344.2 g m<sup>-2</sup>). Among herbicides, topramezone accumulated maximum dry matter (320.8 g m<sup>-2</sup>). This might be due to improved crop growth parameters viz.; (Plant height, Number of branches plant<sup>-1</sup> and Leaf area plant<sup>-1</sup>). Similar findings were seen with Kumar and Hiremath (2018) and Thakur *et al.* (2019).~~

Formatted: Indent: First line: 0"

**YIELD PARAMETERS:** An overview of the yield attributes data from table 2 revealed that there were significant differences among different weed management treatments except pod weight, number of seeds pod<sup>-1</sup> and test weight.

#### **Total number of branches plant<sup>-1</sup> at harvest**

From the data on yield attributes, weed free treatment depicts significantly higher number of branches plant<sup>-1</sup> (30.51) followed by hand weeding treatment (26.32). Among herbicides, topramezonerecorded higher number of branches (25.95). ~~This could be due to lower weed density at critical period of crop that facilitated crop towards optimum utilization of~~

resources (Sunlight, Water, Space and Nutrients) that reflected in higher number of branches plant<sup>-1</sup>. Similar results were earlier documented by Sanketh *et al.* (2021).

#### Number of pods branch<sup>-1</sup>

Data on yield attributes (Table 2) depicted that weed free produced higher number of pods branch<sup>-1</sup> (2.78) was obtained from weed free plots which was comparable with hand weeding (2.52) and among herbicidal treatments higher number of pods branch<sup>-1</sup> observed in topramezone (2.19). This was due to higher number of branches plant<sup>-1</sup> in turn results in higher number of pods branch<sup>-1</sup>.

#### Total number of pods plant<sup>-1</sup>

An overview on yield attributes, weed free plot registered highest total number of pods plant<sup>-1</sup> (33.93) and it was statistically on par with hand weeding plot (31.20). While, among herbicidal treatments highest total number of pods plant<sup>-1</sup> were registered in topramezone (30.97) due to higher number of branches plant<sup>-1</sup> and number of pods branch<sup>-1</sup>. These results find support from those of Tiwari *et al.* (2019). No significant differences among different weed management practices on pod weight or on number of seeds pod<sup>-1</sup>. This could be attributed to the fact that number of seed seeds weight, seeds pod<sup>-1</sup> and test weight of 100 seeds are a genetic characters and hence not be altered by any weed management practices. The results are in accordance with findings of Sanketh *et al.* (2021) and Ratnam *et al.* (2011).

#### Pod weight (g)

Data on yield attributes had shown that there were no significant differences among different weed management practices on pod weight. Because it was also purely a genetical character.

#### Number of seeds pod<sup>-1</sup>

Yield attributes from table, it was clear that different weed management practices had no significant effect on number of seeds pod<sup>-1</sup>. This could be attributed to the fact that number of seeds pod<sup>-1</sup> is a genetic character and hence not be altered by any weed management practices. The results are in accordance with findings of Sanketh *et al.* (2021).

#### Test weight (100 seeds) (g)

It could be inferred that from the data on yield attributes (Table 2) test weight of different weed management practices was shown as non-significant. This might be due to varietal genetic character. These results corroborate with those of Ratnam *et al.* (2011).

**YIELD:** Seed yield of zero till chickpea was significantly influenced by different weed management practices and it ranged from 903 to 1621 kg ha<sup>-1</sup> across the treatments which were presented in Table 2 and Fig 1.

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Highlight

Formatted: Not Superscript/ Subscript

Formatted: Not Superscript/ Subscript

Formatted: Highlight

**Comment [GN7]:** Insert supporting reference to prove that they can't be improved by cultivation practices such as nutrition or other crop husbandry.

**Comment [GN8]:** If these are established factors, the assessment of herbicides for such characters is redundant.

Formatted: Highlight

Formatted: Highlight

Formatted: Indent: First line: 0"

### Seed yield (kg ha<sup>-1</sup>)

Weed free plot recorded significantly higher seed yield (1621 kg ha<sup>-1</sup>) followed by hand weeding (1453 kg ha<sup>-1</sup>) but among the herbicidal treatment topramezone recorded highest seed yield (1397 kg ha<sup>-1</sup>). It was due to the proper utilization of moisture, nutrients, light and space by the chickpea crop in the absence of weed competition during critical period of crop. While, lowest seed yield was recorded from Un-weeded (Check) plot (903 kg ha<sup>-1</sup>) due to highest crop-weed competition. Similar results were reported by those of Kumari *et al.* (2021), Mishra *et al.* (2012), Quddus *et al.* (2020), Banjara *et al.* (2022) and Sesharee *et al.* (1996).

### Haulm yield (kg ha<sup>-1</sup>)

The haulm yield of zero till chickpea (Table 2) was significantly influenced by different weed management practices.

An overview from the data showed that weed free plot registered higher haulm yield (2648 kg ha<sup>-1</sup>) followed by hand weeding plot (2527 kg ha<sup>-1</sup>). On the other hand, topramezone (2518 kg ha<sup>-1</sup>) treatment among herbicides recorded highest haulm yield. The higher stover yield in above treatments might be due to lesser weeds during early crop growth period, higher yield attributes and pod yield which leads to higher stover yield. Quddus *et al.* (2020) and Banjara *et al.* (2022) found similar results.

### Harvest index (%)


From the table 2 it could be inferred that significantly higher harvest index (38.0 %) observed in weed free treatment, which was statistically comparable with hand weeding treatment but different from remaining treatments. This The minimum harvest index was obtained (33.3 %) under un-weeded (check) due to low seed yield and more crop-weed competition. Maximum harvest index under these treatments might be due to proper reproductive growth due aided by timely translocation of photosynthesis from source to sink thus increase increasing the seed production ratio in total produce. Quddus *et al.* (2020) and Thakur *et al.* (2019) observed similar output.

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold



**Fig. 1 Seed yield, Haulm yield and Harvest index as influenced by different weed management practices in zero till chickpea**

#### CROP PHYTOTOXICITY

Visual scoring for crop phytotoxic symptoms such as yellowing, stunting, chlorosis and necrosis were taken at 3<sup>rd</sup> day after spray of herbicide up to 25 days based on the rating scale. It was noted according to 1-9 scale method (Table 3). For chickpea, 1 meant no to very slight phytotoxicity and 9 for complete destruction of plant (Jiddimaniet al., 2017). A score of < 3 was considered as acceptable (Rao, 2000).

Topramezone treated plot showed yellowing and stunting symptoms (Scale < 2) (Table 43) and recovered completely 10 days after application (DAA) due to the new meristematic growth and initiation of branching. Similar findings on yellowing and stunting in chickpea crop were documented earlier by (Nath et al., 2018) and (Sanketh et al., 2021).

In contrast, treatment T<sub>3</sub> and T<sub>4</sub> showed chlorotic symptoms of phytotoxicity (Score < 4) (Table 43) which was higher at 7 DAA leads leading to necrosis due to death of apical tissue. But, Though by at 30 DAA the crop started producing new branches and associated growth by new meristematic activity in the apical buds, recovered. This the prolonged phytotoxicity affected the crop phenological development as evident from lower seed yield. Similar results of phytotoxicity due to application of imazethapyr leads leading to leaf chlorosis followed by and subsequent necrosis have been reported by earlier workers were in line with (Kumar et al., 2015); (Dubey et al., 2018); (Dubey and Chaudhary, 2021); (Lyon and Wilson, 2005); and (Sanketh et al., 2021).

#### CONCLUSION

**Comment [GN9]:** Repeat of part of Table 2.  
Please remove .

**Formatted:** Highlight

Experimental results revealed that weed free plots as well as manual weeding were the best weed management practices for growth and yield contributing characters in zero-tillage chickpea cultivation. However, in the absence of these, topramezone treated plots recorded higher value for growth and yield contributing characters. Hence, topramezone 33.6 % SC @ 20.6 g a.i. ha<sup>-1</sup> applied at 2-3 leaf stage of weeds is suggested for controlling management of the diversified weed flora under zero tillage conditions in rice-fallow chickpea.

Formatted: Font: Italic

**Table. 1 Growth parameters of zero\_ till chickpea as influenced by weed management practices**

Treatment details	Plant height (cm)	Number of branches plant <sup>-1</sup>	Leaf area (cm <sup>2</sup> plant <sup>-1</sup> )	Dry matter accumulation (g m <sup>-2</sup> )
T <sub>1</sub> -Pendimethalin 38.7 % CS @ 700 g a.i. ha <sup>-1</sup> at 48 hours after sowing	35.5	19.61	372.42	277.9
T <sub>2</sub> -Topramezone 33.6 % SC @ 20.6 g a.i. ha <sup>-1</sup> applied at 2-3 leaf stage of weeds	38.7	25.95	403.75	320.8
T <sub>3</sub> -Quizalofop ethyl 7.5 % + Imazethapyr 15 % EC @ 100 g a.i. ha <sup>-1</sup> applied at 2-3 leaf stage of weeds	34.2	16.66	321.29	244.0
T <sub>4</sub> -Propaquizafop 2.5 % + Imazethapyr 3.75 % (w/w) ME @ 125 g a.i. ha <sup>-1</sup> applied at 2-3 leaf stage of weeds	34.6	17.13	337.64	263.9
T <sub>5</sub> -Two hand weeding at 15 and 30	42.0	26.32	410.59	344.2

DAS				
T <sub>6</sub> -Weed free (Season long)	43.6	30.51	433.99	383.5
T <sub>7</sub> -Un-weeded (Check)	31.0	14.14	285.92	233.2
SEm±	0.6	0.8	4.8	13.9
CD (P=0.05)	1.9	2.3	14.8	42.9

**Table. 2 Yield and yield attributes of zero till chickpea as influenced by weed management practices**

Treat No.	Total number of branches plant <sup>-1</sup> at harvest	Number of pods branch <sup>-1</sup>	Total no. of pods plant <sup>-1</sup>	Pod Weight (g)	No. of seeds pod <sup>-1</sup>	Test weight (g)	Seed yield (kg ha <sup>-1</sup> )	Haulm yield (kg ha <sup>-1</sup> )	Harvest index (%)
T <sub>1</sub>	19.61	1.70	26.80	0.27	1	23.00	1129	2253	33.4
T <sub>2</sub>	25.95	2.19	30.97	0.28	1	23.67	1397	2518	35.7
T <sub>3</sub>	16.66	1.08	20.67	0.27	1	23.00	1000	1922	34.2
T <sub>4</sub>	17.13	1.18	21.43	0.27	1	23.00	1067	2014	34.6
T <sub>5</sub>	26.32	2.52	31.20	0.28	1	23.67	1453	2527	36.5
T <sub>6</sub>	30.51	2.78	33.93	0.29	1	24.00	1621	2648	38.0
T <sub>7</sub>	14.14	1.08	18.67	0.27	1	23.00	903	1802	33.3
SEm±	0.8	0.3	1.3	0.008	-	0.4	31.4	37.9	0.5
CD (P=0.05)	2.3	0.8	3.9	NS	-	NS	96.6	116.9	1.6

**Table. 3 Phytotoxicity score**

Rating	Phytotoxicity %	Description
1	0-10	No to very slight discoloration
2	11-20	More severe but not lasting
3	21-30	Moderate and more lasting
4	31-40	Medium and lasting
5	41-50	Moderately heavy
6	51-60	Heavy
7	61-70	Very heavy
8	71-80	Nearly destroyed

Formatted: Highlight

Formatted Table

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

9	81-100	Destroyed
---	--------	-----------

Source: Jiddimaniet *al.* (2017)

**Comment [GN10]:** Since the reference is given, this table can be removed.

**Formatted:** Highlight

**Table. 4 Phytotoxicity rating (Scale 1-9) as influenced by weed management practices on zero till chickpea**

Treatment details	Dosage (ml ha <sup>-1</sup> )	Phytotoxicity rating
T <sub>1</sub> -Pendimethalin 38.7 % CS @ 700 g a.i. ha <sup>-1</sup> at 48 hours after sowing	1808	-
T <sub>2</sub> -Topramezone 33.6 % SC @ 20.6 g a.i. ha <sup>-1</sup> applied at 2-3 leaf stage of weeds	61.3	1-2
T <sub>3</sub> -Quizalofop ethyl 7.5 % + Imazethapyr 15 % EC @ 100 g a.i. ha <sup>-1</sup> applied at 2-3 leaf stage of weeds	444.4	3-4
T <sub>4</sub> -Propaquizafop 2.5 % + Imazethapyr 3.75 % (w/w) ME @ 125 g a.i. ha <sup>-1</sup> applied at 2-3 leaf stage of weed	2000	3-4
T <sub>5</sub> -Two hand weedings at 15 and 30 DAS	-	-
T <sub>6</sub> -Weed free (Season long)	-	-
T <sub>7</sub> -Un-weeded (Check)	-	-

## REFERENCES

- Afzal, M.K. Economic value of chickpea production consumption and world trade. *AFS-Advance in Food Science*, 2021;211.
- Banjara, G.P., Porte, P and Banjara, B. Impact of different broad-spectrum herbicides on yield and economics of chickpea crop (*Cicer arietinum* L.). *The Pharma Innovation Journal*, 2022;11(11):617-620.
- Bergtold, J and Sailus, M. Conservation tillage systems in the southeast-production, profitability and stewardship. *Sustainable Agriculture Research and Education (SARE) program*, 2020;146.
- Chaudhary, B.M., Patel, J.J and Delvadia, D.R. Effect of weed management practices and seed rates on weeds and yield of chickpea. *Indian Journal of Weed Sciences*, 2005;37:271-272.

Choudhary, A.K. Role of phosphorus in pulses and its management. *Indian Farmers Digest*, 2009;42(9):32-34.

Dubey, A.K., Kumar, A., Singh, D., Partap, T and Chaurasiya, A. Effect of different weed control measures on performance of chickpea under irrigated condition. *International Journal of Current Microbiology and Applied Sciences*, 2018;7(5):3103-3111.

Dubey, T.P and Chaudhary, R.S. Yield and Yield Attributes of Chickpea (*Cicer arietinum* L.) as Influenced by Planting Dates and Weed Management Systems. *Asian Research Journal of Agriculture*, 2021;14(4):123-134.

Hemmat, A and Eskandari, I. Tillage system effects upon productivity of a dryland winter Wheat-Chickpea rotation in the northwest region of Iran. *Soil and tillage research*, 2004;78(1):69-81.

Indiastat. 2020-2021. Agricultural production. <http://www.indiastat.com>.

Jimmimani, L., Chandranath, K.T and Chogatapur, S.V. Phytotoxicity ratings and weed control ratings as influenced by chemical weed control treatments in green gram (*Vigna radiata* L.). *Journal of Pure and Applied Biosciences*, 2017;5(5):1578-1581.

Kaushik, S.S., Rai, A.K., Sirothia, P., Sharma, A.K and Shukla, A.K. Growth, yield and economics of rain fed chickpea (*Cicer arietinum* L.) as influenced by integrated weed management. *Indian Journal of Natural Products and Resources*, 2014;5(3):282-285.

Kumar, M.R and Hiremath, S.M. Bio-efficacy of imazethapyr in rainfed green gram. *Annals of Reviews and Research*, 2018;4(1):555-627.

Kumar, N., Hazra, K.K., Yadav, S.L and Singh, S.S. Weed dynamics and productivity of chickpea (*Cicer arietinum*) under pre and post-emergence application of herbicides. *Indian Journal of Agronomy*, 2015;60(4):570-575.

Kumari, S., Kumar, B., Seema, K.L and Kumar, D. Investigation on efficacy of pre and post emergence herbicides of chickpea (*Cicer arietinum*): Productivity, weed dynamics and economics. *The Pharma Innovation Journal*, 2021;10(9):622-626.

Lyon, D.J and Wilson, R.G. Chemical weed control in dryland and irrigated chickpea. *Weed Technology*, 2005;19(4):959-965.

Mishra, J.S., Singh, V.P., Bhanu, C and Subramahmauam, D. Crop establishment, tillage and weed management techniques on weed dynamics and productivity of Rice (*Oryza sativa*)–Chickpea (*Cicer arietinum* L.) cropping system. *Indian journal of weed science*, 2012;82(1):15-20.

Nair, R., Schafleitner, R., Easdown, W., Ebert, A., Hanson, P., D'arros, H.J and Donough, H. K.J. Legume improvement program at AVRDC-the world vegetable center: impact and future prospects. *Ratarstvo I Povrtarstvo*, 2014;51(1):55–61.

**Comment [GN11]:** The content is removed from the manuscript and thus this reference can be removed.

**Formatted:** Highlight

**Formatted:** Highlight

**Comment [GN12]:** Missing in tet

**Formatted:** Highlight

**Formatted:** Highlight

**Comment [GN13]:** Missing in text

**Formatted:** Highlight

**Formatted:** Highlight

Nath, C.P., Dubey, R.P., Sharma, A.R., Hazra, K.K., Kumar, N and Singh, S.S. Evaluation of new generation post-emergence herbicides in chickpea (*Cicer arietinum* L.). *National Academy Science Letters*, 2018;41(1):1-5.

Niranjan, I.K., Tyagi, S., Kumar, B and Pradhan, A.K. Evaluation of different post-emergence herbicides in chickpea (*Cicer arietinum* L.). *International Journal of Agricultural Applied Sciences*, 2020;1(1):87-91.

Nyanga, P.H. Food security, conservation agriculture and pulses: Evidence from small holder farmers in Zambia. *Journal of Food Research*, 2012;1:120.

**Comment [GN14]:** Not found in text.

**Formatted:** Highlight

**Formatted:** Highlight

Panase, V. G and Sukhatme, P. V. Statistical methods for Agricultural workers. (3rd Edition). ICAR, New Delhi, 1985.

Quddus, M.A., Naser, H.M., Siddiky, M.A., Ali, M.R., Mondol, A.A.I., Islam, M.A and Gazipur, B. Impact of Zero Tillage and Tillage Practice in Chickpea Production. *Journal of Agricultural Science*, 2020;12(4):106-118.

Rajkumara, S., Gundlur, S.S., Neelakanth, J.K and Ashoka, P. Impact of irrigation and crop residue management on Maize (*Zea mays*) - Chickpea (*Cicer arietinum* L.) sequence under no tillage conditions. *Indian Journal of Agricultural Sciences*, 2014;84(1):43-48.

**Comment [GN15]:** Missing in text.

**Formatted:** Highlight

**Formatted:** Highlight

Rao, V.S. Weed research methodology-Field experimentation. *Principles of Weed Science*, 2000;497-498.

Ratnam, M., Rao, A.S and Reddy, T.Y. Integrated weed management in chickpea (*Cicer arietinum* L.). *Indian Journal of Weed Science*, 2011;43(1&2):70-72.

Sanketh, G.D., Rekha, K.B., Sudhanshu, K.S and Ramprakash, T. Effect of integrated weed management with new herbicide mixtures on growth, yield and weed dynamics in chick pea. *The Pharma Innovation Journal*, 2021;10(7):1074-1077.

Sesharee, A.P., Prasad, V.N., Rao, K.L and Prasad, H. Integrated weed management in gram (*Cicer arietinum* L.). *Indian Journal of Agronomy*, 1996;41(3):496-497.

Singh, S and Singh, A.N. Crop-weed competition in chickpea: Challenges and strategies for the New Millenium. In *National Symposium on Agronomy*, 2000;15(18):199.

Smart, J.R and Bradford, J.M. Conservation tillage corn production for a semi-arid, subtropical environment. *Agronomy Journal*, 1999;91:116-21.

Thakur, A.K., Thakur, N.S., Kushwaha, B.B and Bangar, K.S. Effect of Different Weed Control Methods on Growth and Yield of Chickpea (*Cicer arietinum* L.) under Irrigated Condition in Malwa Region of Madhya Pradesh. MSc Thesis, Rajmata VijayarajeScindia Krishi Vishwa Vidyalaya, Gwalior, Indore, Madhya Pradesh, 2019.

Tiwari, R., Vivek, Naresh, R.K., Tyagi, S., Shivangi and Kumar, A. Evaluation of effective weed management strategy for enhancing productivity and profitability of chickpea under sub-

tropical climate of western U.P. *International Journal of Chemical Studies*, 2019;7(5):928-933.

Yadav, V.L., Shukla, U.N., Raiger, P.R and Mandiwal, M. Efficacy of pre and post-emergence herbicides on weed control in chickpea (*Cicer arietinum* L.). *Indian Journal of Agricultural Research*, 2019;53(1):112-115.

**Comment [GN16]:** Not found in text.

**Formatted:** Highlight

**Formatted:** Highlight

UNDER PEER REVIEW