

Original Research Article

Impact of Different Levels of NPK and Neem Cake on Soil Health under Cowpea (*Vigna unguiculata* L.) Var. Gomati.

ABSTRACT

A field experiment was conducted during March-June (Late *Zaid*) 2022 at the research farm of Department Soil Science and Agricultural Chemistry, NAI, SHUATS, laid out in Randomized Block Design. To achieve higher growth and yield of rainy cowpea variety *Gomati*, it was observed that application of @ 100% NPK+ @ 100% Neem Cake has shown effective growth under Prayagraj climatic conditions but also improved significantly. physical and chemical properties of soil. This physico-chemical like Bulk density, Partical density, Pore space (%), Water Holding Capacity (%), pH, Electrical conductivity (dS m^{-1}), Organic Carbon (%), Available Nitrogen (kg ha^{-1}), Available Phosphorous (kg ha^{-1}), Available Potassium (kg ha^{-1}) were analysis. Result revealed that Soil Organic carbon content, Water holding capacity and available NPK significantly increased in after harvest of cowpea in T₉, whereas Bulk density (Mg m^{-3}), particle density (Mg m^{-3}) and pH recorded minimum value. Therefore treatment T₉ was best in terms of growth, yield and economic parameters with maximum plant height, number of branches plant⁻¹, pods plant⁻¹, seeds pod⁻¹, pod yield and maximum cost benefit ratio.

Keywords: Physico-Chemical properties, Nitrogen, Phosphorus, Potassium, Neem Cake and Cowpea.

1. INTRODUCTION

Cowpea belongs to the family Leguminosae and sub family Fabaceae. It is a warm season, annual and herbaceous legume. Plant types are often categorized as erect, prostrate or creeper. There is much variability within the species. Growth habit ranges from intermediate to profuse. Flowers are born in multiple racemes on 8-20 inches long flower stalk that arise from the long leaf axils. Cowpea seeds contain 54.4% carbohydrates, 24.2% protein and 0.1% fat. Moreover, it is a rich source of phosphorus, calcium and iron. The crop produces such heavy vegetative growth and completely covers the ground that it prevents soil erosion in problematic areas and can later be ploughed under to produce green manure. This might be the result of the plant receiving more nutrients over time, as well as more leaf fall and field waste. It has considerable promise as an alternative pulse crop (Madukwe *et al.*, 2008).

Pulses are least preferred by farmers because of high risk and less remunerative than cereals consequently, the production of the pulses is sufficiently low. Among the pulse crops, cowpea is more cosmopolite and grown in most of the regions of India which showed very encouraging results and promises to have a far-reaching significant in achieving a breakthrough in the pulse production. Cowpea is cultivated throughout India for its long green pods as vegetables, seeds and pulses, as foliage, as green manure, as well as green fodder. The cultivars grown for their immature pods are variously known as “Asparagus bean”, “Snake bean” and “Yard long bean”. Cowpea originated in India and Ethiopia and it is widely grown in India as well as all over the world. (Sailaja kumari, and Usha kumari, 2002).

Nitrogen is important for plant nutrient and a crucial component of protein, but also found in many other compounds that have a significant physiological impact on plant metabolism. Crop production approaches involve the growth of nitrogen-fixing leguminous crops, such as cowpea (*Vigna unguiculata* L.), which is able to fix up to 337 kg N ha⁻¹. In addition, cowpea can provide vital micronutrients, enhancing nutritional yield. When growing cowpea under subsistence farming system, some rural communities add, for example, inorganic fertilizer in addition to inter-cropping in order to optimize use of limited productive agricultural land. Inter-cropping by rural resource-poor farmers dates back to ancient civilization and is today still being practiced. (Buhlebelive *et al.*, 2020)

Phosphorus is an essential component of numerous plant processes, including energy metabolism, nitrogen fixation, membrane and nucleic acid synthesis, photosynthesis, respiration, and enzyme regulation. Phosphorus is critical to cowpea yield because it stimulates growth, initiates nodule formation as well as influences the efficiency of the *Rhizobium* legume symbiosis. Phosphorus decreases zinc concentration in the cowpea grain, thereby affecting its nutritional quality. It is required for the physiological processes of protein synthesis and energy transfer in plants. Application of phosphorus has been reported by several authors to improve yield of cowpea. Seed yield is, therefore, governed by number of factors which have a direct or indirect impact. Among these factors some are yield components such as number of pods plant⁻¹, number of seeds pod⁻¹ and 100-seed weight over a given land area (Ndakidemi and Dakora, 2007).

Potassium plays an important role in formation of protein and chlorophyll and it provides much of osmotic "pull" that draws water into plant roots. Potassium helps reduce lodging in maize and produces strong, stiff straw. Potassium gives plants more strength and resistance to disease. (Cobbinah *et al.*, 2011).

Neem cake known as non-edible oilcake organic manure. The composition of neem cake is 5.2% N, 1.0% P, 1.4% K. It also acts as a nematicide but also as a nitrogen inhibitor reducing nitrification. It supplies the available nitrogen for a long time in the soil. (Katyayan, 2012). The neem cake obtained from the expeller contains 8-10% oil, which is recovered by solvent extraction. The cake's residual oil and limonoids repel insects, and solvents or water extracts make the product a strong antifeedant and growth inhibitor. The seed cake is rich in plant nutrients (crude protein 13-18%, carbohydrate 24-50%, crude fiber 8-26%, fat 2-13%, ash 5-18% and acid insoluble ash 1-17%, with nitrogen, phosphorus, calcium, and magnesium) and is used as manure for soil amendment and for urea coating. (Manoj Kumar, 2018).

2. MATERIALS AND METHODS

The experiment was conducted at Research Farm of Soil Science and Agricultural Chemistry at Sam Higginbottom University of Agriculture Technology and Sciences which is situated six km away from Prayagraj city on the right bank of Yamuna river, the experimental site is located in the sub-tropical region with 25°58'N latitude, 81°52'E longitude and at an altitude of 98 m above mean sea level. The area of Prayagraj district comes under subtropical belt in the South east of Uttar Pradesh, which experiences extremely hot summer and fairly cold winter. The maximum

temperature of the location reaches up to 46⁰ C-48⁰ C and seldom falls as low as 4⁰ C – 5⁰ C. The relative humidity ranges between 20 to 94 percent. The average rainfall in this area is around 1100 mm annually. The soils of experimental area fall in order of Inceptisol and soil was alluvial. The soil samples were randomly collected from three different sites in the experiment plot prior to tillage operation from a depth of 0-15 cm and 15-30 cm. The size of the soil sample was reduced by conning and quartering the composites soil sample and was air dried passed through a 2 mm sieve for preparing the sample for physical and chemical analysis.

Table. 1 Treatment Combinations of Cowpea

Treatments	Treatment description
T ₁	Absolute Control
T ₂	@50% NPK +@0% Neem Cake
T ₃	@100% NPK +@0% Neem Cake
T ₄	@0% NPK +@50% Neem Cake
T ₅	@50% NPK +@50% Neem Cake
T ₆	@100% NPK +@50% Neem Cake
T ₇	@0% NPK +@100% Neem Cake
T ₈	@50% NPK +@100% Neem Cake
T ₉	@100% NPK +@100% Neem Cake

Table. 2 Results of Physic-chemical properties of pre-soil sample. (Before sowing of crop)

Particulars	Results		Methods
	(0-15) cm	(15-30) cm	
Sand (%)	61.1	59.5	Bouyoucos hydrometer (Bouyoucos 1927)
Silt (%)	24.4	25.3	
Clay (%)	14.5	15.2	
Textural Class	Sandy loam	Sandy loam	
Bulk density	1.322	1.329	Graduated measuring cylinder (Muthuaval <i>et al.</i> 1992)
Particle density	2.184	2.201	
Pore space (%)	45.69	44.77	
Water holding capacity (%)	41.75	39.95	
Soil pH (1:2.5)	7.601	7.609	Digital pH meter (Jackson 1958)
Electrical Conductivity (dS m ⁻¹)	0.175	0.196	Digital conductivity meter (Wilcox 1950)
Organic Carbon (%)	0.253	0.247	Wet oxidation method (Walkley and Black 1947)
Available Nitrogen (kg ha ⁻¹)	167.10	148.64	Modified alkaline permanganate oxidation method (Subbiah and Asija 1956)
Available Phosphorus (kg ha ⁻¹)	23.61	20.37	Spectrophotometric method (Olsen <i>et al.</i> 1954)
Available Potassium (kg ha ⁻¹)	194.51	186.79	Flame photometric method (Toth and Prince 1949)

3. RESULTS AND DISCUSSION

3.1 Physical analysis

The maximum Bulk density of soil was 1.25 Mg m^{-3} and 1.29 Mg m^{-3} at depth 0-15 cm and 15-30 cm, respectively in T_1 (Absolute control) and minimum bulk density was recorded in 1.12 Mg m^{-3} and 1.13 Mg m^{-3} at depth 0-15 cm and 15-30 cm, respectively in T_9 (@ 100% NPK + @100% Neem Cake).

The maximum Particle density of soil was recorded 2.26 Mg m^{-3} and 2.28 Mg m^{-3} at depth 0-15 cm and 15-30 cm in T_1 (Absolute control) and minimum Particle density recorded in 2.13 Mg m^{-3} and 2.17 Mg m^{-3} at depth 0-15 cm and 15-30 cm, respectively in T_9 (@ 100% NPK + @100% Neem Cake).

Treatment T_9 (@ 100% NPK + @100% Neem Cake) recorded maximum Pore space 57.32%, 55.42% at depth 0-15 cm and 15-30 cm. and minimum pore space was recorded in 46.23%, 45.27% at depth 0-15 cm and 15-30 cm, respectively in T_1 (Absolute Control).

The response of Water holding capacity of soil was found to be significant level of NPK and Neem Cake. The maximum water Holding capacity was recorded in T_9 (@ 100% NPK + @ 100% Neem Cake) 52.31%, 39.37% at depth 0-15 cm and 15-30 cm and minimum Water holding capacity was recorded in 41.29%, 32.29% at depth 0-15 cm and 15-30 cm, respectively in T_1 (Absolute Control).

3.2 Chemical analysis

The maximum pH of soil was recorded 7.53, 7.65 at depth 0-15 cm and 15-30 cm in T_1 (Absolute control) and minimum pH of soil was recorded in 7.15,7.25 at depth 0-15 cm and 15-30 cm, respectively in T_9 (@ 100% NPK + @100% Neem Cake).

The maximum EC of soil was recorded $0.26, 0.28 \text{ dS m}^{-1}$ at depth 0-15 cm and 15-30 cm in T_9 (@ 100% NPK + @100% Neem Cake) and minimum EC of soil was recorded in $0.15, 0.21 \text{ dS m}^{-1}$ at depth 0-15cm and 15-30 cm, respectively in T_1 (Absolute Control).

The maximum Organic carbon of soil was recorded in 0.65%, 0.57% at depth 0-15c m and 15-30 cm in T_9 (@ 100% NPK + @100% Neem Cake) and minimum organic carbon was recorded in 0.42, 0.36 at depth 0-15 cm and 15-30 cm, respectively in T_1 (Absolute Control).

The maximum Available Nitrogen of soil was recorded in 281.39, 273.24 kg ha^{-1} at depth 0-15 cm and 15-30 cm in T_9 (@ 100% NPK + @100% Neem Cake) and minimum Available Nitrogen of soil was recorded in 256.71, 251.12 at depth 0-15 cm and 15-30 cm, respectively in T_1 (Absolute Control).

The maximum Available Phosphorous of soil was recorded in 27.89, 25.96 kg ha^{-1} at depth 0-15 cm and 15-30 cm in T_9 (@ 100% NPK + @100% Neem Cake) and minimum Available of Phosphorous of soil was recorded in 20.26, 18.36 kg ha^{-1} , at depth 0-15 cm and 15-30 cm, respectively in T_1 (Absolute Control).

The maximum Available Potassium of soil was recorded in 158.15, 156.32 kg ha⁻¹ at depth 0-15 cm and 15-30 cm in T₉ (@ 100% NPK + @100% Neem Cake) and minimum Available Potassium of soil was recorded in 124.16, 120.65 kg ha⁻¹ at depth 0-15 cm and 15-30 cm, respectively in T₁ (Absolute Control).

UNDER PEER REVIEW

Table. 3 Physical properties of soil sample after harvesting of Cowpea.

Symbols	Treatments	Bulk Density (Mg m ⁻³)		Particle Density (Mg m ⁻³)		Pore space (%)		Water holding capacity (%)	
		Depth (cm)		Depth (cm)		Depth (cm)		Depth (cm)	
		0-15	15-30	0-15	15-30	0-15	15-30	0-15	15-30
T₁	Control	1.25	1.29	2.26	2.28	46.23	45.27	41.29	32.29
T₂	@50% NPK +@0% Neem Cake	1.21	1.26	2.21	2.25	48.56	46.35	43.67	33.35
T₃	@100% NPK +@0% Neem Cake	1.19	1.21	2.19	2.22	52.24	51.29	48.56	31.16
T₄	@0% NPK +@50% Neem Cake	1.14	1.17	2.17	2.23	51.78	50.78	44.36	30.56
T₅	@50% NPK +@50% Neem Cake	1.18	1.19	2.20	2.21	52.39	51.58	50.30	38.45
T₆	@100% NPK +@50% Neem Cake	1.16	1.17	2.15	2.18	54.68	53.65	48.63	36.35
T₇	@0% NPK +@100% Neem Cake	1.17	1.18	2.12	2.20	53.67	52.32	46.68	34.26
T₈	@50% NPK +@100% Neem Cake	1.13	1.15	2.16	2.19	55.49	54.26	49.56	37.63
T₉	@100% NPK +@100% Neem Cake	1.12	1.13	2.13	2.17	57.32	55.42	52.31	39.37
	F-test	S	S	S	S	S	S	S	S
	S.Em±	0.018	0.017	0.026	0.021	0.664	0.836	0.564	0.737
	C.D.@5%	0.056	0.072	0.080	0.064	1.999	2.517	1.700	2.219

Fig.1 Physical properties of soil sample after harvesting of Cowpea.





Table. 4 Chemical properties of soil sample after harvesting of Cowpea.

Symbols	Treatments	Soil pH		Electrical Conductivity (dS m ⁻¹)		Organic Carbon (%)		Available Nitrogen (kg ha ⁻¹)		Available Phosphorus (kg ha ⁻¹)		Available Potassium (kg ha ⁻¹)	
		Depth (cm)		Depth (cm)		Depth (cm)		Depth (cm)		Depth (cm)		Depth (cm)	
		0-15	15-30	0-15	15-30	0-15	15-30	0-15	15-30	0-15	15-30	0-15	15-30
T₁	Control	7.53	7.65	0.15	0.21	0.42	0.36	256.71	251.12	20.26	18.36	124.16	120.65
T₂	@50% NPK +@0% Neem Cake	7.48	7.46	0.19	0.23	0.51	0.45	263.65	256.36	24.18	21.45	137.26	134.46
T₃	@100% NPK +@0% Neem Cake	7.37	7.39	0.21	0.25	0.56	0.51	274.84	263.48	26.38	23.17	141.18	139.84
T₄	@0% NPK +@50% Neem Cake	7.42	7.45	0.17	0.19	0.48	0.39	259.74	249.26	22.19	19.26	125.12	123.58
T₅	@50% NPK +@50% Neem Cake	7.31	7.32	0.23	0.27	0.54	0.43	267.65	261.75	23.16	21.37	138.32	135.20
T₆	@100% NPK +@50% Neem Cake	7.41	7.40	0.22	0.24	0.59	0.52	278.51	267.86	25.78	22.18	145.37	142.76
T₇	@0% NPK +@100% Neem Cake	7.27	7.31	0.18	0.22	0.50	0.46	263.79	257.41	21.37	23.01	137.76	135.69
T₈	@50% NPK +@100% Neem Cake	7.25	7.29	0.24	0.26	0.62	0.54	271.86	254.47	26.78	18.68	149.10	147.64
T₉	@100% NPK +@100% Neem Cake	7.15	7.25	0.26	0.28	0.65	0.57	281.39	273.24	27.89	25.98	158.15	156.32
	F-test	S	S	S	S	S	S	S	S	S	S	S	S
	S.Em±	0.062	0.074	0.002	0.003	0.007	0.006	3.250	3.507	0.408	0.365	2.201	2.345
	C.D.@5%	0.187	0.223	0.008	0.009	0.022	0.018	9.786	10.559	1.228	1.099	6.625	7.060

Fig.2 Chemical properties of soil sample after harvesting of Cowpea.



4. CONCLUSIONS

It was concluded that the treatment T₉ was the best in terms of all soil parameters like Bulk density (Mg m⁻³), Particle density (Mg m⁻³), Pore Space (%), Soil pH, Electrical Conductivity (dS m⁻¹), Organic Carbon (%), Available Nitrogen, Phosphorus and Potassium (kg ha⁻¹), as well as helps to increase yield of cowpea, thus overall benefit to Indian farmers.

ACKNOWLEDGEMENTS

The authors are thankful to Advisor, Department of Soil Science and Agriculture chemistry, Sam Higginbottom University of Agriculture, Technology and Sciences Prayagraj, U. P. for taking their keep interest and encouragement to carry the research work.

REFERENCES

- Buhlebelive, Mndzebele., ORCID.Bhekumthetho. Ncube., Melvin, Nyathi., Sheku, Alfred.Kanu. and Melake, F. (2020) Nitrogen Fixation and Nutritional Yield of Cowpea-Amaranth Intercrop. *Agronomy* 2020 **10**(4):565.
- Cobbinah, FA., Addo-Quaye, A.A. and Asante, I.K. (2011) Characterization, evaluation and selection of cowpea accessions with desirable traits from eight regions of Ghana. *ARPJ Agric. Biol. Sci.* 2011; **6**:21-32.
- Joshi D, Gediya KM, Patel jS, Birai MM. Effect of organic manures on growth and yield of summer cowpea (*vigna unguiculata* L.). *Journal of Agricultural science digest.* 2016; 36(2):134-137.
- Khandelwal R. Effect of inorganic and biofertilizers on productivity and nutrients uptake in cowpea [*Vigna unguiculata* (L.) Walp]. *Legumes Research.* 2012 35(3): 235-238.
- Kumar Yadav, Ramawat Naleeni and Singh Dashrath. Effect of organic manures and biofertilizers on growth and yield parameters of cowpea (*Vigna unguiculata* (L.) Walp.). *Journal of Pharmacognosy and Phytochemistry* 2019; 8(2): 271-274.
- Madukwe, D. K., Christo, I. E. C. and Onuh, M. O. (2008). Effect of organic manure and cowpea walp varieties on the chemical properties of the soil and root nodulation. *Sel World.*3 (1):210-214.
- Manoj Kumar KP, Tarence Thomas, Amreen Hasan P, Smriti Rao. Effects of vermicompost and inorganic fertilizers on physico-chemical properties of soil in Indian mustard. *Journal of Pharmacognosy and Phytochemistry* 2018; **7**(3): 1999-2001.

- Ndakidemi, PA. and Dakora, FD. (2007) Yield components of nodulated cowpea (*Vigna unguiculata*) and maize plants grown with exogenous phosphorus in different cropping systems. *Aust. J. Exp. Agric.* 47:583-589.
- Subbiah, BV, and Asija, GL. A rapid procedure for the estimate of available Nitro gen in Soil
Current Sci. 1956;25: 259-260.
- Sailaja kumari, M.S. and Usha kumari, K. (2002). Effect of vermicompost enriched with rock phosphate on yield and uptake of nutrients in cowpea. *Journal of Tropical Agriculture*, **40**: 27-30.
- Taniya Mistri, Narendra Swaroop, AA David and A Barthwal. Assessment of organic and inorganic fertilizers on physico-chemical properties of soil of tomato. *The Pharma Innovation Journal*. 2021; 10(8): 703-707.
- Toth, SJ, and Prince AL. Estimate of cation exchange capacity and exchangeable Ca, K, Na, content of soil by flame photometer technique soil sci. 1949; 67(6): 439-445.
- Olsen, SR, Cole CV, Watanabe, F.S. and Dean, LA. (1954) Estimate of available Phosphorus In soils by extraction with sodium bicarbonate. United States Department of Agriculture circular.1954; 939:1-19.
- Jackson ML. Soil chemical analysis Second edition Indian reprint, prentice hall of India, New Delhi.1973; 498.
- Sailaja kumari, M.S. and Ushakumari, K. (2002). Effect of vermicompost enriched with rock phosphate on yield and uptake of nutrients in cowpea. *Journal of Tropical Agriculture*, **40**: 27-30.
- Wilcox LV. Electrical Conductivity Journal American Water works Association 1950; 42: 775-776.
- Walkley, A. and Black, IA. Critical examination of rapid method for determining organic carbon in
soils, effect of variance in digestion conditions and inorganic soil constituents. *Soil science*.1947; 632:251.