

Original Research Article

FEATURE OF IRON ACCUMULATION IN THE GRAIN OF SOFT WHEAT BELONGING TO DIFFERENT ECOLOGICAL AND GEOGRAPHICAL REGIONS

Abstract:

Among the grain-growing countries in the world, countries such as Canada, the United States of America, Mexico, Brazil, Australia, China, India, Turkey, and Russia, in the direction of selection for the creation of new varieties of wheat resistant to abiotic factors, are paying great attention to creating new wheat varieties by obtaining new genotypes as a result of identifying donors with high quality indicators and with positive indicators of valuable economic traits and introducing them into modern selection methods. In this direction, some progress has been made in the world, today many varieties of wheat with valuable economic traits and with high grain quality have been created and are being introduced to large areas.

During our research, 23 genotypes were selected from 45 genotypes of soft wheat varieties and lines. It was observed that the growth period of this nursery lasted 233-238 days. It was also noted that the lines were more mature than the model varieties. Compared to the model varieties, among the biometric indicators of the plant: 15 lines manifested a positive result in terms of plant height indicators, 10 lines by last node length indicators, 5 lines by spike length indicators, 1 line by spike number indicator, and 1 line showed a positive result on resistance for lodging. The results of the statistical analysis of grain yield and grain quality by the Dospekhov method showed that the experimental error on the yield index is 0.888%, on the weight of 1000 grains is 3.018%, on the nature of the grain index is 0.627%, in terms of the protein content index is 2.028%, and gluten content index is 1.519%, in terms of the IDK indicator is 2.001%, and in terms of grain glassiness index the error is 4.01%, and it was noted that the experiment was conducted correctly in terms of repetitions and showed a positive result.

Key words: soft wheat, varieties and lines, 1000 grain weight, protein content indicator, iron content in the grain.

Introduction: Wheat is considered one of the most important grain crops in world agriculture, occupying 17% of the total agricultural land, and about 750 million tons of grain is produced annually. Globally, a total of 240.8 million hectares are planted with wheat, and it is predicted that the demand for wheat grain will increase even more in the coming years.

Triticum aestivum L. is a biennial crop grown in spring and fall, and it is a spike-producing crop with a flat leaves and small flowers. Wheat stem is a common straw with five to seven joints and nodes, three to four leaves, the length of the leaf is 20-37 cm, and the width is 1-2 cm [1; 13].

According to Sh. Dilmurodov [2], effective use of newly created intensive type varieties is one of the main factors in increasing soft wheat grain yield, which is competitive, suitable for regional soil and climate conditions.

During the period of shooting of wheat, productive stalks, spikes inside the stalks and grains in the spike are formed. The duration of the shooting phase in wheat is 25–30 days. During this period, the plant accumulates 50–60% of the dry matter that it accumulates during the entire vegetation period [3; 14].

Winter wheat is divided into classes according to the height as follows:

dwarf (lower from 60 cm), semi-dwarf (60-85 cm), lower height (85-105 cm), medium height (105-120 cm) and tall (higher than 120 cm) [4].

D.T. Juraev [5] stated that the height of the stalk of winter soft wheat changes depending on the weather conditions and is grouped as: dwarf-stemmed (50-75 cm), short-stemmed (76-90 cm), medium-stemmed (91- 110 cm), and long-stemmed (higher than 110 cm).

The grain yield can be reduced by 30-50% due to the lodging of the wheat plants, and the photosynthetic activity of the wheat leaves is reduced as a result of lodging, it becomes difficult to harvest the grain with machines [6].

According to O.A. Amanov [7], the length of the spike and the number of spikelets in the spike are mainly related to the characteristics of the variety, which is determined by some differences. Productive tillering, the number of grains per spike and the weight of 1000 grains are of great importance in the high productivity, although they interact with the external environment.

The growing importance of wheat in the population requires increasing its gross yield and quality. In solving the problem, it is important to adapt new varieties of wheat to different soil-climate zones and agrometeorological conditions. The dependence of the variety and weather conditions on the accumulation of protein and crude gluten was studied in early-ripening and medium-ripening soft wheat varieties grown in the gray soils of Kemerovo region. As a result, it was determined that the protein content in early-ripening varieties was from 8.6 to 13.8%, and in the medium-ripening was from 9.0 to 14.0%. As the scientists emphasized, the effect of the genotype is not always the same, it changes a lot depending on the weather [8; 9].

The relationship between the height of the stalk and the lodging does not always appear. Mostly lodging is caused depending on biological property of the variety and, mainly, on the anatomorphological structure. Below

Comment [PG1]: Grain is a broader term, use cereal or wheat instead

Comment [PG2]: Neither the meaning of the line is clear nor is it grammatically correct

Comment [PG3]: Use genotypes instead of line as it is a more scientific word

Comment [PG4]: Elaborate about the nursery. It is not clear which nursery and in which context it is used?

Comment [PG5]: Positive result in which context. Explain? The abstract is incomplete without it

Comment [PG6]: ??? Meaning not clear...syntax error

nodes of the varieties susceptible to lodging are longer, the plant is tall, the stalk and nodule is small or thin, while the resistant varieties to lodging are short in height and below nodes are short as well [10].

The formation of grains, a high weight of 1000 grains is a decisive factor for obtaining abundant and stable grain yield. Lack of moisture in the soil, high temperature, infestation by fungal diseases lead to a decrease in the weight of 1000 grains [11].

Under the influence of drought, the decrease in the yield was caused by stunted growth point of the plant, and a decrease of the assimilation process on the surface of the plant leaves. Dry weather has a negative effect on grain glassiness, grain quality, and different nitrogen combinations in the grain [12].

As A. Amanov [19] stated, the amount and quality of gluten is a defining indicator for assessing the technological and nutritional quality of wheat grain and is determined by the IDK (index of gluten deformation) tool. If IDK indicator is 0-15, then the gluten is very unsatisfactory belonging to group III, if IDK is 20-40 then gluten is satisfactory in the group II, when IDK is 45-75 the gluten is good belonging to group I, when 80-100 it is satisfactory, belonging to group II, and if IDK is 105-120 then gluten is unsatisfactory, belonging to group III [5].

D. Juraev, O. Amanov [20] in their scientific research, stated that the grain weight in the spike is considered one of the important indicators for the high yield, and the grain weight in the spike of the varieties and samples changes over the years depending on the weather conditions [6].

Materials and methods:

Field experiments were conducted in the experimental plot of the Karshi district branch of the Southern Agricultural Research Institute. Experiments in field conditions were carried out in the field experiment area of "Genetics and selection of grain crops" department. Experiments in laboratory conditions were carried out in the "Laboratory of Plant Biochemistry and Evaluation of Quality Indicators" and "Laboratory of Organo-Mineral Fertilizers and Agrochemical Gross Analysis" of the institute.

The experimental layout was done based on Complete block design and Alpha lattice design of GenStat 13 software. Phenological observations, calculations and analyzes were carried out according to the method of the All-Union Plant Science Institute (1984) and biometric analyzes were carried out according to the methods of the Center for Agricultural Crops Variety Trial (1985, 1989).

The technological quality indicators of the grain of winter soft wheat crops grown in the experimental field were determined according to the methodological manuals "Guidelines for assessing grain quality", "Methods of biochemical research of plants".

Statistical analyzes was done based on the method of B.A. Dospekhov (1985).

RESULTS AND DISCUSSION

The research was carried out in Ya. Omonov farm in Karshi district and the soil under the experiment was light gray soil. It constituted 19.2% of the total land area of Kashkadarya region. The irrigated area is 24.6%.

Table 1. The duration of the growing season of soft wheat varieties and ridges belonging to different ecological-geographical regions (Karshi 2021-2022).

№	Name of Genotypes	Germination date	Tillering date	Shooting date	Heading date	Days to heading	Days to maturity	Vegetation period
1	Gozgon (check)	12.10.21	15.11.21	07.02.22	10.04.22	180	03.06.22	234
2	KR20-27-FAWIR-67	12.10.21	15.11.21	08.01.22	02.04.22	172	03.06.22	234
3	Antonina (check)	12.10.21	15.11.21	14.02.22	07.04.22	178	02.06.22	233
4	KR20-BWF5IR-3301	11.10.21	18.11.21	16.02.22	04.04.22	175	03.06.22	235
5	KR20-27-FAWIR-73	12.10.21	08.11.21	10.02.22	05.04.22	175	03.06.22	234
6	KR20-BWF5IR-2083	12.10.21	19.11.21	02.02.22	05.04.22	176	03.06.22	235
7	KR20-BWF5IR-2460	11.10.21	16.11.21	02.02.22	04.04.22	176	05.06.22	237
8	KR20-BWF5IR-3380	12.10.21	15.11.21	09.02.22	02.04.22	172	05.06.22	236
9	KR20-BWF5IR-2095	11.10.21	18.11.21	14.02.22	05.04.22	176	01.06.22	233
10	KR20-BWF5IR-2625	11.10.21	18.11.21	16.02.22	06.04.22	177	04.06.22	236
11	KR20-BWF5IR-3529	13.10.21	16.11.21	16.02.22	05.04.22	174	02.06.22	233
12	KR20-27-FAWIR-84	12.10.21	19.11.21	13.01.22	05.04.22	175	07.06.22	238
13	KR20-BWF5IR-94	12.10.21	19.11.21	08.01.22	07.04.22	177	06.06.22	237
14	KR20-BWF5IR-2113	11.10.21	16.11.21	07.02.22	05.04.22	176	03.06.22	236
15	KR20-27-FAWIR-27	11.10.21	18.11.21	01.02.22	05.04.22	176	04.06.22	236
16	KR20-BWF5IR-144	11.10.21	16.11.21	15.02.22	03.04.22	174	04.06.22	236
17	KR20-27-FAWIR-138	12.10.21	19.11.21	12.01.22	03.04.22	173	04.06.22	235
18	KR20-27-FAWIR-39	11.10.21	17.11.21	14.02.22	03.04.22	174	01.06.22	233
19	KR20-27-FAWIR-142	12.10.21	17.11.21	11.01.22	02.04.22	172	02.06.22	233
20	KR20-27-FAWIR-46	12.10.21	16.11.21	01.02.22	03.04.22	174	01.06.22	233
21	KR20-27-FAWIR-154	12.10.21	17.11.21	12.01.22	08.04.22	178	03.06.22	234
22	KR20-BWF5IR-246	13.10.21	19.11.21	07.02.22	03.04.22	173	03.06.22	234

Comment [PG7]: Spelling mistakes and grammatical errors need to be improved

Comment [PG8]: Introduction is incomplete as nothing about iron content and its importance is mentioned here which is the main centre line of the manuscript

Comment [PG9]: Explain about the longitude, latitude and soil type of the area

Comment [PG10]: Indicate from which geographical regions the genotypes are procured along with their pedigree

23	KR20-BWF5IR-3235	12.10.21	15.11.21	10.02.22	03.04.22	173	02.06.22	233
Minimum		11.10.21	08.11.21	08.01.22	02.04.22	172	01.06.22	233
Mean		11.10.21	16.11.21	01.02.22	04.04.22	175	03.06.22	234
Maximum		13.10.21	19.11.21	16.02.22	10.04.22	180	07.06.22	238

Source: [Compiled by the authors]

The control variety trial nursery has 45 varieties and lines, they planted in 10m² area in 2 replications. It was observed that the germination of varieties and samples lasted until October 11-13. Standard variety Gozgon and Antonina variety germinated on October 12. The other 8 samples were observed to germinate on October 11. During the tillering period, the plant stops growing, and as a result of the cooling of the temperature, side stalks grow from the main stalk. It was observed that the tillering phase lasted from November 8 to November 19. Standard varieties Gozgon and Antonina manifested tillering on November 15. Compared to standard varieties, KR20-27-FAWIR-73 line tillered on November 8, and was found as earlier tillering variety (Table 1). During the shooting phase, the plant growth point continues to grow in soft wheat varieties and samples. In this process, the spike is formed. The shooting phase of varieties and samples lasted from January 8 to February 16. It was noted that the spike formation period lasted from April 2-10.

DHD




Fig.1 Differentiation of germination-heading period between replications and genotypes

Source: [Compiled by the authors]

The day from germination to spike formation lasted 172-180 days. It was noted that the standard variety Gozgon formed spikes in 180 days, Antonina variety in 178 days. Compared to the standard variety KR20-27-FAWIR-67, KR20-BWF5IR-3380 lines formed spikes in 172 days. Full ripening lasted until June 1-7. It was observed that the days from germination to maturity of the varieties and samples lasted for 233-238 days. It was found that the lines ripened 2-3 days earlier in comparison to the standard variety.

Table 2 Analysis of variance for various traits in winter wheat genotypes evaluated in 2022

Source of variation	df	Mean square			
		Days to heading	Days to maturity	Plant hieght	Grain yield
Replication	1	113,34	0,278	1448,01	1,188
Genotype	44	12,7	4,577	99,03*	6,12**
Error	44	12,78	3,164	58,94	6,961
Total	89	138,82	8,019	1506,95	8,149
CV (%)		2	0,8		0,4

ns Noon significant

*Significant at P=0,05

**Significant at P=0,01

Source: [Compiled by the authors]

DMD




Fig.2 Differentiation of germination-maturing period between replications and genotypes

Source: [Compiled by the authors]

It was observed that the plant height index of soft wheat varieties and lines ranged from 91.5 to 123.0 cm. The indicator of plant height in standard Gozgon variety was 104.0 cm and in Antonina variety 98.5 cm. It was noted that the KR20-BWF5IR-2460 line was 120.0 cm, and the KR20-27-FAWIR-46 line was 123.0 cm higher than the standard variety. The length of the last node was 35.0-51.0 cm in varieties and lines, and compared to standard Gozgon variety, 5 lines and compared to standard Antonina variety, 10 lines showed higher results. Spike length serves as part of the crop element in the plant. It was found that the length of the spike in varieties and lines was 10.5-13.0 cm. It was observed that the number of spikelets ranged from 19-23 pieces, in the standard Gozgon variety, this index was 21 spikelets, and in the Antonina variety 22 spikelets. Compared to the standard variety, KR20-BWF5IR-3529 line showed a higher index of 23 spikelets with a difference of 1-2 pieces (Table 3). When the resistance to lodging in soft wheat varieties and lines was evaluated as a percentage, it was observed that it was up to 3-40% in the varieties and lines.

Table 3 Biometric indicators of soft wheat varieties and lines belonging to different ecological and geographical regions (against 2021-2022).

Sl. No.	Name of Genotypes	Plant height, cm	Peduncle length, cm	Spike length, cm	Number of spikelets of per spike	Lodging resistance, %
1	Gozgon (check)	104.0	41.5	12.0	21	10
2	KR20-27-FAWIR-67	108.5	37.5	11.0	22	5
3	Antonina (check)	98.5	35.0	10.5	22	25
4	KR20-BWF5IR-3301	108.0	45.5	12.0	22	18
5	KR20-27-FAWIR-73	108.5	38.0	11.0	20	25
6	KR20-BWF5IR-2083	115.0	40.0	10.5	20	13
7	KR20-BWF5IR-2460	120.0	51.0	13.0	21	15
8	KR20-BWF5IR-3380	114.5	37.5	10.5	22	25
9	KR20-BWF5IR-2095	110.5	42.5	11.5	20	25
10	KR20-BWF5IR-2625	96.5	38.0	10.5	20	3
11	KR20-BWF5IR-3529	103.0	40.0	12.5	23	8
12	KR20-27-FAWIR-84	111.5	37.0	10.5	19	5
13	KR20-BWF5IR-94	108.5	41.5	11.0	21	5
14	KR20-BWF5IR-2113	112.5	43.5	12.5	21	20
15	KR20-27-FAWIR-27	101.5	41.0	11.0	20	3
16	KR20-BWF5IR-144	106.0	40.5	12.0	21	8
17	KR20-27-FAWIR-138	96.5	42.0	10.5	20	10
18	KR20-27-FAWIR-39	112.0	43.0	11.5	21	10
19	KR20-27-FAWIR-142	91.5	35.0	11.0	21	15
20	KR20-27-FAWIR-46	123.0	42.0	10.5	21	15
21	KR20-27-FAWIR-154	110.0	38.5	11.0	21	40
22	KR20-BWF5IR-246	100.0	36.5	11.5	22	15
23	KR20-BWF5IR-3235	106.5	41.5	10.5	21	15
	Minimum	91.5	35.0	10.5	19	3
	Mean	107.2	40.4	11.2	21	14
	Maximum	123.0	51.0	13.0	23	40

Source: [Compiled by the authors]

The standard Gozgon variety has 10% lodging resistance, and Antonina variety has 25%. It was observed that the KR20-27-FAWIR-154 line has 40 percent resistance, higher than the standard varieties.

PH_{cm}




Fig.3 Differentiation of plant height between replications and genotypes

Source: [Compiled by the authors]

Under the influence of drought, a decrease in productivity was caused by stopping of growth points, reduction of assimilation processes on the plant leaf surface. Dry weather had a negative effect on grain glassiness, grain quality, and various nitrogenous compounds in grain. When determining the yield index in varieties and lines, it showed 59.7-96.7 c/ha. The standard Gozgon variety showed 80.5 c/ha, and Antonina variety 76.9 c/ha. Compared to the standard varieties, it was observed that the KR20-27-FAWIR-67 line showed a high rate of 90.9 c/ha, and the KR20-BWF5IR-2113 line showed a high rate of 96.7 c/ha. When the results of the statistical analysis by the Dospekhov method on the productivity indicator were carried out, it was observed that the experiment error was small, 0.888% and showed a positive result (Table 5).

Comment [PG11]: The results need to be elaborated more




Fig.4 Differentiation of grain yield and TKW between replications and genotypes

Source: [Compiled by the authors]

Table 4 Analysis of variance for various traits in winter wheat genotypes evaluated in 2022

Source of variation	df	Mean square			
		1000 kernel weight	Test weight	Protein	Gluten
Replication	1	3,873	23,511	0,093	0,004
Genotype	44	22,64**	1665,6**	2,648**	5,554**
Error	44	0,331	6,329	0,03	0,047
Total	89	4,204	29,84	0,123	0,051
CV (%)		1,5	0,3	1	

**Significant at P=0,01

Source: [Compiled by the authors]

A high weight of 1000 grain in grain formation is a decisive sign of abundant and stable harvest. Lack of moisture in the soil, high temperature, infestation with fungal diseases lead to a decrease in the weight of 1000 grains. It was found that the weight of 1000 grains was 33.8-43.2 g in varieties and lines. Compared to the standard Gozgon variety, 14 lines, and compared to the Antonina variety, 3 lines showed higher results.

Grain nature is the mass of grain per liter. Grain nature is determined according to GOST 3040-55. It was observed that grain nature was up to 755.0-835.0 g/l in varieties and lines. KR20-BWF5IR-2083 line had 835.0 gr/l grain nature, KR20-27-FAWIR-84 line had 821.0 gr/l, KR20-BWF5IR-2113 line had 828.5 gr/l, KR20-27-FAWIR-39 line had was 834.0 g/l grain nature, and all showed good results compared to standard varieties. The experimental error was 0.627 % and it is proved that it showed a positive result.

Table 5 Grain quality indicators of soft wheat varieties and ridges belonging to different ecological and geographical regions (Karshi 2021-2022).

№	Name of Genotypes	Grain yield, c/ha	1000 kernel weight, g	Test weight, g/l	Protein content, %	Gluten content, %	IDK	Vitreosity, %
1	Gozgon (check)	80.5	33.8	783.5	17.3	28.1	98.0	73.0
2	KR20-27-FAWIR-67	90.9	36.0	783.0	19.3	26.6	105.3	61.0
3	Antonina (check)	76.9	41.7	818.5	17.7	31.2	96.0	76.3
4	KR20-BWF5IR-3301	76.2	36.7	783.0	15.6	26.4	95.4	53.8
5	KR20-27-FAWIR-73	59.7	41.0	801.5	17.3	30.1	110.7	86.3
6	KR20-BWF5IR-2083	73.0	36.8	803.0	18.4	29.2	104.8	80.5

7	KR20-BWF5IR-2460	73.7	40.8	835.0	18.5	28.7	106.0	80.8
8	KR20-BWF5IR-3380	76.3	35.4	765.5	16.3	27.4	103.5	62.3
9	KR20-BWF5IR-2095	75.2	35.2	755.0	16.1	27.0	98.0	59.8
10	KR20-BWF5IR-2625	63.6	37.1	775.0	16.6	26.7	94.4	56.3
11	KR20-BWF5IR-3529	73.0	34.2	759.5	16.8	25.8	93.1	58.8
12	KR20-27-FAWIR-84	73.3	41.1	821.0	15.5	28.5	86.6	82.3
13	KR20-BWF5IR-94	66.2	43.0	811.5	15.5	29.2	92.1	84.8
14	KR20-BWF5IR-2113	96.7	43.2	828.5	15.4	30.9	104.8	72.8
15	KR20-27-FAWIR-27	62.2	39.7	809.0	16.2	28.6	95.9	70.3
16	KR20-BWF5IR-144	72.6	37.9	766.5	15.7	26.3	104.9	81.8
17	KR20-27-FAWIR-138	73.6	42.2	821.5	16.2	27.5	97.0	70.8
18	KR20-27-FAWIR-39	79.4	41.0	834.0	17.7	28.7	102.3	77.3
19	KR20-27-FAWIR-142	79.4	42.8	822.0	18.1	30.4	105.8	82.0
20	KR20-27-FAWIR-46	63.3	34.2	762.5	15.2	26.4	92.4	53.3
21	KR20-27-FAWIR-154	75.2	41.3	824.0	18.0	30.3	97.6	76.8
22	KR20-BWF5IR-246	69.0	42.1	820.5	18.6	29.7	98.0	85.3
23	KR20-BWF5IR-3235	67.8	36.8	822.5	18.4	28.7	118.8	86.8
	Minimum	59.7	33.8	755.0	15.2	25.8	86.6	53.3
	Mean	73.8	38.9	800.3	16.9	28.3	100.0	72.7
	Maximum	96.7	43.2	835.0	19.3	31.2	118.8	86.8
	LSD0,05	0.52	1.14	4.98	0.35	0.43	2	2.92
	LSD0,05%	0.888	3.018	0.627	2.028	1.519	2.001	4.01
	CV%	0.4	1.5	0.3	1	0.8	1	2

Source: [Compiled by the authors]

According to the data, as the wheat fields move from north to south, from west to east, the amount of protein in the grain increases. The amount of protein depends on the amount of nitrogen content and moisture in the soil. For this reason, the amount of protein in grain depends on 30% heredity and 70% on agrotechnical measures. If the plant is supplied with enough nutrients, especially nitrogen, the protein will accumulate more in the grain. The reason for this is that protein is made up of amino acids, and amino acids have an amino group in their name, and the amino group holds nitrogen in its radical. Nitrogen increases protein, excess moisture causes it to decrease.

When determining the indicator of protein content in the grain of the high-yielding, high-grain quality of the studied varieties and lines of the control variety trial nursery on the range of project, it was 15.2-19.3%. Compared to the standard variety, 7 lines with higher protein content were selected among the lines.




Fig.5 Differentiation of grain yield and TKW between replications and genotypes

Source: [Compiled by the authors]

The baking properties of wheat flour are mainly evaluated by the amount and quality of gluten. The amount and quality of gluten refers to the hydrated gel-rubbery mass, which consists mainly of water-insoluble protein when wheat dough is washed in water. It was noted that the gluten content of soft wheat varieties and lines was 25.8-31.2%. This index was 28.1% in standard variety Gozgon, and 31.2% in Antonina variety. It was found that 12 lines showed a higher index compared to the standard Gozgon variety. While the standard Antonina variety showed a higher index compared to the lines. It was observed that the experiment error was 1.519% in varieties and lines and showed a positive result. One of the necessary indicators characterizing the technological properties of the grain depends on the amount of gluten in the wheat grain and mainly on the IDK index of the gluten in the bread making process. The gluten quality of soft wheat varieties and lines the IDK unit indicator according to the state standard is: 1st class (excellent) up to 45-75; 2nd class (good) up to 80-100; 3rd class (unsatisfied) up to 105-120.

It was found that the IDK index of soft wheat varieties and lines in the control variety trial nursery ranged from 86.6 to 118.8. Standard varieties Gozgon and Antonina were accepted to the 2nd class (good) with 96.0 to 98.0 indicators. 11 lines were classified as 2nd class (good), 5 lines as 3rd class (unsatisfactory). The glassiness or hardness of the grain is one of the characteristics of the wheat variety. Nevertheless, these signs can change according to the growing conditions of the wheat plant. The glassiness quality of the grain decreases in conditions of excess moisture and lack of nitrogen.

The glassiness of grain is determined according to GOST 10987-76. According to our research, the glassiness of soft wheat varieties and lines was determined and showed was 73.0-76.3 indicators in the standard varieties Gozgon and Antonina. It was observed that 9 lines with 80.5-86.8 indicators showed higher results compared to standard varieties. The experimental error in grain glassiness according to the Dospekhov method was 4.01% and showed a positive result.

Table 6 Correlation of characteristics of genotypes

Correlations	DHD	DMD	GY_c_ha	Gluten_%	IDK	PH_cm	PL_cm	Protein_%	SL_cm	TKW_g	TST_g_l
DMD	0,12	-									
GY_c_ha	-0,13	0,08	-								
Gluten_%	0,08	-0,04	-0,01	-							
IDK	-0,01	-0,10	0,03	-0,09	-						
PH_cm	0,12	0,01	0,03	0,04	-0,02	-					
PL_cm	-0,04	0,00	0,18	-0,01	0,12	0,48					
Protein_%	0,01	-0,23	-0,04	0,38	0,24	0,08	0,06	-			
SL_cm	0,05	-0,03	0,09	0,06	-0,04	0,00	0,17	0,09	-		
TKW_g	-0,07	0,15	0,30	0,44	-0,05	0,04	0,17	0,09	-0,12	-	
TST_g_l	-0,04	-0,07	0,20	0,44	0,09	0,12	0,17	0,30	-0,07	0,66	-
Vitreosity_%	0,05	0,09	0,03	0,62	0,12	0,04	-0,05	0,41	0,01	0,33	0,50

Source: [Compiled by the authors]

Currently, due to the widespread use of new methods and, first of all, molecular genetics, significant progress has been made in understanding these mechanisms. It should be noted that a number of aspects of plant resistance to various metals and their absorption and adaptation by plants are still not sufficiently studied and require additional research. In recent times, the level of use of reserve proteins with a high degree of polymorphism as a marker of useful agricultural traits of cereal crops is increasing.

As a result of the determination of proteins in wheat plants using the electrophoresis method, it was found that gliadin and glutelin are storage proteins with a complex structure, composed of many structural components. It is noted that these proteins can represent the unique genetic system of wheat in a relatively detailed state. Therefore, the study of these structural components can serve as markers of wheat grain quality and specific genes.

Isolation of gliadin proteins and their electrophoretic study. Electrophoretic analysis of gliadin from storage proteins in wheat was carried out on polyacrylamide gel (PAAG) in an acidic medium by the method of V.A. Bushuk and R.R. Zilman. The electrophoretic spectrum of varieties with high iron content was used as a standard. The electrophoretic formulas of gliadin proteins were divided into four (α , β , γ and ω) fractions according to method of V.G. Konarev.

In the studies, depending on the low or high iron content of grain and flour, reserve proteins in grain were analyzed according to the movement of proteins in polyacrylamide gel towards the poles and the bonds they left behind. For the analysis, one hundred grains of the varieties were threshed separately. After becoming flour form, each of them was mixed with 200 μ l of 70% ethyl alcohol in a separate test tube and extracted for 30-40 minutes in a thermostat at a temperature of 40°C.




Fig.6 Varieties in the exhibition nursery. 1. Yaksart, 2. Gozgon, 3. Antonina, 4. Bunyodkor, 5. Shams, 6. Krasnodarskaya-99, 7. Hisorak, 8. Turkistan, 9. Bezostaya-100, 10. Brigada, 11. .Navbahor, 12.Davr, 13.Grom, 14.Chillaki, 15.Mars, 16.Dostlik, 17.Kroshka, 18.Polovchanka, 19.Starshina, 20.Tanya.

Source: [Compiled by the authors]

According to the results of the analysis, it was known that there was 1.4 mg of Fe in the grain of Yaksart variety, and 1.1 mg of Fe in the flour of this variety, and its α -, β -, γ - and ω - bonds can be compared with other varieties. It is known that the grain of Bunyodkor variety under number 4 contains 1.6 mg Fe, and the flour of this variety contains 1.2 mg Fe. Although the amount of iron in Bunyodkor grain is somewhat higher, the amount of iron in the flour is 1.2 mg, and the iron in the grain is lost through the husk (bran). The grain of Bezostaya-100 variety studied in PAAG gel contains 1.6 mg Fe, its flour 1.2 mg. The variety Davr under number contains 121.4 mg Fe in the grain, 1.2 mg in flour. The variety Chillaki under number 14, contains 1.6 mg of iron, these varieties contain more iron than other varieties, and their β -zone protein bands are similar (Figure 6).

It is known that the electrophoretic spectra of gliadin proteins soluble in 70% ethanol spirit of wheat grain in 3.5% PAAG gel are genetically determined, these spectra are specific for each variety and do not change regardless of the conditions under which wheat varieties are grown. In several subsequent electrophoretic analyses, there is information that some varieties consist of certain biotypes according to their electrophoretic spectra, and based on this information, the varieties with high Fe content were selected and analyzed.

In polyploid cereal species, prolamins have been found to be controlled by several independent (unlinked) gene clusters, and analysis of a small number of grains (around 100 grains) from a single cultivar population allows you to think with confidence about the degree of genetic homogeneity within the cultivar population. Nevertheless, genetic heterogeneity at prolamin gene loci was reported to be up to 17% in prolamin protein selection studies of wheat and oat varieties. In order to use several valuable economic traits of the studied wheat varieties for the purpose of selection and breeding and to study their genetic nature, their genotypes were identified.

The electrophoretic spectrum of gliadin proteins is divided into 4 zones, including those designated by α -, β -, γ - and ω -zones. It is noted that there are a number of bands in each zone, on the basis of which intra- and inter-variety differences are determined.




Figure 7. Varieties in the nursery of demonstrative varieties. 1. Alekseyvich, 2. Nodr, 3. Asr, 4. Babur, 5. Qadr, 6. Yog'du, 7. Pervitsa, 8. Drujba, 9. Velena, 10. Zvezda, 11. Uzb. Mus.-25, 12. Aziz, 13. Durdona, 14. Gurt, 15. Omad, 16. Yuka, 17. Andijan-2, 18. Ishanch, 19. Andijan-4, 20. Aleksiy-2
Source: [Compiled by the authors]

Figure 7 shows the 20 varieties in the exhibition nursery of the studied varieties. According to the results of the analysis, Alekseyvich variety is listed in number 1 with bands. This variety contains 1 mg of iron in the grain and flour, which indicates that the main part of iron is located in the endosperm. It was found that the analyzed grains of the variety Gurt under number 14 and Omad variety under number 15 contain 1.6 and 1.8 mg of iron, respectively, and their electrophoretic spectra on PAAG gel were genetically determined, and these spectra were specific for each variety.

When analyzing the 20 varieties whose electrophoretic spectra are presented in Figure 5, it was found that Matonat variety under number 1 contains 1.4 mg of iron and 1.1 mg of iron in its flour, and their electrophoretic spectra on PAAG gel are genetically different. According to the results of the analysis, it was found that there is 1.6 and 1.5 mg of iron in Yuksalish variety grain and its flour respectively, and electrophoretic spectra in PAAG gel are genetically determined (Fig. 7).

In our experiments, the number of bands in the electrophoretic spectrum of gliadin proteins in the variety samples of ancient local soft wheat varieties in Uzbekistan were analyzed by dividing them into minor, moderately active or major groups. According to the results obtained when the electrophoretic spectra of the varieties with high iron content in grain and flour were analyzed by morphological characteristics and separated as individual variety samples, compared to the electrophoretic spectrum of the control, ancient local variety samples were divided into homogeneous or heterogeneous varieties according to the electrophoretic content of gliadin proteins.




Figure 8. Varieties in the nursery of demonstrative varieties. 1. Matonat, 2. Vekha, 3. Yangi hayot, 4. Sardar, 5. Shijoat, 6. Ziyokor, 7. Rokhat, 8. Turon, 9. Yuksalish, 10. Sarbon, 11. Shukrona, 12. Kesh-2016, 13. Dovan, 14. Istikbol, 15. Navroz, 16. Flight, 17. J. Gavhari, 18. Gallakor, 19. Ravon, 20. Aksaroy

Source: [Compiled by the authors]

In soft wheat (*Triticum aestivum*), gliadin proteins have been found to be controlled by several independent (unlinked) clusters, and analysis of 100 grains from a single cultivar population provides a reliable estimate of the degree of genetic homogeneity or heterogeneity within that cultivar population.

In addition to several valuable economic traits of the studied wheat varieties, it is necessary to identify their genotype in order to use them as varieties with high iron content and for breeding purposes, and to study their genetic nature (Figure 8).

Therefore, it is important to analyze the electrophoretic content of gliadin proteins in samples of ancient local soft wheat varieties.

The flour obtained from threshing grain and grain sample was burned and the amount of Fe in the resulting ash was determined using an Atomic Adsorption Spectrometer AAS 200.

The 20 varieties in the nursery of locally created soft wheat lines, and according to the results of the analysis, the presence of iron was detected and their electrophoretic spectra on the PAAG gel were genetically determined, and it was found that these spectra are specific for each line.

Comment [PG12]: Explain about the safe levels of range of iron content and clarify which varieties are found in this research with safe levels of iron content that can be used in breeding program

Conclusions

According to the conducted research, it was found that soft wheat varieties and lines were early-ripening. Varieties and lines with high grain yield and grain quality indicators were selected, and during the research, the amount of iron contained in grain and flour was determined in laboratory conditions and concluded as follows: the grain of Bezostaya-100 in PAAG gel contains 1.6 mg iron, its flour contains 1.2 mg, Davr variety grain has 1.4 mg of iron, flour 1.2 mg, and the variety Chillaki grain contained 1.6 mg iron. These varieties have more iron than other varieties, and it was found that their protein bands in the β - zone are similar. The grain of Gurt and Omad varieties under analysis contained 1.6 and 1.8 mg of iron, respectively, and their electrophoretic spectra on PAAG gel were genetically determined, and these spectra were specific for each variety. The variety Yuksalish grain contains 1.6 mg iron and its flour 1.5 mg, and their electrophoretic spectra on PAAG gel were genetically different. It was recommended to use these varieties in selection and breeding works as a variety with a high iron content.

Reference:

1. Beebe S. Biofortification of common bean for higher iron concentration //Frontiers in Sustainable Food Systems. – 2020. – T. 4. – P. 573449.
2. Dilmurodovich DS, Bekmurodovich BN, Shakirjonovich KN, Shomiljonovich SS & Raxmatullaevich AJ. (2021). Productivity, quality and technological characteristics of bread wheat (*Triticum aestivum* L.) variety and lines for the southern regions of the Republic of Uzbekistan. Plant cell biotechnology and molecular biology, 63-74. Retrieved from <https://www.ikppress.org/index.php/PCBMB/article/view/5935>
3. Diyor Juraev T, Amanov OA, Dilmurodov SD, Meyliev AK, Boysunov NB, Kayumov N.S & Ergashev ZB. (2021). Heritability of Valuable Economic Traits in the Hybrid Generations of Bread Wheat. Annals of the Romanian Society for Cell Biology, 2008-2019. Retrieved from <https://www.annalsofscb.ro/index.php/journal/article/view/2730>
4. Gupta O. P. et al. Identifying transcripts associated with efficient transport and accumulation of Fe and Zn in hexaploid wheat (*T. aestivum* L.) //Journal of biotechnology. – 2020. – V. 316. – P. 46-55.
5. Juraev DT, Amanov OA, Dilmurodov SD, Boysunov NB & Odirovich JF. (2020). To study the heat resistance features of bread wheat varieties and species for the southern regions of the republic of Uzbekistan. European Journal of Molecular & Clinical Medicine, 7(2), 2254-2270.
6. Juraev DT, Amanov OA, Dilmurodov SD, Boysunov NB, Kayumov NS, Ishankulova GN & Togaeva KR. (2021). The influence of hot-dry wind on farm valuable traits of wheat genotypes in southern regions of Uzbekistan. Plant cell biotechnology and molecular biology, 22(35-36), 34-49. Retrieved from <https://www.ikppress.org/index.php/PCBMB/article/view/6300>
7. Khazratkulova S, Sharma RC, Amanov A, Ziyadullaev Z, Amanov O, Alikulov S & Muzafarova D. (2015). Genotype \times environment interaction and stability of grain yield and selected quality traits in winter wheat in Central Asia. *Turkish Journal of Agriculture and Forestry*, 39(6), 920-929. Doi:10.3906/tar-1501-24
8. Rehman A. et al. Agronomic biofortification of zinc in Pakistan: Status, benefits, and constraints //Frontiers in sustainable food systems. – 2020. – V. 4. – P. 591722.
9. Safdar L. B. et al. Genome- wide association study identifies five new cadmium uptake loci in wheat //The Plant Genome. – 2020. – T. 13. – №. 2. – P. e20030.
10. Shang J. et al. Comparative studies on physicochemical properties of total, A- and B-type starch from soft and hard wheat varieties //International journal of biological macromolecules. – 2020. – V. 154. – P. 714-723.
11. Sharma D. et al. Biofortification of wheat: Genetic and agronomic approaches and strategies to combat Iron and Zinc deficiency //International Journal of Environment, Agriculture and Biotechnology. – 2020. – V. 5. – №. 4.
12. Velu G., Singh R. P., Joshi A. K. A decade of progress on genetic enhancement of grain zinc and iron in CIMMYT wheat germplasm //Wheat and barley grain biofortification. – Woodhead Publishing, 2020. – P. 129-138.
13. Wang Y. et al. Cytogenetic analysis and molecular marker development for a new wheat–*Thinopyrum ponticum* 1Js (1D) disomic substitution line with resistance to stripe rust and powdery mildew //Frontiers in Plant Science. – 2020. – V. 11. – P. 1282.
14. Wani S. H. et al. Improving zinc and iron biofortification in wheat through genomics approaches //Molecular Biology Reports. – 2022. – V. 49. – №. 8. – P. 8007-8023.

Comment [PG13]: The conclusion needs to be more clear, write about future prospects and conflict of interests