

Adult onset Still's disease revealed by a cardiac tamponade: Case report

Abstract:

Adult-onset Still's disease (AOSD) is a rare systemic inflammatory disease of unknown origin with various clinical manifestations. In this article, we describe a case of 57-year-old presenting with Still's disease complicated by cardiac tamponade. This case highlights the importance of a prompt diagnosis of cardiac involvement in inflammatory diseases, as it can be fatal, and underlines the utility of echocardiographic evaluation not only in symptomatic patients, but also for the systematic detection of pericardial effusions. The case of hemodynamic compromise, it allows the provider to determine the timing, approach, and method of pericardial drainage due to cardiac tamponade being a very rare complication that requires an invasive approach.

Keywords: Diagnosis; treatment; cardiac tamponade; Adult Onset Still Disease : AOSD.

Introduction:

Still's disease was first described by Still(1) in 1897 in children, but it was not until the 1970s that Bywaters(2) and Bujak et al(3) described similar symptomatology in adult patients. Despite the similarity of symptoms, these two entities remained distinct, and the latter was named adult-onset Still's disease (AOSD). It is characterized by daily fever, evanescent rash, arthritis, pharyngitis, leucocytosis, lymphadenopathy, and less frequently, polyserositis and hepatitis. Pericarditis occurs in approximately 30-40% of patients while cardiac tamponade is relatively rare complication. The diagnosis of AOSD is purely clinical and lacks evidence, hence the need for diagnostic criteria that have been proposed and modified over time.

We report a case of cardiac tamponade that presented as a complication of AOSD adult the onset Still's disease.

Presentation of the case:

A 57-year-old male resident of rural area was admitted to the cardiac intensive care unit presenting with symptoms of NYHA stage IV dyspnea and retrosternal chest pain, both of which had persisted for 5 days alongside a febrile state. The patient had type 2 diabetes, which was being treated with metformin. It is noteworthy that the patient had a 2-year history of untreated polyarthralgia with an inflammatory pattern and, on presentation, had salmon colored rash infectious history was determined to be negative upon inquiry.

1- Clinical findings:

Upon initial evaluation, the patient presented with acute dyspnea and had a peripheral oxygen saturation level of 95% while breathing room air. Vital signs showed a blood pressure BP (100/60 mmHg), a regular heart rate of 110bpm, (the paradoxical pulse). The examination also revealed signs of right heart failure, including spontaneous jugular vein distention and pitting oedema in the lower extremities. The thoracic examination indicated dullness on percussion of bases and muffled heart sounds, with no murmurs or rubs detected. Despite the patient's high fever at 39°, its infectious history was determined to be negative upon review with no clinical suggestive of arthritis.

2- Diagnostic assessment and therapeutic intervention:

Initial laboratory data revealed a leukocytosis of 16 220 with 92% segmented neutrophils, hemoglobin of 16.22g/dL, elevated inflammatory markers; erythrocyte sedimentation rate [ESR] at 97s, C-reactive protein [CRP] at 255m/dl and fibrinogen at 7. Empiric antibiotic therapy was initiated for a presumed infectious etiology, however fever and leukocytosis persisted despite suitable antibiotic therapy. The results of thorough infectious investigation were unremarkable, including blood cultures, fungal cultures and serologies. A chest X-ray revealed marked cardiomegaly, cardiothoracic index (CTI) 0.8. with minor bilateral pleural effusions. The electrocardiogram (ECG) demonstrated sinus tachycardia and diffuse microvoltage. Shortly thereafter, the patient presented with hemodynamic distress as his BP dropped to 85/54. An emergency bedside transthoracic echocardiography (TTE) was performed and revealed a large pericardial effusion, with abnormal respiratory variation in transvalvular blood flow velocities, a paradoxical septum on inspiration, and a significant dilatation of the inferior vena cava, all consistent with the diagnosis of tamponade. In accordance with the institution's protocol on managing cardiac tamponade, an urgent pericardiocentesis was performed, analysis of the pericardial fluid revealed an exudate with negative microbiologic examination results and a large amount of haemorrhagic fluid was drained. The patient's distress abated following the pericardiocentesis fluid was exudates, it yielded negative microbiologic examination including cytology and cultures (aerobic, anaerobic, acid-fast and fungal) and was unremarkable upon histologic analysis. An immunologic screening for rheumatoid factor and anti-nuclear antibodies (ANA) was performed and was also negative. The patient received aspirin 3g/d during the investigation phase with no clinical nor biological improvement.

Once the diagnosis of AOSD was established based on the presence of fever, arthralgia, skin rash and leukocytosis clinically and hyperferritinemia with a decreased glycosylated fraction in the patient's antibodies were discontinued once cultures were finalized as being negative. Corticosteroids were prescribed to induce symptom remission were required to induce symptom remission starting with an intravenous infusion of high-dose methylprednisone for 3 days in a row which was tapered down to prednisolone oral route (1mg/kg/d) with remarkable clinical and biological improvement during the first week with a resolution of fever and the normalization of leukocyte count. Following a rapid prednisolone a rapid prednisolone tapering scheme, the patient was discharged home and was kept on a regimen of prednisone 20mg/dl orally.

3- Follow-up and outcomes:

During the several months of follow-up, the patient was asymptomatic and TTE revealed a complete resolution of pericardial effusion. During the few months of follow-up, the patient was asymptomatic, and CT examination revealed complete resolution of the pericardial effusion. Eight months after hospitalization, the patient was clinically stable and asymptomatic, and was successfully weaned off steroids. Approximately one year after hospitalization, he developed a recurrence of arthralgia and was put back on maintenance doses of steroids).

A close follow up was maintained during the initial months following hospitalization, and it was determined through a transthoracic echocardiogram (TTE) that there was a complete disappearance of the pericardial effusion as well as the pleural effusion, with the patient remaining asymptomatic throughout this time period.

Eight months after the initial hospitalization, the patient continued to exhibit excellent health, and it was deemed appropriate to begin a gradual weaning process off of the steroid medication. Despite this positive progression, approximately one year after the initial hospitalization, the patient experienced a recurrence of arthralgias as he was weaned off prednisone.

In light of this development, it was necessary to restart the patient on a maintenance dose of prednisone in order to manage the symptoms and maintain the overall well-being.

Discussion:

1. "Estimates indicate that the incidence of AOSD in the general population falls within a range of 0.16 to 1.47 per 100,000 individuals. Among patients with AOSD, fluid in the pericardial sac has been identified in 30 to 40% of cases. However, it should be noted that not all individuals with AOSD will exhibit symptoms of pericarditis"(5).
2. "The onset of Still's disease can be sudden and is characterized by a triad of symptoms, including a high fever accompanied by a transient salmon-colored rash appearing during fever peaks and arthralgia".(2) "The diagnosis is established based a constellation of clinical manifestations, laboratory findings, and exclusion of other systemic infections, conditions, and malignancies. Despite the lack of a specific diagnostic test for Still's disease, several authors have attempted to develop criteria that aid in its diagnosis based on observations of various patient groups".(5) "Many diagnostic criteria were proposed, however, the criteria proposed by Yamaguchi et al remain the most commonly used, characterized by a high sensitivity of 93.5%".(6) The constellation of clinical manifestations and laboratory findings in our case fit in the diagnostic criteria of AOSD proposed by Yamaguchi et al. (6)
3. "Serosal involvement is seen in 25% to 60% of all AOSD patients, and cardiac involvement is frequently observed and can be severe. Pericarditis is present in 10% to 40% of patients, with 20% of these patients experiencing pericardial effusion or cardiac tamponade".(7)
"Pericarditis is sometimes discovered accidentally in a patient diagnosed with AOSD on a routine ECG, chest x-ray, or TTE. It is often accompanied by a pleural effusion in 60 to 80% of cases, and presents during the initial flare-up of the disease. Pericarditis or its complications can be the first symptom of AOSD, however, this does not appear to negatively impact the prognosis. Myocarditis is less common, affecting about 3% of patients, and can lead to various complications such as complete atrioventricular block, tachyarrhythmia, heart failure, or cardiogenic shock. Endocardial involvement is uncommon, and can manifest as non-infectious endocarditis".(7)
The occurrence of cardiac tamponade is unusual, especially during a flare-up of AOSD, and only a small number of cases was reported in the literature. When the disease presents with tamponade. The diagnosis of AOSD may be difficult, as other symptoms may be overlooked. If the patient presents with signs of tamponade, the pericardial drainage may be required, either through, ultrasound-guided puncture or a sub-xiphoid surgical procedure. Treatment with high doses of corticosteroids, often in the form of methylprednisolone, has been successful when the diagnosis of AOSD is suspected with no hemodynamic distress. (8)
In AOSD, the pericardial fluid is commonly sero-hematous and exudative. While pericardial biopsies are rarely performed, they often reveal either an acute, non-specific pericarditis or a chronic, oedematous pericarditis. Congestive heart failure has been reported in few cases, and limited endomyocardial biopsies have shown only diffuse, non-specific mono-nuclear inflammation, along with fibroblast proliferation and hypertrophy in certain myocardial fibers. However, endocardial lesions are uncommon in this disease. In rare cases, mitral or aortic valve involvement has been reported, requiring valve replacement in some patients.(9)

Due to the frequency and severity of cardiac involvement in AOSD, it is recommended that all patients undergo regular echocardiographic evaluation to monitor for these complications

4. *"The primary line of treatment for patients with symptoms that warrant medication typically involves the use of nonsteroidal anti-inflammatory drugs (NSAIDs). When necessary, other options such as corticosteroids, immunosuppressive medications like methotrexate or cyclophosphamide, or biologic drugs such as tumor necrosis factor inhibitors, interleukin-1 receptor antagonists, and anti-B-cell antibodies may be prescribed".(10) "In general, cardiac involvement tends to respond well to conservative therapies, but in cases of cardiac tamponade, percutaneous or surgical pericardial drainage may be required". (4) "The prognosis for adult-onset Still's disease is generally favorable, with low mortality rates and a 10% annual recurrence rate during follow-up".(11)*
5. *This case highlights the importance of considering a broad differential diagnosis for patients presenting with cardiac tamponade, including rare conditions like adult onset Still's disease. Additionally, this case underscores the need for a thorough medical evaluation to uncover underlying causes, even in the absence of characteristic symptoms. This is particularly important for patients with pre-existing medical conditions, such as diabetes, who may be at an increased risk for developing secondary conditions.*

Conclusion:

This case report re-emphasizes the importance of considering Still disease in the differential diagnosis of the life-threatening emergency of pericardial tamponade. AOSD often poses a diagnostic and therapeutic challenge and clinical guidelines are lacking. The emergence of validated diagnostic criteria, discovery of better serological markers, and the application of new biological agents may all provide the clinician with significant tools for the diagnosis and management of this complex systemic disorder.

Consent

As per international standard or university standard, patient(s) written consent has been collected and preserved by the author(s).

Ethical Approval:

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

References:

1. Still GF. On a Form of Chronic Joint Disease in Children. *Med Chir Trans.* 1897;80:47-60.9.
2. Bywaters EG. Still's disease in the adult. *Ann Rheum Dis.* 1971 Mar;30(2):121-33.
3. Bujak JS, Aptekar RG, Decker JL, Wolff SM. Juvenile rheumatoid arthritis presenting in the adult as fever of unknown origin. *Medicine (Baltimore).* 1973 Sep;52(5):431-44.
4. Parvez N, Carpenter JL. Cardiac Tamponade in Still Disease: A Review of the Literature: *Southern Medical Journal.* 2009 Aug;102(8):832-7.
5. Bilska A, Wilińska E, Szturmowicz M, Wawrzyńska L, Fijałkowska A, Oniszh K, et al. Recurrent Exudative Pericarditis in the Course of Adult-Onset Still's Disease—Two Case Reports. *Advances in Respiratory Medicine.* 2011 Apr 21;79(3):215-21.
6. Yamaguchi M, Ohta A, Tsunematsu T, Kasukawa R, Mizushima Y, Kashiwagi H, et al. Preliminary criteria for classification of adult Still's disease. *J Rheumatol.* 1992 Mar;19(3):424-30.
7. Carrilho-Ferreira P, Silva D, de Jesus Silva M, André R, Varela MG, Diogo AN. Adult-Onset Still's Disease and Cardiac Tamponade: A Rare Association. *Tex Heart Inst J.* 2015 Jun 1;42(3):277-80.
8. Bodard Q, Langlois V, Guilpain P, Le Quellec A, Vittecoq O, Noel D, et al. Cardiac involvement in adult-onset Still's disease: Manifestations, treatments and outcomes in a retrospective study of 28 patients. *Journal of Autoimmunity.* 2021 Jan;116:102541.
9. Ben Ghorbel I, Lamloum M, Miled M, Aoun N, Houman MH, Pouchot J. Tamponnade révélant une maladie de Still de l'adulte : à propos de deux observations. *La Revue de Médecine Interne.* 2006 Jul;27(7):546-9.

10. Fautrel B. Adult-onset Still disease. *Best Practice & Research Clinical Rheumatology*. 2008 Oct 1;22(5):773–92.

11. Ruscitti P, Cipriani P, Masedu F, Iacono D, Ciccia F, Liakouli V, et al. Adult-onset Still's disease: evaluation of prognostic tools and validation of the systemic score by analysis of 100 cases from three centers. *BMC Med*. 2016 Dec 1;14:194.

UNDER PEER REVIEW