

## Original Research Article

# Relationship between flower galls and girth class of *Prosopis cineraria*

### Abstract

Khejri (*Prosopis cineraria*) recognized as a multipurpose tree species of Thar Desert gets infested with flower galls caused by *Eriophyes prosopidis*. The heavy infestation turns the flowers into galls without forming pods causing high losses in the yield of pods locally known as sangri. The present work was conducted to find out the relation between girth class of tree and average number of galls per inflorescence in the flower gall infested trees. In the present study we concluded that the relation between girth class of tree and average number of galls per inflorescence is negative. The average number of mite induced galls per inflorescence was highest in the girth class 70-100 cm and least in girth class 130-160 cm.

**Key words:** *Prosopis cineraria*, *Eriophyes prosopidis*, girth class, galls, inflorescence

### Introduction

The life-supporting tree of the Thar Desert called khejri (*Prosopis cineraria*) is the most crucial element of the traditional farming systems used in the arid and semi-arid regions of India's northwest. It is a key source of premium feed, fuel, and lumber for the rural economy. It has thick, green foliage that tolerates harsh edapho-climatic conditions, and it produces fruit even during the driest season. Its pods (sangri) and nourishing leaf-fodder (loong) are widely used. Sangri is the primary ingredient in the vegetables Panchkutta and Trikuta and it is offered both fresh and dehydrated at high prices. Its beneficial effects on the ecology and the development of soil fertility are well known. Typically, 4-5 kg of pods is produced per plant in healthy trees (Rathore, 2009). Despite the considerable correlation between pod production and tree diameter at breast height, arid areas have been shown to generate roughly 1.4 quintals of pods/ha with a variance of 10.7% (Rathore, 2009). Flower galls are to be blamed for the loss of pods and seed production in *P. cineraria*. In Khejri trees these galls are the result of an eriophyid mite infestation called *Eriophyes prosopidis*, which causes severe pod reduction and gall formation. The effects of eriophyid feeding on their hosts may not always be predictable in advance. Pale yellow flowers turned green at the bases and the size of the ovary began to bulge at the beginning stages of gall growth. Instead of pods, various kinds of galls, such as circular, rectangular, egg-shaped, spindle-shaped, and vase-shaped, forms. A great number of dangling, disorganised, and deformed green galls give gall-infested trees a sickly appearance during the summer. Flower galls significantly reduce the pod yield. As the infestation of flower galls causes economic losses to the farmers therefore, the studies on the relation between girth class of khejri tree and average number of galls per inflorescence deserve to be discussed.

The current study was carried out to investigate the effect of tree girth on floral gall formation in *Prosopis cineraria*.

## Review of literature

Gall mites and other arthropod species, as well as other herbivores, commonly attack crop plants in both natural and farmed environments (Glase *et al.*, 2014). Gall mites are also difficult to control in agricultural settings due to their small size (Lindquist, 1996), high reproductive potential (Sabelis and Bruin, 1996), and the inefficiency of existing miticides (Chong, 2013). Galls can be caused by a wide range of species, including mites, aphids, wasps, and midges (Redfern *et al.*, 2011). Eriophyoid mites are generally phytophagous arthropods that form intimate relationships with the plants that serve as their hosts (Acari: Eriophyoidea). These highly specialized mites are found on both annual and perennial plants, and they have a potent monophagy. Gall production is triggered by insects or mites that feed on plants or by the release of growth hormones, which then interact with the host plant's defense mechanism (Ananthkrishnan, 1984). The pests' interactions with one another lower plant defenses (Kiekiewicz *et al.*, 2019). Eriophyoids are obligate plant-feeders with extraordinary morphological, biochemical, and behavioural specialisation in comparison to other Acari members, according to Skoracka *et al.*, (2010). Despite the fact that many of them are major pests of commercial and ornamental crops, as well as wild plants, grasses, and plants in urban and community forestry, they rarely cause the mortality of these plants (Lindquist *et al.*, 1996). The types of plant modifications brought on by eriophyoids depend significantly on the genotypes, age, and lifestyle of the host plants as well as the species and strains of the mites (Petanovi and Kielkiewicz, 2010a,b; Skoracka *et al.*, 2010; Chakrabarti *et al.*, 2011; Cvrkovi, 2012).

## Methodology:

Five distinct sites in Rajasthan were chosen to evaluate the effect of tree girth on flower gall formation in Khejriviz., Phalodi, Lohawat, Osian, Pipar and Baori. Ten flowering trees of three groups (70-100, 100-130cm and 130-160 cm) each based on girth were selected at each site. By comparing the presence and absence of flower gall in the chosen trees, the number of trees with flower gall infestation was determined. Also for collecting data on flower gall formation per inflorescence, canopy of each selected tree was divided into four horizontal quadrants, representing all the four directions namely, North, South, East and West. From each quadrant 10 inflorescences were collected and number of healthy pods and flower galls were counted in each inflorescence and average no. of galls per inflorescence was calculated.

## Statistical analysis

The raw data from every location that was sampled over the course of five years was converted in an electronic format and laid up in a spreadsheet (Microsoft Excel, 2016). The data thus collected were subjected to statistical analysis using analysis of mean, standard error, standard deviation, correlation and variance by using software NCSS 2023 Version 23.0.1.

## Results

The data was collected for consecutive five years from five locations Phalodi, Lohawat, Osian, Pipar and Baori Tehsil in District Jodhpur, Rajasthan (Fig 4 & 5). It was observed during the study that the average no. of galls

per inflorescence (Table 1; Fig 1, 2) was found to be highest in the year 2020-21 at Phalodi 13.57(70-100 cm), 9.3(100-130cm) and 9.15 (130-160 cm) followed by lohawat viz., 13.15 (70-100 cm), 8.88 (100-130cm) and 8.7 (130-160 cm) and least in Baori viz., 11.9 (70-100 cm), 8.08 (100-130cm), 7.75 (130-160 cm). The average no. of galls per inflorescence of all the sites was highest in the year 2020-21 viz., 12.62 (70-100 cm), 8.52 (100-130cm) and 8.26(130-160 cm). Thus we conclude that the galls per inflorescence is highest in girth class 70-100 cm followed by girth class 100-130cm and least in girth class 130-160 cm. At all of the research sites, the average number of galls per inflorescence was larger on younger trees with girth classes of 70 to 100 cm than it was on older trees with girth classes of 130 to 160 cm. On taking the combined averages of all the sites and years, the average number of galls per inflorescence was found to be  $11.53 \pm 0.31$  which is highest among all girth classes under study (Table 2).

The average no. of galls per inflorescences was calculated taking all the sites and using regression analysis, the number of flower gall per inflorescence was compared with DBH.


**Table 1. The average no. of galls per inflorescence**

Girth class (cm)	TREE	Average per inflorescences (Yearly)				
		2017-2018	2018-2019	2019-2020	2020-2021	2021-2022
		No. of Galls	No. of Galls	No. of Galls	No. of Galls	No. of Galls
70-100	Phalodi	11.7	11.75	12.46	13.57	11.3
	Lohawat	11.33	11.70	11.45	13.15	11.10
	Osian	11.03	11.70	11.50	12.35	10.82
	Pipar	10.65	11.60	11.40	12.13	10.35
	Baori	10.45	11.23	11.08	11.90	10.50
	Mean	11.03	11.59	11.58	12.62	10.82
	Correlation	-0.40	-0.86	-0.32	-0.31	-0.47
	Variance	0.25	0.04	0.27	0.50	0.15
100-130	Phalodi	9.15	7.82	8.63	9.30	8.43
	Lohawat	8.35	7.35	7.85	8.88	7.83
	Osian	7.83	6.93	7.6	8.18	7.20
	Pipar	7.58	6.83	7.38	8.18	6.88
	Baori	7.58	6.75	7.45	8.08	7.13
	Mean	8.09	7.13	7.78	8.52	7.49
	Correlation	-0.23	-0.29	-0.37	-0.271	-0.31
	Variance	0.44	0.19	0.25	0.29	0.39
130-160	Phalodi	8.95	7.70	8.48	9.15	8.28
	Lohawat	8.30	7.34	7.75	8.70	7.78
	Osian	7.60	6.68	7.25	7.93	7.00
	Pipar	7.43	6.55	7.10	7.80	6.90
	Baori	7.40	6.58	7.03	7.75	6.60
	Mean	7.936	6.97	7.522	8.266	7.312
	Correlation	-0.21	-0.262	-0.102	-0.117	-0.23
	Variance	0.454	0.270	0.365	0.391	0.482

**Table 2.** The average number of galls per inflorescence and standard deviation based on girth class.

Girth Class	Average no. of galls per inflorescence	Standard Deviation
<b>70-100cm</b>	11.53±0.31	0.69
<b>100-130cm</b>	7.80±0.24	0.54
<b>130-160cm</b>	7.60±0.23	0.51

Regression analysis showed a highly significant inverse relationship between DBH and average number of galls per inflorescence in Khejritrees in Agro-forestry habitats ( $R^2 = 0.796$ ). The normal distribution curve (fig.3) represents that the number of galls per inflorescence is maximum in the girth class 70-100cm. As we know that normal distribution graph is a continuous probability function therefore from the graph it is clear that occurrence of a flower galls are more in younger trees of girth class 70-100 in comparison to older trees.



**Fig 1.** Graphical representation of Average number of galls per inflorescence per year




Fig 2. Graphical representation of Average number of galls with Girth class



Fig 3. Graphical distribution of Distribution Mean vs Year.



**Fig 4: Khejri flower gall**



**Fig 5: Girth wise data collection**

### **Discussion**

The lower defense mechanisms of young trees may be the cause of the larger number of galls per inflorescence in the girth class of 70-100 cm which is supported by findings of Mutuet *al.* (2010) that *L. invasa* infestation on young (1-3 year old) trees is more severe than on older trees. According to the findings of Madoffe (1989), young pine trees were more susceptible to Pine Woolly Aphid infection than older trees. According to Rohfritch (1981), the accumulation of secondary defense chemicals, such as tannins, their phenolic precursors, and lignin, is a form of defense against injury or the presence of invading organisms. According to Rohfritch (1981), all secondary defense compounds rise in concentration with tree age. Additionally, infestation may be greater in young trees due to their weaker insect defense mechanisms. In contrary, According to Ruohomäkiet *al.*, (2000), *Epirrita autumnata* outbreaks primarily occurred in mature birch trees due to minimal parasitism or good foliage quality as well as the availability of more advantageous oviposition locations in mature trees.

Depending on the genotype of the mite and plant, the density of the mites, the feeding period, the age of the plant, and the environment, some of them displayed a noticeable degree of variability (Petanovi and Kielkiewicz, 2010a, b; Duso *et al.*, 2008; Royalty and Perring, 1996; Westphal and Manson, 1996). Salivary secretions must play a significant role because mechanical action only triggers the synthesis of chitosan and callose at the feeding site as a plant's wound response (Petanovi and Kielkiewicz, 2010a, b). It is possible that pectinolytic and cellulolytic enzymes are involved in the production of galls given that the middle lamella and cell walls can be destroyed by the saliva of piercing and sucking insects. Because of this, it is crucial to understand the physiological changes in infected trees, their genetic foundation, and heredity as well as host plant relationships and preferences within

various girth classes in order to develop resistant varieties (Mitchell *et al.*, 2016; Stenberg and Muola, 2017). This will provide a base for future studies to develop a resistant variety against the flower gall mite.

## Conclusion

Present study concludes that the eriophid mites induced flower galls per inflorescence were highest in girth class 70-100 cm followed by girth class 100-130cm and least in girth class 130-160 cm showing a highly significant inverse relationship between DBH and average number of galls per inflorescence in Khejri (*Prosopis cineraria*) trees.

## References

1. Ananthkrishnan T.N. (1984): Biology of Gall Insects. Edward Arnold, London, 376 pp.
2. Chong, J.H. (2013). A witch's brew of troubles with the Bermuda grass mite. USGA Green. Sect. Rec., 51, 4p.
3. Cvrkovic, T., Chetverikov, P., Vidovic, B., and Petanovic, R. (2012). "Molecular Analysis of COI mtDNA in Phytoptus (Phytoptidae) and Eriophyes (Eriophyidae) species associated with galls of Tilia spp. (Tiliaceae): preliminary results," in Proceedings of the International Symposium on Current Trends in Plant Protection, (Belgrade, Serbia).
4. Glas, J.J.; Alba, J.M.; Simoni, S.; Villarroel, C.A.; Stoops, M.; Schimmel, B.C.; Schuurink, R.C.; Sabelis, M.W.; Kant, M.R. Defense suppression benefits herbivores that have a monopoly on their feeding site but can backfire within natural communities. BMC Biol. 2014, 12, 98.
5. Chakrabarti, S., Chakrabarti, S., and Chakrabarti, S. (2001). Effect of infestation of *Acalitushibisci* (Eriophyoidea: Eriophyidae), a gall forming mite on age and hairs of leaves of *Hibiscus vitifolius*. Acarologia 41, 313–316.
6. Kielkiewicz, M.; Barczak-Brzyzek, A.; Karpińska, B.; Filipecki, M. Unravelling the Complexity of Plant Defense Induced by a Simultaneous and Sequential Mite and Aphid Infestation. Int. J. Mol. Sci. 2019, 20, 806.
7. Lindquist, E. E., Sabelis, M. W., and Bruin, J. (1996). Eriophyoid Mites-Their Biology, Natural Enemies and Control. World Crop Pests, Volume 6. Amsterdam: Elsevier Science Publishers.
8. Lindquist, E.E. (1996). External anatomy and notation of structures. In Eriophyoid Mites: Their Biology, Natural Enemies and Control; Lindquist, E.E., Sabelis, M.W., Bruin, J., Eds.; Elsevier: Amsterdam, The Netherlands, 6:3–32.
9. Mitchell, C., Brennan, R. M., Graham, J., and Karley, A. J. (2016). Plant defense against herbivorous pests: exploiting resistance and tolerance traits for sustainable crop protection. Front. Plant Sci. 7:1132.
10. Madoffe, S.S. (1989). Infestation densities on the pine woody aphid (*Pineuspini*) on *Pinuspatula* as related to site productivity at Sao-Hill Forest Plantation (Unpublished master's thesis). University of Dares Salaam, Dares Salaam, Tanzania.
11. Mutitu, K. E., B.O. Otieno, P. Nyeko and G.N. Ngae (2010). Variability in the infestation of *Leptocybeinvasa* (Hymenoptera: Eulophidae) on commercially grown Eucalyptus germplasm in Kenya. In M. Imo, H. Ipara, L. Etiegni, C.M. Mulewa, F. Muisu, J.M. Njiru and B.B. Kirongo (Eds.). Natural resource management for improved livelihoods (pp. 115–120). Eldoret, Kenya: Moi University, School of Natural Resource Management.

12. Petanović, R., and Kielkiewicz, M. (2010a). Plant-eriophyoid mite interactions: cellular biochemistry and metabolic responses induced in mite-injured plants. *Exp. Appl. Acarol.* 51, 61–80.
13. Petanović, R., and Kielkiewicz, M. (2010b). Plant-eriophyoid mite interactions: specific and unspecific morphological alterations. Part II. *Exp. Appl. Acarol.* 51, 81–91.
14. Rathore, M. (2009). Nutrient content of important fruit trees from arid zone of Rajasthan. *Journal of Horticulture and Forestry*, 1: 103-08.
15. Redfern M., Shirley P. & Bloxham M. 2011: *British Plant Galls*. FSC Publications, Schremsbury, 436 pp.
16. Ruohomäki, K., M. Tanhuanpää, M.P. Ayres, P. Kaitaniemi, T. Tammaru and E. Haukioja (2000). Causes of cyclicity of *Epirrita autumnata* (Lepidoptera, Geometridae): Grandiose theory and tedious practice. *Society Population Ecology*, 42: 211–223.
17. Rohfritsch, O. (1981). A “defense” mechanism of *Picea excelsa* L., against the gall former *Chermes abietis* L. (Homoptera: Adelgidae). *Journal of Applied Entomology*, 92: 18–26.
18. Royalty, R. N., and Perring, T. M. (1996). “Nature of damage and its assessment,” in *Eriophyoid Mites- Their Biology, Natural Enemies and Control*, eds E. E. Lindquist, M. W. Sabelis, and J. Bruin (Amsterdam: Elsevier Science B.V), 493–512.
19. Sabelis, M.W.; Bruin, J. (1996). Evolutionary ecology: Life history patterns, food plant choice and dispersal in *Eriophyoid Mites: Their Biology, Natural Enemies and Control*; Lindquist, E.E., Sabelis, M.W., Bruin, J., Eds.; Elsevier: Amsterdam, The Netherlands; Volume 6, pp. 329–366.
20. Skoracka, A., Smith, L., Oldfield, G., Cristofaro, M., and Amrine, J. W. Jr. (2010). Host-plant specificity and specialization in eriophyoid mites and their importance for the use of eriophyoid mites as biocontrol agents of weeds. *Exp. Appl. Acarol.* 51, 93–113.
21. Stenberg, J. A., and Muola, A. (2017). How should plant resistance to herbivores be measured? *Front. Plant Sci.* 8:663.