

Minireview Article

Virulence and resistance factors associated with the specie of *Proteus vulgaris*

ABSTRACT

Proteus vulgaris belongs to the Enterobacteriaceae family, colonizing mainly the gastrointestinal tract of animals, including humans. However, when there is some deregulation in the host's immune system, this pathogen spreads, causing several clinical manifestations. Currently in the literature little is discussed about the virulence and resistance profile of this species. Thus, the study aimed, through a literature review, to describe the main virulence factors and resistance of this strain. Studies have shown that *P. vulgaris* is associated with most Urinary Tract Infections (UTIs), despite being less isolated from *P. mirabilis*, exhibiting high motility ability related to its fimbriae, enzyme and endotoxin secretion, ability to produce urease and biofilm formation. As for resistance, the pathogen is intrinsically resistant to several antibiotics, requiring further studies for the development of new therapeutic alternatives.

Formatted: Font: Italic

Formatted: Font: Italic

Keywords: *Infections; Proteus vulgaris; Resistance; Virulence.*

1. INTRODUCTION

The genus *Proteus* spp., belongs to the Enterobacteriaceae family, characterized as facultative anaerobic Gram-negative bacilli, having five species well described taxonomically: *Proteus mirabilis*, *Proteus vulgaris*, *Proteus penneri*, *Proteus hauseri* and *Proteus myxofaciens*[1]. These are part of the enteric microbiota of animals, regardless of cold-blooded or warm-blooded species, including humans, in addition to being found in contaminated water and food. Among the species, *P. mirabilis* and *P. vulgaris* stand out, as they are clinically relevant due to their pathogenicity [2].

P. mirabilis and *P. vulgaris* have the ability to colonize and cause infectious processes in humans, being characterized as opportunistic microorganisms [reference](#). Such pathogens make use of conditions that make the individual's immunity vulnerable to cause various manifestations, from a simple urinary tract infection to more serious infections, such as meningitis and generalized infections [3]. Several factors contribute to the pathogenicity of these species, facilitating the onset of infection and the establishment of the microorganism at the site of infection.

Thus, the present study aimed to analyze the literature regarding the main characteristics of the bacterium *P. vulg* well as to investigate the main virulence and resistance factors involved with the microorganism. ~~The information was obtained through data collection carried out by bibliographic survey, where platforms such as Google Scholar (<https://scholar.google.com.br/>), PUBMED (<https://pubmed.ncbi.nlm.nih.gov/>), SciELO (<https://www.scielo.org/>), Science Direct (<https://www.sciencedirect.com/>), Scopus® (<https://www.scopus.com/>), LILACS (<https://lilacs.bvsalud.org/>) and BDTD (<https://bdtd.ibict.br/>). As an inclusion criterion, we considered articles published in the previously mentioned journals between the period of 2013 to 2023, which addressed the theme in question, and which were in Portuguese and English.~~

Formatted: Font: Italic

MATERIAL AND METHODE

The information was obtained through data collection carried out by bibliographic survey, where platforms such as Google Scholar (<<https://scholar.google.com.br/>>), PUBMED (<<https://pubmed.ncbi.nlm.nih.gov/>>), SciELO (<<https://www.scielo.org/>>), Science Direct (<<https://www.sciencedirect.com/>>), Scopus@ (<<https://www.scopus.com/>>), LILACS (<<https://lilacs.bvsalud.org/>>) and BDTD (<<https://bdtb.ibtci.br/>>). As an inclusion criterion, we considered articles published in the previously mentioned journals between the period of 2013 to 2023, which addressed the theme in question, and which were in Portuguese and English.

2. DEVELOPMENT

2.1 General enterobacteria and epidemiology of *P. vulgaris*

Proteus vulgaris is a short rod-shaped Gram-negative opportunistic enterobacterium, 0.4-0.8 µm in diameter and 1.0-3.0 µm in length, has motility through peritrichous flagella, is lactose and lysine decarboxylase negative and have the ability to hydrolyze urea [reference](#). It is commonly found colonizing the gastrointestinal tract of animals, forming part of the normal microbiota, and in food and water contaminated with waste [4]. Its pathogenic form is mainly associated with urinary tract infections (UTI), being responsible for more severe forms with formation of kidney stones [reference](#). However, this is also related to more serious conditions such as meningitis, wound infection, rheumatoid arthritis and septicemia in patients with burns [5].

In a study carried out by scientists, the isolation of 11 strains identified as *P. vulgaris* was verified, eight from manure, two from animals (dog) and only 1 from human origin (nasal mucosa) [5]. Having already investigated the bacteria that cause urinary infections, it was reported that of the 136 isolates at the Center for Clinical Analysis Diagnostics between the years 2017-2018, 4.5% (6 strains) were identified as *P. vulgaris* among female patients and males ranging in age from 3 to 89 years [6]. Furthermore, researchers point out that the main agents isolated from the urine of domestic animals and humans are *Escherichia coli* (37.64%), *Staphylococcus* spp. (19.66%) and *Proteus* spp. (13.48%) [7]. Reinforced by studies indicating *E. coli* and *Proteus* sp. as the main causes of UTIs [8].

Analyses also reported the isolation of *P. vulgaris* from UTIs in patients treated at a metropolitan hospital, it was seen that *P. vulgaris* and *P. mirabilis* per se corresponded to 5.63% of isolations in patients aged 0 to > 60 years of age [9]. However, what has been observed is that despite its pathogenicity, *P. vulgaris* is still less isolated than *P. mirabilis*. In studies, 73 strains were isolated from the urine culture of domestic animals, of which 94.51% (64 strains) corresponded to *P. mirabilis* and only 5.49% (4 strains) were identified as *P. vulgaris* [10]. Even so, despite this difference, *P. vulgaris* has several virulence factors that contribute to its pathogenicity. In addition, it is important to emphasize that the high number of isolates from domestic animals alert us to a public health problem due to the high ease of transmission to humans, especially those with impaired immune systems, children and the elderly [11].

2.2 Virulence Factors

Several conditions favor the transition from the commensal to the pathogenic form, facilitating the onset of infection and the establishment of the pathogen, from factors that are in the cell structure itself, such as fimbriae and flagella, to the secretion of enzymes and endotoxins [4]. The simultaneous expression of different types of fimbriae is directly associated with the ability of the microorganism to adhere to different surfaces, including epithelial cells [reference](#). These structures are proteinaceous, filamentous with high specificity for cell-cell and bacteria-host adhesion. In addition, the mobility of this species in cycles by peritrichous flagella helps in the dissemination of the pathogen at the site of infection [12-13].

Thus, *Proteus vulgaris* exhibits the ability to swarm due to its high mobility. Studies point out that this bacterium can trigger the differentiation between cells with short rods to extremely elongated swarm cells with many flagella, which facilitates their displacement and dissemination on surfaces [14].

Furthermore, the presence of LPS (Lipopolysaccharide) represents great virulence in Gram-negative bacteria, including *P. vulgaris*, responsible for stimulating the action of prostaglandins and pro-inflammatory cytokines that can result in toxic shock syndromes [10]. Another characteristic of this species is the production of hemolysin, responsible for forming pores in eukaryotic membranes and the action of an amino acid deaminase that acts as a siderophore due to its affinity and capture of the iron ion, which is essential for the metabolism of *P. vulgaris* [15].

83 Another virulence factor that has been much discussed in urinary tract infections due to the dissemination of *P. vulgaris* is
84 its ability to produce urease, causing the formation of stones [reference](#). This is due to the hydrolysis of urea into ammonia
85 and CO², changing the pH of the urine and causing the precipitation of ions with consequent formation of stones
86 [reference](#). These, in turn, obstruct the normal flow of urine, in addition to the high concentrations of urea being extremely
87 harmful to the urinary epithelium [16]. *Proteus* urolithiasis is difficult to treat, given that the bacteria can be located inside
88 the calculus, where it keeps multiplying and can easily spread to other organs [reference](#).

89 As in other bacteria, biofilm formation also configures an important virulence factor, favoring the permanence of the
90 pathogen at the site of infection and making it difficult to eradicate the microorganism. These microbial communities are
91 recognized for becoming up to 1000 times more resistant when compared to their planktonic form, because the
92 microorganisms are surrounded by an exopolymeric matrix composed of polysaccharides, proteins, extracellular DNA and
93 lipids, which hinders the action of the cells of the ~~system~~ immune system and the introduction of antibiotics [17].

94 Biofilms are formed through multiple stages, from bacterial adhesion to biotic and abiotic surfaces, through attachment,
95 formation of small colonies, production of exopolymer matrix, to the mature biofilm stage, where cells that are in the upper
96 layers begin to come off to colonize other surfaces [17]. In the case of *Proteus*, biofilms are formed from the initial
97 adhesion of adhesins present in the fimbriae, resulting in crystalline structures that are mainly associated with obstruction
98 of urinary catheters [18].

100 2.3 Resistance

101 Several studies have shown the resistance profile of *P. vulgaris* to some antibiotics, pointing to the urgent need for new
102 therapeutic alternatives that can contribute to the control of these infections. According to the Clinical & Laboratory
103 Standards Institute, *Proteus vulgaris* presents intrinsic resistance to some antibiotics, among which are: ampicillin,
104 cephalosporin I – cefazolin and cefotetan, cephalosporin II – cefuroxime, tetracyclines – tigecycline, nitrofurantoin,
105 polymyxin B and colistin [19]. However, this feature is commonly found in bacterial genomes, without the influence of
106 selective pressure exerted by antimicrobials, and is not even related to horizontal gene transfer [20].

108 Investigations demonstrated the isolation of 17 strains of *P. vulgaris* from urinary infections, of these, about 94% showed
109 resistance to the antibiotics ampicillin, tigecycline and chloramphenicol, while 12% showed resistance to imipenem [4]. In
110 addition to this, other studies also reported the resistance of this species, with approximately 70% of the isolates
111 presenting this characteristic to nitrofurantoin [21]. However, in contrast to these findings, a study with isolates of *Proteus*
112 spp. from domestic animals, showed high in vitro sensitivity to the antibiotics imipenem, norfloxacin, amikacin,
113 levofloxacin, ceftriaxone, florfenicol, marbofloxacin and amoxicillin/clavulanic acid, suggesting the efficacy of
114 carbapenemics, fluoroquinolones, aminoglycosides, [3rd generation cephalosporins](#) and amphenicols, while
115 73 isolates showed resistance to novobiocin and sulfamethoxazole-trimetopine [10]. In general, the study indicates a
116 percentage of resistance to multidrugs of 45.21% (33 strains). It is noteworthy that the high sensitivity profile is associated
117 with the restricted use of these drugs in veterinary medicine, which is not reflected in human medicine [22].

118 Studies evaluating the resistance profile of *P. vulgaris* isolated from UTIs in a metropolitan hospital. It was seen that 100%
119 of the isolates showed resistance to ampicillin, cefepime, fluoroquinolones ciprofloxacin and levofloxacin, nitrofurantoin
120 and cephalothin and sensitivity to amikacin, cefoxitin, gentamicin, imipenem, norfloxacin, ceftriaxone, piperacillin-
121 tazobactam and tobramycin. These findings suggest the importance of correctly identifying the microorganism and using
122 the TSA (Antimicrobial Sensitivity Test), ensuring the best treatment targeting and preventing the development of more
123 resistance [9].

125 4. CONCLUSION

126 *Proteus vulgaris* is one of the most important pathogens that cause Urinary Tract Infections, exhibiting several virulence
127 factors that facilitate and condition its permanence in the site of infection, related to mobility, secretion of enzymes and
128 endotoxins, in addition to biofilm formation. However, this species has intrinsic resistance to several antibiotics, which
129 demonstrates the need for the development of new therapeutic alternatives. In general, the current literature does not
130 have many studies regarding the pathogenicity of this species, which also reinforces the need for further epidemiological
131 and clinical studies.

133
134
135
136

REFERENCES

- 137 1. KishanRaj, S., Sumitha, S., Tang, TH, Citartan, M., & Chinni, SV (2021). Identificação e caracterização genômica
138 comparativa de homólogos de npcRNA em *Proteus vulgaris*. *Journal of biosciences*, 46, 1-10.
- 139 2. Ribeiro, MG, Zappa, V., Bolaños, CAD, de Paula, CL, Callefe, JLR, Alves, AC, ... & Melville, PA (2017). Índice de
140 resistência múltipla antimicrobiana, concentrações inibitórias mínimas e produtores de beta-lactamase de espectro
141 estendido de cepas de *Proteus mirabilis* e *Proteus vulgaris* isoladas de animais domésticos com várias manifestações
142 clínicas de infecção. *Semina: Ciências Agrárias*, 38 (2), 775-789.
- 143 3. Ronanki, S. P., Ramya, P., Babu, A. J., & Sreedevi, B. (2022). Multidrug resistant-*Proteus mirabilis* and *Proteus*
144 *vulgaris* isolated from milk and meat samples in Tirupati, Andhra Pradesh: An emerging public health threat.
- 145 4. Bilal, S., Anam, S., Mahmood, T., Abdullah, RM, Nisar, S., Kalsoom, F., ... & Anjum, FR (2019). Perfil antimicrobiano e
146 caracterização molecular de genes resistentes a antibióticos de *Proteus vulgaris* isolados de hospital terciário, Islamabad,
147 Paquistão. *Jornal de ciências farmacêuticas do Paquistão*, 32.
- 148 5. Serafim, MLRC. Identificação e perfil de resistência a antimicrobianos de bactérias isoladas de diferentes amostras
149 provenientes do aterro controlado da cidade de Campos dos Goytacazes- RJ. 2013. Dissertação (Mestrado em Ciência
150 Animal), Universidade Estadual do Norte Fluminense, Campos dos Goytacazes, 2013.
- 151 6. Cabeço, A. L. B., & Colombo, T. E. (2019). Bactérias causadoras de infecções urinárias e seu perfil de resistência aos
152 antimicrobianos. *Journal of the Health Sciences Institute*, 37(2), 113-8.
- 153 7. Furini, A. A. C., dos Santos Silva, B. T. O., Chiaparin, J., Ramos, M. P. S. C. M., Martins, E. A., Atique, T. S. C., ... &
154 de Castro, K. F. (2013). **Análise epidemiológica, identificação e perfil de susceptibilidade a antimicrobianos de isolados de**
155 **cães com infecção do trato urinário**. *Acta Veterinaria Brasílica*, 7(4), 288-293.
- 156 8. Marques, C., Belas, A., Aboim, C., Triqueiro, G., Cavaco-Silva, P., Gama, L. T., & Pomba, C. (2019). Clonal
157 relatedness of *Proteus mirabilis* strains causing urinary tract infections in companion animals and humans. *Veterinary*
158 *microbiology*, 228, 77-82.
- 159 9. Araujo, K. L., & Queiroz, A. C. D. (2012). Análise do perfil dos agentes causadores de infecção do trato urinário e dos
160 pacientes portadores, atendidos no Hospital e Maternidade Metropolitano-SP. *J Health Sci Inst*, 30(1), 7-12.
- 161 10. Zappa, V. Índice de resistência múltipla aos antimicrobianos, concentração inibitória mínima e beta-lactamases de
162 espectro estendido em linhagens de *Proteus mirabilis* e *Proteus vulgaris* isoladas de diferentes afecções em animais
163 domésticos. 2015. Tese (Doutorado em Medicina Veterinária). Universidade Estadual Paulista, Botucatu, 2015.
- 164 11. Bahashwan, S. A., & El Shafey, H. M. (2013). Antimicrobial resistance patterns of *Proteus* isolates from clinical
165 specimens. *European Scientific Journal*, 9(27).
- 166 12. Quinn, P. J., Markey, B. K., Leonard, F. C., Hartigan, P., Fanning, S., & Fitzpatrick, E. (2011). *Veterinary microbiology*
167 *and microbial disease*. John Wiley & Sons.
- 168 13. Beltrão, EMB. Análise do perfil genético de virulência, resistência e relação clonal de isolados clínicos de *Proteus*
169 *mirabilis* carreadores e não carreadores do gene blaKPC e blaNDM provenientes de um hospital de Recife-PE. 2021.
170 Tese (Doutorado em Medicina Tropical). Universidade Federal de Pernambuco, Recife, 2021.
- 171 14. Wasfi, R., Abdellatif, G. R., Elshishtawy, H. M., & Ashour, H. M. (2020). First-time characterization of viable but
172 non-culturable *Proteus mirabilis*: Induction and resuscitation. *Journal of Cellular and Molecular Medicine*, 24(5), 2791-
173 2801.
- 174 15. Siqueira, A. K., Ribeiro, M. G., Salerno, T., Takahira, R. K., Lopes, M. D., Prestes, N. C., & Silva, A. V. (2008). Perfil
175 de sensibilidade e multirresistência em linhagens de *Escherichia coli* isoladas de infecção do trato urinário, de piometra e
176 de fezes de cães. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia*, 60, 1263-1266.

Formatted: Font color: Red

- 177 16. Malinowski, E. (2021). MOST FREQUENT BACTERIA IN URINARY TRACT INFECTIONS. *Revista Saúde e Meio*
178 *Ambiente*, 12(1), 121-134.
- 179 17. Wasfi, R., Hamed, S. M., Amer, M. A., & Fahmy, L. I. (2020). Proteus mirabilis biofilm: Development and therapeutic
180 strategies. *Frontiers in cellular and infection microbiology*, 10, 414.
- 181 18. Filipiak, A., Chrapek, M., Literacka, E., Wawszczak, M., Głuszek, S., Majchrzak, M., ... & Adamus-Białek, W. (2020).
182 Pathogenic factors correlate with antimicrobial resistance among clinical Proteus mirabilis strains. *Frontiers in*
183 *microbiology*, 11, 579389.
- 184 19. CLSI, E. (2016). Clinical and Laboratory Standards Institute. *Performance standards for antimicrobial susceptibility*
185 *testing*, 19087-1898.
- 186 20. Cox, G., & Wright, G. D. (2013). Intrinsic antibiotic resistance: mechanisms, origins, challenges and
187 solutions. *International Journal of Medical Microbiology*, 303(6-7), 287-292.
- 188 21. Lazm, A. M., Jebur, M. S., & Alomashi, G. B. (2018). Sequencing of HpmA Gene in Proteus mirabilis of UTIs among
189 rheumatoid arthritis patients. *J. Pharm. Sci. Res*, 10(2), 265-271.
- 190 22. Greene, C. E. (2012). Infectious Diseases-of the dog and cat. St Louis. *MO: Elsevier Saunders*.
191

Formatted: Font color: Red

UNDER PEER REVIEW