

Deciphering variability and genetic parameters in traditional scented rice (*Oryza sativa* L.) genotypes of Kerala

ABSTRACT

Gandhakasala and *Jeerakasala*, the most popular traditional scented rice cultivars of Wayanad district in Kerala was used to assess the nature and magnitude of variability and genetic parameters in this investigation. The study on twenty yield and associated quantitative, cooking and biochemical characters revealed significant differences among genotypes for each character except for leaf width indicating the presence of high genetic variability among the genotypes. Phenotypic and genotypic coefficient of variations showed high level of variability for grain yield (42.19%), number of spikelets panicle⁻¹ (31.36%) and test weight (26.23%) and moderate level of variability for leaf length, culm number, culm length, number of productive tillers, number of spikelets panicle⁻¹, panicle length, straw yield, grain length, grain width, amylose content and protein content. For most of the characters studied, the difference between GCV and PCV was narrow, indicating that environmental factors have little influence on these characters. Heritability estimations ranged from low to high (17.86% to 99.90%). Characters like leaf length, culm number, number of spikelets panicle⁻¹ and straw yield showed high heritability coupled with high genetic advance reflecting that these characters were under additive gene action and selection for improvement might be effective. The present study suggests that diverse scented rice germplasm in Kerala could be used for the improvement in grain yield and excellent quality traits in short grain scented rice breeding programmes.

Keywords: Scented rice, *Gandhakasala*, *Jeerakasala*, variability, heritability, genetic advance

1. INTRODUCTION

Rice (*Oryza sativa* L.) belonging to the family Poaceae is the second largest cereal crop in the world and serves as the staple food of more than half of the global population. Scented rice, constitutes a small but special group of rice, which is considered best in aroma, quality and are highly priced [1]. Despite their low yield, are endowed with aroma, taste, excellent cooking and eating quality attributes and enjoy local consumer preference with the market being meagre. Non-Basmati traditional scented rice cultivars are generally small and medium grained [2]. All the major rice growing countries including India have their own traditional and evolved scented rice varieties. The genetic diversity of scented rice in India is largest in the world having more than 300 scented rice genotypes cultivated in different states. Traditional scented rice cultivars of Kerala include *Gandhakasala*, *Jeerakasala*, *Velumbala*, *Chomala*, *Kayama*, *Kothampalariikkayama*, *Pookkilathari*, *Mullanchanna*, *Poothadikayama*, *Kunjinellu Neycheera* and *Urunikazama* [3,4,5]. The scented rice cultivation is on the foothills and requires ideal climatic situations like lower temperature, soil conditions and humidity, which are essential for developing and retaining high aroma [6]. Wayanad district which adorns the northern hills of Kerala, ranks first in the cultivation of traditional scented rice cultivars *Gandhakasala* and *Jeerakasala* which are the registered Geographical Indications [7,8].

Traditional scented rice varieties are thermo-photosensitive, long duration and very low yielding due to their poor harvest index and tendency for lodging. Hence these rice genotypes need to be improved by combining the grain quality attributes with high yield potential. Yield being a complex trait, the knowledge about genetic parameters and correlation between different yield attributing and quality characters is essential while formulating an efficient breeding program. Therefore, keeping the above in view, the current study aims to screen the scented rice genotypes to study the genetic variability and determine the genetic parameters among the selected genotypes. Biometric tools such as genotypic and phenotypic coefficients of variability, heritability and genetic advance are essential

for measuring genetic divergence among genotypes and can be used to improve the scented rice genotypes through suitable breeding methods.

2. MATERIAL AND METHODS

The present investigation to study genetic variability, heritability and genetic gain was conducted at the Regional Agricultural Research Station, Kerala Agricultural University, Ambalavayal, Wayanad district, Kerala in a randomized block design (RBD) with three replications in a plot size of (3 X 2) m² with (20 x 10) cm² spacing with the recommended package of practices. The study consists of nine scented rice genotypes *Gandhakasala* types (GT 1 – GT 8), *Jeerakasala* type (JT 9) and a non-scented local check variety *Aiswarya* collected from different parts of Wayanad district in Kerala. Observations were recorded on twenty yield and associated quantitative, cooking and biochemical characters from ten randomly selected plants of each genotype in each replication and the mean was used for the statistical analysis. The characters that were evaluated include seedling height (cm), leaf length (cm), leaf width (cm), culm number, culm length (cm), days to 50 per cent flowering, number of productive tillers, panicle length (cm), number of spikelets panicle⁻¹, days to maturity, straw yield (kg/ha), grain yield (kg/ha), test weight (g), grain length (mm), grain width (mm), milling recovery (%), kernel elongation ratio, volume expansion ratio, total carbohydrate (%), amylose content (%) and protein content (%).

Analysis of Variance was done using MSTAT package and treatments were compared using Duncan's Multiple Range Test (DMRT). The variance components were estimated using the method pioneered by Lush [9]. The phenotypic coefficient of variation (PCV) and genotypic coefficient of variation (GCV) were calculated by the formula suggested by Burton [10] and classified as suggested by Sivasubramanian and Madhavamenon [11]. Heritability (h^2) in broad sense was estimated by the formula suggested by Lush. The expected genetic advance (GA) under selection among the various characters were calculated according to the formula suggested by Johnson et al. [12].

3. RESULTS AND DISCUSSION

Significant difference has been revealed through ANOVA among the ten genotypes for all characters studied indicating significant inherent genetic differences (Table 1). These differences can be utilized for the genetic improvement of the scented rice genotypes for yield and quality traits through selection. The mean sum of squares for the genotypes showed high significant differences for all the traits under study except for leaf width and similar results has been reported earlier [13].

3.1 Genetic variability

In the present investigation, phenotypic coefficients of variation (PCV) were higher than genotypic coefficients of variation (GCV) for all the traits studied, indicating the impact of the environment on the appearance of these traits (Table 2). Similar findings were reported earlier [14,15]. The results of genetic variability indicated that, the highest GCV and PCV for GY (40.42% and 42.19%) followed by NSP (28.08% and 31.36%) and TW (26.13% and 26.23%), suggesting that these traits were controlled mainly by genetic factors and are less affected by environment and hence these traits can be utilized for further improvement of scented rice genotypes through selection. The estimates were found to be in accordance with Hasib and Kole [16] for TW and GY; Dinkar et al. [17] for GY, NSP and TW. Moderate estimates of both GCV and PCV were exhibited by CL (17.32% and 17.88%), LL (17.29% and 17.97%), SY (17.02% and 19.59%), GW (12.81% and 14.83%), PC (12.70% and 12.71%), GL (12.64% and 12.89%), AC (12.51% and 12.51%), PL (12.25% and 13.98%), NPT (12.13% and 13.40%), CN (11.72% and 12.61%) and VER (10.96% and 11.28%). SH (5.27% and 8.07%), MR (6.09% and 6.10%), (DF (8.75% and 8.77%), DM (8.87% and 8.88%) and TC (9.80% and 9.80%) had the lowest GCV and PCV exhibiting the influence of the environment and the narrow genetic base for these characters. The possibility of improving these characters is less through selection and can be made through combination breeding methods or induced mutagenesis to widen the genetic base followed by selection in advanced generations.

3.2. Heritability and genetic advance

h^2 determines the extent to which a trait is inherited or the value that is transferred from an individual to its progeny [18]. High estimates of h^2 were witnessed for LL (93.35%), CN (88.17%), CL (94.40%), CD (82.44%), DF (99.63%), NPT (84.13%), PL (78.77%), NSP (81.88%), DM (99.79%), GY (92.59%), SY (78.02%), TW (99.29%), GL (95.97%), GW (75.82%), MR (99.94%), KER (63.77%), VER (94.97%), TC (99.97%), AC (99.99%) and PC (99.99%). GA is a measure that explains the degree of gain that may be obtained in a trait under a particular selection pressure. Estimates of high heritability with high genetic advance are more reliable for selection [19]. The highest value of GA was observed for GY (80.25%) followed by TW (53.61%) NSP (52.545), CL (34.71%), LL (34.46%), SY (31.24%), PC (26.12%), AC (25.78%), GL (25.53%), NPT (23.08%), GW (22.96%), CN (22.59%), PL (22.53%), VER (22.03%) and TC (20.18%) which stated the presence of the additive gene action while moderate GA was detected for DM (18.26%), DF (17.99%), KER (16.27%) and MR (12.54%), which revealed non-additive gene action. LW had the lowest GA (4.07%) followed by SH (7.59%). High h^2 coupled with high genetic gain was observed for LL, CN, CL, NPT, PL, NSP, GY, SY, TW, GL, GW, VER, TC, AC and PC. High heritability coupled with high genetic gain indicates that the heritability of these characters was due to additive gene action and the negligible influence of environment in the expression of traits and hence selection based on these characters would be highly effective for the development of elite genotypes. These results conform with Pratap et al. [20], Gupta, et al. [21] and Krishnan et al. [22]. Moderated GA with high h^2 was detected for DM, MR and KER, indicating the scope of individual plant selection for further improvement.

Table 1: Analysis of variance for grain yield and associated quantitative, cooking and biochemical characters in scented rice genotypes

Source of variation (Mean sum of squares)			
Characters	Genotype	Replication	Error
SH	8.6*	2.02	2.68
LL	170.74**	12.24	4.40
LW	0.012	0.02	0.01
CN	2.37**	0.11	0.12
CL	738.18**	22.79	15.91
DF	415.29**	0.63	0.56
NPT	2.51**	0.22	0.17
PL	28.99**	8.19	2.66
NSP	5986.60**	89.37	456.97
DM	725.43**	0.63	0.54
SY	1479466.24**	146855.00	141113.06
GY	1390348.34**	21169.67	40096.72
TW	50.13**	0.09	0.13
GL	2.11**	0.03	0.03
GW	0.23**	0.01	0.02

MR	49.49**	0.01	0.01
KER	0.16**	0.02	0.03
VER	1.04**	0.11*	0.02
TC	177.50**	0.06	0.02
AC	19.27**	0.002	0.001
PC	3.13**	0.00001	0.0001

* Significant at 5% level, ** Significant at 1% level of probability

SH: Seedling height; LL: Leaf length; LW: Leaf width; CN: Culm number; CL: Culm length; DF: Days to 50 per cent flowering; NPT: Number of productive tillers; PL: Panicle length; NSP: Number of spikelets panicle⁻¹ DM: Days to maturity; SY: Straw yield; GY: Grain yield; TW: test weight; GL: Grain length; GW: grain width; MR: Milling recovery %; KER: Kernel elongation ratio; VER: Volume expansion ratio; TC: Total carbohydrate; AC: Amylose content; PC: Protein content

Table 2: Estimates of variability parameters of different characters in scented rice genotypes

Characters	Range	Mean±S.E	PCV%	GCV%	h ² %	GA %
SH	25.18 - 30.8	27.11 ± 1.34	8.01	5.27	46.92	7.59
LL	24.24 - 52.33	43.06 ± 1.71	17.97	17.29	93.35	34.46
LW	1.01 - 1.24	1.08 ± 0.09	11.26	4.14	17.86	4.07
CN	6.46 - 9.34	7.39 ± 0.28	12.61	11.72	88.17	22.59
CL	48.05 - 101.56	89.58 ± 3.26	17.88	17.32	94.40	34.71
DF	101 - 140.67	134.37 ± 0.61	8.77	8.75	99.63	17.99
NPT	6.34 - 9.27	7.28 ± 0.338	13.40	12.13	84.13	23.08
PL	16.38 - 27.1	24.19 ± 1.33	13.98	12.25	78.77	22.53
NSP	64.07 - 216.13	152.92 ± 17.45	31.36	28.08	81.88	52.54
DM	179.0 - 180.33	175.17 ± 0.61	8.88	8.87	99.79	18.26
SY	2923.98 - 5043.73	3910.83±306.72	19.59	17.08	78.02	31.24
GY	1077.87- 3055.33	1659.59±163.49	42.19	40.42	92.59	80.25
TW	12.74 - 25.81	15.63 ± 0.29	26.23	26.13	99.29	53.61
GL	5.48 - 8.47	6.58 ± 0.14	12.89	12.64	95.97	25.53

GW	1.69 -2.39	2.05 ± 0.13	14.83	12.81	75.82	22.96
MR	61.81 – 72.12	66.65 ± 0.08	6.10	6.09	99.94	12.54
KER	1.71 – 2.49	2.12 ± 0.14	12.52	9.81	63.77	16.27
VER	5.21 – 6.07	5.54 ± 0.08	11.28	10.96	94.97	22.03
TC	64.89 – 87.09	78.48 ± 0.12	9.80	9.80	99.97	20.18
AC	17.82 – 26.36	20.25 ± 0.03	12.51	12.51	99.9	25.78
PC	6.38 – 9.70	8.04 ± 0.012	12.71	12.70	99.9	26.12

Mean±S.E: Means±Standard error; PCV %: Percent phenotypic coefficient of variation; GCV %: Percent genotypic coefficient of variation; h^2 : Broad sense heritability; GA%: Genetic Advance as percent of mean.

4. CONCLUSION

The scented rice genotypes differ significantly for different yield attributing and cooking quality characters under this study. *Jeerakasala* types recorded significantly higher yield (2735.22 kg/ha), grain length and test weight than *Gandhakasala* (1350.64 kg/ha) types. The scented rice genotypes had less milling recovery than check variety indicating the need for specially designed milling machines for maximum milling recovery. *Gandhakasala* types showed high kernel elongation ratio and low volume expansion ratio which gives finer appearance compared to *Jeerakasala* type and check variety. Total carbohydrate content was low for scented genotypes than the check variety and can be recommended for low carbohydrate diets. Based on genetic parameters, characters like grain yield, number of spikelets panicle⁻¹ and test weight exhibited high GCV, PCV and high heritability coupled with high genetic advance indicating little influence of environment on the expression of these characters and responsiveness under selection to improve traits to increase the grain yield and quality parameters. Among scented genotypes, JT 9, GT 3 and GT 8 showed better performance based on the high grain yield, high straw yield, intermediate amylose content and moderate aroma which are the preferred qualities for specialty rice. These genotypes could be used for the development of high yielding scented rice varieties with better cooking and biochemical characters. Also, superior genotypes identified can be further evaluated and used for combination breeding. Hence, we suggest to conduct further studies on these genotypes to validate the results and use for varietal development.

REFERENCES

1. Sarika M, Asif S, Renuka N, Kantilal W, Narendra J, Ratnakar T, Altafhusain N. Characterisation of aromatic rice (*Oryza sativa* L.) germplasm and correlation between their agronomic and quality traits. *Euphytica*. 2011;179:237–246.
2. Singh RK, Gautam PL, Sanjeev SS, Singh S. Scented rice germplasm: conservation, evaluation and utilization. In: Singh RK, Singh US, Khush GS, (Eds.). *Aromatic rices*. 1st ed. New Delhi: Oxford & IBH Publishing Co. Pvt Ltd; 2000.
3. Ahuja U, Subhash CA, Thakrar R, Rani NS. Scented Rices of India. *Asian Agri-History*. 2008;12(4):267-283.
4. Latha M, Abdul NM, Abraham Z, Joseph JK, Asokan NR, Mani S, Dutta M. Rice landraces of Kerala State of India: A documentation. *Int. J. Biodivers. Conserv.* 2013;5(4):250-263.
5. Girigan G and Manjula M. Speciality rice biodiversity of Kerala: Need for incentivising conservation in the era of changing climate. *Curr. Sci.* 2018;114(5):997-1006.

6. Mathure S, Shaikh A, Renuka N, Wakte K, Jawali N, Thengane R, Nadaf A. Characterisation of aromatic rice (*Oryza sativa* L.) germplasm and correlation between their agronomic and quality traits. *Euphytica*. 2011;179:237–246.
7. Elsy CR and Adheena RA. Geographical Indications - A marketing tool for unique goods from specific environments. In: Sabu T, Vinod TR, Subramonia IM, Bhaskaran C, Babu A. (Eds.). *Proc. of Kerala Environ. Cong. Thiruvananthapuram: Centre for Environment and Development*; 2012.
8. Blakeney M, Krishnankutty J, Raju RK, Siddique KHM. Agricultural Innovation and the Protection of Traditional rice varieties: Kerala a case study. *Front. Sustain. Food Syst.* 2020; 3:116.
9. Lush JL. Intra-sire correlation and regression of offspring on dams as a method of estimating heritability of characters. *Proc. Am. Soc. Anim. Prod.* 1940;33:293-301.
10. Burton GW. Quantitative inheritance in grasses. In: *Proc. Sixth Int. Grassl. Cong. Pennsylvania, USA, 1952*;1: 227-283.
11. Sivasubramanian V and Madhavamenon P. Path analysis for yield and yield components of rice. *Madras Agric. J.* 1973;60:1217-1221.
12. Johnson HW, Robinson HF and Comstock RE. Estimates of genetic and environmental variability in soybean. *Agron. J.* 1955;47:314-318.
13. Manjunatha GA, Elsy CR, Rajendran P, Jiji J, Rose MF and Krishnan S. DUS Characterization of rice (*Oryza sativa* L.) landraces of Wayanad, Kerala. *Electron. J. Plant Breed.*, 2018; 9(2):617-630.
14. Kishore NS, Srinivas T, Nagabhushanam U, Pallavi M, Sameera SK. Genetic variability, correlation and path analysis for yield and yield components in promising rice (*Oryza sativa* L.) genotypes. *SAARC J. Agri.*,2015;13(1):99-108.
15. Mamta S, Kumar K and Singh, RP. Study of co-efficient of variation, heritability and genetic advance in hybrid rice. *Oryza*. 2007;44(1):160-162.
16. Hasib, KM and Kole PC. Genetic variability for panicle characters in segregating populations of aromatic rice. *Madras Agric. J.* 2008;95: 174-178.
17. Dinkar, Anand K, Ravi RK, Mankesh K, Singh SP, Satyendra and Namrata D. Genetic variability, correlation and path analysis for selection in elite breeding materials of aromatic rice (*Oryza sativa* L.) *Pharma Innovation*. 2023;12(3):5733-5740.
18. Acquaah, G. *Breeding Genetics and Biotechnology*. In: Thomas, B, Murray, BG, Murphy, DJ. (Eds.). *Encyclopedia of Applied Plant Science*. 2nd ed. New York: Elsevier Inc.; 2017.
19. Ogunniyan, J and Olakojo SA. Genetic variation, heritability, genetic advance and agronomic character association of yellow elite inbred lines of maize (*Zea mays* L.). *Niger. J. Genet.* 2014; 28:24-28.
20. Pratap A, Prashant B, Bapsila L, Sandhya, Singh PK. Assessment of genetic variability for yield and yield components in rice (*Oryza sativa* L.) germplasms. *Int. J. bio-resour. stress manag.* 2018; 9(1):087-092.
21. Sneha G, Sameer U, Ganesh KK, Sanket RR, Prashant B, Bapsila L, Pawan KS, Brajesh S. Trait association and path analysis studies of yield attributing traits in rice (*Oryza sativa* L.) germplasm. *Int. J. bio-resour. stress manag.* 2020;11(6):508-517.

22. Krishnan V, Sivaranjani V, Tamilzharasi M, Anandhan T. Characterization of morpho-phenological traits in the traditional landraces of rice. *Electron. J. Plant Breed.* 2023;14(1):234 – 245.

UNDER PEER REVIEW