

**INFLUENCE OF ORGANIC MANURES ON GROWTH, FLOWERING,
QUALITY AND YIELD OF CARNATION
(*Dianthus caryophyllus*) CV. MASTER UNDER
NATURALLY VENTILATED POLYHOUSE CONDITIONS OF
PRAYAGRAJ**

Comment [u1]: Rephrase please. Much words in topic may be confusing

ABSTRACT

An experiment entitled Influence of organic manures on growth, flowering, quality and yield of carnation (*Dianthus caryophyllus*) cv. Master under naturally ventilated polyhouse conditions of Prayagraj was conducted in naturally ventilated polyhouse of Department of Horticulture, Naini Agricultural Institute, SHUATS, Prayagraj, during November, 2022 to April, 2023 with an aim to identify the most suitable organic manure combination under the agroclimatic conditions of Prayagraj. There were thirteen treatments replicated thrice in Randomized Block Design (RBD). It was observed that treatment T₁₁(RDN + FYM 2.5kg/m² + Vermicompost 1.25kg/m² + Neem cake 1kg/m²) was significantly better in all the parameters studied like plant height (48.5cm), number of primary branches(7.6), days taken to first bud initiation (93.5), flower diameter (6.3 cm), flower stalk length (77.2 cm), vase life (6.2 days), number of cut flowers per plant (5.6), number of cut flowers per 200 square meter(27,833.3).

Keywords: Carnation, Organic manures, FYM, Vermicompost, Neem Cake, Trichoderma

1. INTRODUCTION:-

Carnation (*Dianthus caryophyllus*L.) belongs to the Caryophyllaceae family and holds a significant position as one of the world's most popular cut flower crops, consistently ranking among the top ten cut flowers in terms of demand and commercial value. It is a half-hardy perennial cultivated in various regions across the globe, believed to be native to the Mediterranean. It is characterized by its diploid chromosome number of $2n = 30$. They have a sturdy structure and can endure long-distance transportation without significant damage or loss of quality, making them an excellent choice for international and long-haul markets. Exceptional

ability to rehydrate after continuous shipment. Carnations have a remarkable ability to recover and rehydrate quickly after being transported, ensuring they remain fresh and presentable upon arrival at their destination (Bhatia et al., 2007). Excessive use of inorganic fertilizers for increasing the cut flower production has led to the leaching of nutrients causing soil hazards, altering the soil fertility and leading to pollution of soil and water bodies. They imposed disturbance in the ecosystem. Chemical fertilizers are commonly employed in carnation cultivation due to their ability to rapidly release essential elements to the crop. However, their usage has been associated with certain negative effects on flower quality, as well as adverse impacts on soil health, water, and the environment. Organic manure is characterized as the product resulting from the controlled biological decomposition of organic matter. Organic matter has significant effect on the physico-chemical and biological characteristics of soil. Organic manures increase the organic matter in the soil which in turn releases the plant nutrients in easily available form for the use of crops. Application of the organic manures and microbial agents make easy uptake of nutrients when required by the crop contrasting with chemical fertilizers (Vanilarasu and Balakrishnamurthy, 2014). Organic manures like FYM, Neem cake, Vermicompost etc. has great potential to boost the yield and can play an important role in enhancing flower yield, maintain soil health and also to sustain productivity. Neem Cake is highly valued for its diverse composition of micro and macro nutrients. These nutrients play a crucial role in supporting plant growth and development. Its organic nature makes it an environmentally friendly choice for sustainable agriculture practices. One of the significant benefits of neem cake is its ability to control soil-borne pathogens and nematodes. Vermicompost is the end product of the decomposition process involving various species of worms, such as red wigglers, white worms, and earthworms. These worms work to break down a combination of decomposing vegetable or food waste, bedding material, and vermicast. The process, known as vermicomposting, results in nutrient-rich organic matter suitable for enhancing soil fertility and plant growth. It has more nutrient contents and the availability of growth stimulating substances, in addition, many helpful microorganisms which assist in activities like nitrogen fixation, phosphorus solubilization and reduction of harmful micro-organisms (Barik and Barik 2009). FYM is an important component of traditional farming systems and organic agriculture. It has been used for centuries as a natural fertilizer and soil conditioner. Its primary purpose is to enhance soil health, promote plant growth and sustainably manage agricultural land.

Comment [u2]: Not clear please

Comment [u3]: How?

Comment [u4]: No reference, what essential element

Comment [u5]: rephrase

Comment [u6]: not necessary

Comment [u7]: break this sentence

Comment [u8]: break sentence

Comment [u9]: you can't start a sentence this way

Trichoderma has been widely studied and utilized as a biocontrol agent against plant pathogens, including fungi, in various agricultural systems. In case of carnation cultivation, *Trichoderma* species have shown promising results as organic fungicides. It can be applied as a preventative measure to protect carnations from fungal diseases, such as Fusarium wilt (caused by *Fusarium oxysporum*) and Botrytis blight (caused by *Botrytis cinerea*).

Comment [u10]: How do you intend to link your introduction to material and methods?
Try to improve on your sentence linkage to enhance the beauty of your introduction
Link your objective with your introduction

2. MATERIALS AND METHODS:-

2.1 Geographical location and climatic conditions

Geographically, Prayagraj is situated in the South-Eastern part of Uttar Pradesh. It lies between the parallels of 24°77' and 25°47' north latitudes and 81°19' and 82°21' east longitudes. The area of Prayagraj district comes under agro climatic zone V (Upper Gangetic Plain region) and sub-zone of Central Plains. The climate ranges from dry sub-humid to semi-arid and the soil is alluvium calcareous sandy loam. This District experiences average maximum temperature range between 43°– 47°C which may go as high as 48°C during peak summers (May-June). The minimum average temperature is 2-4°C, which may fall as low as 1°C during peak winter months (December-January). The average rainfall of the district is 1042 mm and the monsoon season is spread between July-September.

2.2 Experimental details

The experiment was conducted in Randomized Block Design (RBD) with 13 treatments of organic manures with three replications in the Departmental Research Field of Department of Horticulture, Sam Higginbottom University of Agriculture, Technology And Sciences, Prayagraj during November, 2022 to April, 2023. Total number of treatments were thirteen viz. T₀- Control RDN (30:20:10 g/m² NPK), T₁ (RDN + FYM 5kg/m²), T₂ (RDN + Vermicompost 2.5 kg/m²), T₃ (RDN + Trichoderma 10g/m²), T₄ (RDN + Neem cake 1kg/m²), T₅ (RDN + FYM 2.5 kg/ m² + Vermicompost 1.25 kg/m²), T₆ (RDN + FYM 5 kg/m² + Trichoderma 10g/m²), T₇ (RDN + FYM 5kg/m² + Neem cake 1kg/m²), T₈ (RDN + Vermicompost 2.5kg/m² + Trichoderma 10g/m²), T₉ (RDN + Vermicompost 2.5kg/m² + Neem cake 1kg/m²), T₁₀ (RDN + FYM 2.5kg/m² + Vermicompost 1.25kg/m² + Trichoderma 10g/m²), T₁₁ (RDN + FYM 2.5kg/m² + Vermicompost 1.25kg/m² + Neem cake 1kg/m²), T₁₂ (RDN + FYM 2.5kg/m² + Vermicompost 1.25kg/m² + Trichoderma 10g/m² + Neem cake 1kg/m²). Carnation cultivar Master was planted on 16th November, 2022 at a spacing of 20 cm x 20 cm.

Comment [u11]: What is the initial state of the soil before adding the treatment?

3. RESULTS AND DISCUSSION

Plant height and Number of primary branches

Maximum plant height (48.5 cm) and number of primary branches was observed from the plants grown in treatment T₁₁ containing RDN + FYM 2.5 kg/m² + Vermicompost 1.25 kg/m² + Neem cake 1 kg/m² significantly superior to rest of the treatments. The treatment combination of RDN (Recommended Dose of Nutrients) + FYM (Farm Yard Manure) + Vermicompost + Neem cake has shown significantly better plant height in carnations compared to other treatment combinations. This can be attributed to the synergistic effects of the various components. RDN provides essential nutrients in optimal proportions, promoting overall plant growth. FYM enhances soil structure and nutrient retention, leading to improved root development. Vermicompost enriches the soil with organic matter and beneficial microorganisms, facilitating nutrient uptake. Neem cake acts as a natural pesticide and also provides supplementary nutrients. Together, these components create a favorable environment for root expansion and nutrient absorption, resulting in taller, healthier carnation plants. The combination's balanced nutrient supply and disease-resistant properties contribute to the superior plant height observed. Similar findings were reported by **Barman et al., (2005)**; **Pooja et al., (2012)**; **Bohra et al., (2019)**.

Days taken to first flower bud initiation

In terms of days taken to first bud initiation treatment T₁₁ containing RDN + FYM 2.5 kg/m² + Vermicompost 1.25 kg/m² + Neem cake 1 kg/m² recorded minimum days taken to first bud initiation (93.5) whereas the maximum days were recorded in the treatment T₀ – Control (116.4). The earliness might be due to the effect of organic fertilizers, creating a conducive source sink relationship and ultimately causing an increase in the synthesis of cytokine in the root tissue and its simultaneous transport to auxiliary buds would have resulted in better sink for mobilization of photo assimilates at a rapid rate and have helped in the early transformation from vegetative to reproductive phase. Similar results were observed by **Gupta et al., (2014)** in marigold; **Bohra et al., (2019)**; **Mogalet al., 2006**.

Comment [u12]: Keep sentence short. Earliness?

Flower diameter (cm)

Data revealed that higher flower diameter was recorded in treatment T₁₁ containing RDN + FYM

2.5kg/m² + Vermicompost 1.25kg/m² + Neem cake 1kg/m² (5.3cm) whereas the minimum flower diameter was recorded in the treatment T₀- Control (4.2cm). This increase in flower diameter might be due to better nutrient uptake, higher photosynthesis and excellent physiological, biological activities due to presence of organic manures which have resulted in rapid synthesis and translocation of photosynthetic from the source to developing flower bud and finally increase in flower diameter **Pooja et al., (2012); Swaroop and Janakiram, (2010); Pandey et al., (2017).**

Stalk length (cm)

Maximum flower stalk length was recorded in treatment T₁₁ containing RDN + FYM 2.5kg/m² + Vermicompost 1.25kg/m² + Neem cake 1kg/m² significantly superior to rest of the treatments.

All the treatments involving in combination with various levels of phosphorus were effective considerably as compared to control due to enhanced absorption in organic manures inoculated plants, leading to increased availability of assimilates that needed for the improvement in flower stalk length. **Bohra et al., (2019); Gupta et al., (2008)** reported similar results.

Comment [u13]: Rephrase please

Vase life (Number of days)

In terms of vase life treatment T₁₁ containing RDN + FYM 2.5kg/m² + Vermicompost 1.25kg/m² + Neem cake 1kg/m² recorded maximum vase life (6.2 days) whereas the minimum vase life was recorded in the treatment T₀- Control (3.7 days). Organic fertilizer increased vase life of carnation. The Increased vase life might be due to by triggering of such metabolic activity and Narrowing of the C:N ratio by the significant Accumulation of carbohydrates. Furthermore, the use of the treatment combination RDN (Recommended Dose of Nutrients) + FYM (Farm Yard Manure) + Vermicompost + Neem cake has resulted in an improved vase life of carnation flowers compared to other treatment combinations. This improvement can be attributed to the combined effects of these components on flower quality and longevity. RDN supplies essential nutrients in optimal proportions, ensuring better flower health and resilience. FYM enhances soil structure and nutrient retention, allowing the flowers to access nutrients even after being cut. Vermicompost enriches the soil with organic matter and beneficial microorganisms, contributing to the flowers' overall vitality and vase life extension. Neem cake's presence helps in reducing post-harvest pests, minimizing petal damage and decay. The synergy of these components creates a favorable environment for cut flowers, resulting in longer-lasting and aesthetically pleasing carnations. These findings are well supported by **Pooja et al., (2012); Bohra et al., (2019); Kumar et al., (2022)** in tuberose.

Comment [u14]: Please sentences properly

Comment [u15]: were

Comment [u16]: remove

Number of cut flowers per plant/200m²

Yield is an important parameter to decide the efficacy of a treatment. Data recorded on maximum number of flowerstalks per plant (5.6) and maximum number of flower stalks per 200m² (27,833) in treatment T₁₁ containing RDN + FYM 2.5kg/m² + Vermicompost 1.25kg/m² + Neem cake 1kg/m². The treatment combination of RDN (Recommended Dose of Nutrients) + FYM (Farm Yard Manure) + Vermicompost + Neem cake has led to a higher number of cut flowers per plant/200m² in carnations compared to other treatment combinations. This increase can be attributed to the combined effects of these components on plant growth and reproductive development. RDN supplies essential nutrients in optimal proportions, promoting robust flower bud initiation and development. FYM enhances soil fertility and water retention, ensuring continuous nutrient availability to support flower production. Vermicompost enriches the soil with organic matter and beneficial microorganisms, promoting healthy flowering and increased flower bud formation. Neem cake's presence provides natural pest resistance, preventing potential flower damage and loss. The synergy of these components creates an ideal growth environment, resulting in more abundant flower production in carnations, leading to higher yields of cut flowers per plant and per square meter. Findings were in accordance with findings of **Raha et al., (2015); Bohra et al., (2019).**

Table 1. Effect of organic manure on plant height, number of primary branches, days taken to first flower bud initiation and flower diameter of carnation

Treatment Symbols	Treatment Combinations	Plant height (cm)	No. of primary branches	Days taken to first flower bud initiation	Flower diameter (cm)
		60 Days*	60 Days*		
T ₀	Control RDN (30:20:10 g/m ² NPK)	31.9	4.4	116.4	4.2
T ₁	RDN + FYM 5kg/m ²	45.1	6.1	114.0	5.1
T ₂	RDN + Vermicompost 2.5 kg/m ²	34.1	6.2	100.9	4.9
T ₃	RDN + Trichoderma 10g/m ²	40.5	5.9	102.9	4.8
T ₄	RDN + Neem cake 1kg/m ²	34.6	6.2	102.4	4.3
T ₅	RDN + FYM 2.5 kg/m ² + Vermicompost 1.25 kg/m ²	39.5	4.5	111.4	5.3
T ₆	RDN + FYM 5 kg/m ² + Trichoderma 10g/m ²	36.4	6.4	101.5	4.9
T ₇	RDN + FYM 5kg/m ² + Neem cake 1kg/m ²	37.5	4.8	101.3	5.0
T ₈	RDN + Vermicompost 2.5kg/m ² + Trichoderma 10g/m ²	35.9	4.7	107.4	5.3
T ₉	RDN + Vermicompost 2.5kg/m ² + Neem cake 1kg/m ²	46.5	5.8	99	6.0
T ₁₀	RDN + FYM 2.5kg/m ² + Vermicompost 1.25kg/m ² + Trichoderma 10g/m ²	35.5	5.1	115.7	5.5
T ₁₁	RDN + FYM 2.5kg/m ² + Vermicompost 1.25kg/m ² + Neem cake 1kg/m ²	48.5	7.6	93.5	6.3
T ₁₂	RDN + FYM 2.5kg/m ² + Vermicompost 1.25kg/m ² + Trichoderma 10g/m ² + Neem cake 1kg/m ²	46.1	6.3	110.0	4.7
F-Test		S	S	S	S
SEd (±)		0.75	0.46	0.34	0.11

CV	2.33	9.80	0.55	2.64
CD _{0.05}	1.54	0.94	0.98	0.23

*Daysafterpinching

Table 2: Effects of organic manures on stalk length, vase life of flowers, no. of flowers per plant and number of flowers per 200m² of carnation

Treatment Symbols	Treatment Combinations	Stalk Length (cm)	Vase Life (days)	Number of flower stalks per plant	Number of flower stalks per 200m ²
T ₀	Control RDN (30:20:10 g/m ² NPK)	61.0	3.7	4.2	21000.0
T ₁	RDN + FYM 5kg/m ²	68.0	4.8	5.0	25166.7
T ₂	RDN + Vermicompost 2.5 kg/m ²	65.7	5.2	5.1	25333.3
T ₃	RDN + Trichoderma 10g/m ²	68.4	4.9	4.9	24500.0
T ₄	RDN + Neem cake 1kg/m ²	67.9	5.1	4.5	22666.7
T ₅	RDN + FYM 2.5 kg/ m ² + Vermicompost 1.25 kg/m ²	68.1	4.9	4.9	24666.7
T ₆	RDN + FYM 5 kg/m ² + Trichoderma 10g/m ²	64.9	4.8	4.8	23833.3
T ₇	RDN + FYM 5kg/m ² + Neem cake 1kg/m ²	66.8	4.8	4.8	23833.3
T ₈	RDN + Vermicompost 2.5kg/m ² + Trichoderma 10g/m ²	70.6	4.5	4.4	22166.7
T ₉	RDN + Vermicompost 2.5kg/m ² + Neem cake 1kg/m ²	73.9	5.3	5.3	26333.3
T ₁₀	RDN + FYM 2.5kg/m ² + Vermicompost 1.25kg/m ² +Trichoderma 10g/m ²	66.3	4.9	4.9	24500.0
T ₁₁	RDN + FYM 2.5kg/m ² + Vermicompost 1.25kg/m ² + Neem cake 1kg/m ²	77.2	6.2	5.6	27833.3

T₁₂	RDN + FYM 2.5kg/m ² + Vermicompost 1.25kg/m ² + Trichoderma 10g/m ² + Neem cake 1kg/m ²	64.5	5.1	5.1	25500.0
F-Test		S	S	S	S
SEd (±)		0.69	0.24	0.13	660.76
CV		1.24	5.94	3.32	3.32
CD_{0.05}		1.42	0.49	0.27	1363.16

UNDER PEER REVIEW

Conclusion:-

Based on the present investigation it is concluded that treatment T₁₁ (RDN + FYM 2.5kg/m² + Vermicompost 1.25kg/m² + Neem cake 1kg/m²) was found to be best in all the parameters like plant height, number of primary branches, days taken to first bud initiation, flower diameter, flower stalk length, vase life, number of cut flowers per plant, number of cut flowers per 200 square meter followed by T₉ (RDN + Vermicompost 2.5kg/m² + Neem cake 1kg/m²), whereas minimum is recorded in treatment T₀ (Control).

Comment [u17]: improve on this conclusion

Comment [u18]: A research paper is different from report. Improve on this write up.

References:-

Barik, T. and Barik, K. C. (2009). Vermicomposting practices for its quality maintenance. *Indian farming*, **58** (12): 22-25.

Barman, D., Rajni, K. and Upadhyaya, R.C. (2005). Corm multiplication of gladiolus as influenced by application of potassium and spike removal. *Journal of Ornamental Horticulture*, **8**(2):104-107.

Bhatia, S., Gupta, Y.C. and Dhiman, S.R., (2007). Effect of growing media and fertilizers on growth and flowering of Carnation under protected condition. *Journal of Ornamental Horticulture* **8**(3): 239-240.

Bohra, M., Rana, A., Punetha, P., Upadhyay, S. and Nautiyal, B.P. (2019). Effect of organic manures and biofertilizers on growth and of Karnini China aster. *Indian Journal of Horticulture* **76**(2): 329-333.

Gupta, P., Rajwal, N., Dhaka, V.K. and Rajwal, D., (2008). Effect of different levels of vermicompost, NPK and FYM on performance of Gladiolus (*Gladiolus grandiflorus* L) cv. Happy End. *The Asian Journal of Horticulture*, **3**(1): 142-143.

Gupta, R., Yadav, A., & Garg, V. K. (2014). Influence of vermicompost application in potting

media on growth and flowering of marigold crop. *International Journal of Recycling of Organic Waste in Agriculture*, **3**, 47.

Kumar, P., Sheoran, S., & Beniwal, B. S. (2022). Growth and yield parameters of rose as influenced by different organic manures and their levels. *The Pharma Innovation Journal*, SP-11(6), 394-398

Mogal, S.A., Khiratkar, S.D., Chopde, N.K., Dalvi, A.M., Kuchanwar, O.D., Khobragade, Y.R. (2006). Effect of organic manures and biofertilizers with reduced doses of nitrogen on growth, yield and quality of China aster. *Journal of Soils and Crops* **16**(1): 180-185.

Pandey, S.K., Kumari, S., Singh, D., Singh, V.K. and Prasad, V. M. (2017). Effect of Biofertilizers and Organic Manures on Plant Growth, Flowering and Tuber Production of Dahlin (*Dahlia variabilis* L.) Cv. SP. Kamala, *Indian Journal of Pure & Applied Biosciences*, **5**(2): 549-555.

Pooja G., and Kumari, S., Dikshit, S.N. (2012). Response of African marigold (*Tagetes erecta* L.) to integrated nutrient management. *Annals of Biology*, **28**(1):66-67.

Raha, S. (2015). Studies on the Effect of Vermicompost on the Growth Yield and Quality of Chrysanthemum (*Chrysanthemum coronarium* L.) cv. Kasturba Gandhi. *International Journal of Environmental Sciences*, **4**(2), ISSN: 2277-1948.

Swaroop, k and Janakiram, T. (2010). Response of Bio fertilizers, FYM and their combined application with different levels of inorganic fertilizers in gladiolus (*Gladiolus hybridus* L.) *International Journal of Tropical Agriculture* **28**(3-4): 435-438.

Vanilarasu, K and Balakrishnamurthy, G., (2014). Influences of organic manures and amendments in soil physiochemical properties and their impact on growth, yield and nutrient uptake of banana. *The Bioscan*, **9**(2): 525-529.