

Assessment of Radiological Health Risks in Agricultural Soil Samples within Bitumen Belt of Ondo State, Nigeria

Abstract:

Naturally occurring radionuclides (^{238}U , ^{232}Th and ^{40}K) present in agricultural soil samples collected within the bitumen belt of Ondo State, Nigeria was measured using gamma spectroscopy. The radiological health risk parameters; absorbed dose rate (D), Annual Gonadal Equivalent Dose (AGED), External Hazard Index (H_{ex}), Internal Hazard Index (H_{in}) and Excess Lifetime Cancer Risk (ELCR) were estimated using standard analytical method. The estimated results of D ranged from 3.56 nGyh^{-1} (Omotosho) to 10.55 nGyh^{-1} (Igbotako) with a mean value of $5.83 \pm 2.19 \text{ nGyh}^{-1}$. While AGED ranged from 24.07 mSvy^{-1} (Omotosho) to 70.72 mSvy^{-1} (Igbotako) with a mean value of $40.39 \pm 14.26 \text{ mSvy}^{-1}$. Also, H_{ex} varies between 0.021 (Omotosho) to 0.063 (Igbotako) with a mean value of 0.034 ± 0.01 . Similarly, H_{in} ranged from 0.030 (Omotosho) to 0.089 (Igbotako) with a mean value of 0.047 ± 0.02 . Whereas, estimated ELCR were far lower than the safe limit of 0.29×10^{-3} and implies that though the radiation hazard due to exposure to natural radionuclides is minimal, utilization of the studied soil may not cause any immediate health hazard to the inhabitants but prolonged exposure might pose radiologically induced health challenges.

Keywords: External and Internal Hazard Indices, ELCR, Absorbed Dose Rate, Bitumen, Ondo

1. Introduction

Bitumen occurs naturally and is found typically on the surface and sub-surface as seepages and outcrops, or as tar sands. This implies that they occur in the earth's uppermost layer and within strata below its surface. The tar sand is a combination of clay, sand, heavy oil and water in varying proportions of 2%, 84%, 12%, and 4%, respectively [1]. Generally, bitumen is believed to have originated from fossil deposit, which represents the product of reservoir transformation of conventional oil by microorganism.

Ondo State bitumen belt spans six of the eighteen Local Government Areas: Odigbo, Irele, Okitipupa, Ilaje, Ese-Odo and Idanre [2]. The exploitation sites of bitumen in Nigeria were distributed mainly along a stretch beginning at the outcrop belt northeast of Lekki Lagoon in Ogun State and extend toward the Southeast in Ondo State [3].

Usually, oil exploration is known to have great impact on the environment due the presence of natural radionuclides in crude oil and bitumen [4]. The experience of tar sand and petroleum exploration and exploitation had created oil pollution and spillages affecting land, crop, water and welfare of the host communities. Ordinarily, bitumen pollution occurs when there is seepage from the ground reservoir to the ground surface through cracks or faults on the ground due to changes in temperature and/or bacterial degradation [2]. Spilling of the lower viscous bitumen through seepage on farmlands and rivers is common within Ondo State bitumen belt.

However, the exploration and exploitation of bitumen may bring economic benefits to a country, but its activities may be destructive to the environment even at the safest and best operating practices and such unsafe acts may include the redistribution of Naturally Occurring Radioactive Materials (NORMs) which are within the earth crust and brought to the surface during the processes [5]. In

addition, contamination of soil compartments by bitumen disrupts the activities of soil microorganisms and nutrient availability to plants [6].

The soil, a major sink of environmental contaminants, comprises several organic and mineral components and acts as a repository for many environmental pollutants including radionuclides. This accounts for the presence of some levels of radioactive elements which are primarily dependent on the parent rock type of the soil. The physicochemical properties of soils also influence the behavior, concentration, and distribution of radioactive materials [7]. Lower levels of potassium and thorium are usually associated with sedimentary rocks while higher levels are associated with igneous rocks [8, 9].

Consequently, consumption of radionuclide contaminated food crops can lead to the radiological exposure of a person [10]. More so, radionuclides in soils and other environmental matrices can directly pose significant human exposure especially for local population through various pathways, depending on their concentration. Although radionuclides exposure rate in soil is low but the effect of these radiations overtime can lead to radiological health risk, which can be estimated [11].

Anthropogenically enhanced naturally occurring radionuclides ^{238}U , ^{232}Th and ^{40}K present in soils around tar-sand deposit area of Ogun state were measured by Gbadamosi *et al.*, [12]. Their report showed that the average activity concentrations of ^{238}U , ^{232}Th and ^{40}K ranges from BDL to 76.00 ± 12.00 , BDL to 204.48 ± 13.02 and BDL to $755.6 \pm 40.15 \text{ Bqkg}^{-1}$ respectively. While the values obtained for their associated radiological health hazard parameters were all higher than the world's average set by the United Nations Scientific Committee on Effect of Atomic Radiation (UNSCEAR). However, RESRAD computer code was used to calculate the probability of excess lifetime cancer incurred by dwellers of the tar-sand deposit site, the level of which was determined to be 2.3×10^{-3} . Hence the tar-sand soil samples were considered to pose serious radiological hazard and cancer risk to humans through the multiple routes of exposure.

Isinkaye *et al.*, [4] evaluated radiological hazards due to natural radioactivity in bituminous soils from tar-sand belt of Southwest Nigeria using HpGe Detector. They concluded that the activity contents of soil and viscous bitumen from the tar-sand belt are lower than those in soils of non-bituminous area. Meanwhile, Fasasi *et al.*, [1] studied natural radioactivity of the tar-sand deposits areas of Ondo State, Southwestern Nigeria and their report revealed that the measured activity in the bituminous sand layer is so low that it can be said to be non-radioactive. However, the result of the Energy Dispersive X-Ray Fluorescence (EDXRF) supports the presence of radioelements in the overburden.

It is of concern that there might be a high level of radionuclides presence in the tar-sand deposit and bitumen contaminated agricultural soil within the study area, and future exploration or exploitation can elevate this level hence pose negative radiological impact on the environment and the population. It might expose individuals to both stochastic and deterministic effects through various pathways due to the ionizing radiation emitted by these radionuclides. These effects include; sterility, tumor necrosis and cancer.

However, the renewed interest in the exploration and exploitation of the vast bituminous sand deposit of Nigeria, the suspected radiological health risk and recent reported cases of possible high levels of radioactive materials in some bituminous nodules in Czech Republic [13] and in northern Saskatchewan, Canada [14] account for the need to quantify the presence and level of radionuclides in agricultural soil within the Ondo State bitumen belt.

2. Materials and Methods

2.1 Study Location

The study area is located on the eastern margin of a coastal sedimentary Benin basin which lies on the onshore regions of Eastern Dahomey between the coordinates; longitude $6^{\circ} 15' 0''$ N & $6^{\circ} 45' 0''$ N and latitude $4^{\circ} 30' 0''$ E & $5^{\circ} 10' 0''$ E (Figure 1). It is the most noted area of bitumen activities in Nigeria, and falls within the tropical rainforest region with two distinct climatic seasons, which are; dry season from November to April and wet (rainy) season from May to October. The sedimentary rocks are mainly of the postCretaceous sediments and the Cretaceous Abeokuta Formation. Although exploitation of the bitumen is yet to commence, seepages of the naturally occurring bitumen within the shallow subsurface contaminates soils, farmlands and rivers within the study area, hence constitute another source of radioactivity due to the presence of naturally occurring radioactive materials in the bitumen. The soil samples were collected one each from Okitipupa, Iletitun, Igbotako, Omotosho, Ode-Aye, Agbabu, Ode Irele, Iyasan, Akotogbo, Loda, Ibekegbo, Igbobini and Araromi Sea-Side.




Figure 1: Map of the Study Area

2.2 Samples Collection

Thirteen (13) soil samples were collected from a pre-determined depth of 0.5m – 1.0m [15], with the aid of hand auger across agricultural fields within the study area. The samples were sealed in a transparent polythene bag and carefully labeled to prevent sample mix-up. They were then properly marked and taken to the Nigerian Institute of Radiation Protection and Research (NIRPR), University of Ibadan, Ibadan, Oyo state Nigeria for processing and analysis.

2.3 Sample Preparation

Each soil sample was dried under the laboratory condition until constant weight was achieved. The dried samples were pulverized and homogenized using a motorized grinder and allowed to pass through a sieve of 200 μ m mesh size. The homogenized soil samples were then dried in a temperature-controlled oven at 105°C for about 24 hrs in order to eliminate organic matter content of the soil samples. They were then placed in Marinelli beakers (size 500ml each) and sealed

accordingly to maintain its in-situ characteristics. The weights of the sealed samples were recorded using electronics weighing balance and then kept for twenty-eight (28) days in order to achieve radioactive secular equilibrium between parent radionuclides and their respective daughters.

2.4 Gamma Spectrometry

The radioactivity measurement was done using Sodium Iodide (NaI-TI) scintillation detector. The detector is a lead shield Canberra 76mm x 76mm NaI (TI) crystal model number 802 series. It is a compatible sealed assembly which contains high-resolution NaI (TI) crystal, a photomultiplier tube that detects the small visible light photons produced in the crystal and converts them into amplified electrical pulses, which is fed into analyzer systems through a preamplifier base. The detector system was calibrated before carrying out actual measurement of the soil samples. In order to commence counting, three gamma standard sources Cs-137, Am-241 and Co-60 were placed into 6 cm lead shield of the detector chamber. This set up is aimed to minimize the effects of background and scattered radiation. The energy calibration was done by obtaining the relationship between peak position in the spectrum and the corresponding gamma-ray. The height of each pulse output from a photomultiplier tube which was viewed on the display output and the channel corresponding to it, is directly proportional to the initial gamma energy producing the pulse. The calibration was done using gamma emitter sources of known energies, these are Cs-137 and Co-60 source that emits gamma rays with energies of 662 keV, 1332 keV and 1173 keV, and Am-241 which is an alpha emitter but also emits some gamma rays with energies 26.3 keV and 59.6keV. The gamma emitter sources were exposed to the NaI (TI) detector and gamma spectrum was acquired. These were done with the amplifier gain that gives 72% energy resolution for the 662 keV of Cs-137 and counted for 30 minutes. The net area corresponding to the photopeak's in the energy spectrum was computed by subtracting count from the background source from the total area of the photopeak's. The identification of individual radionuclides was performed using their gamma ray energies and the quantitative analyses of radionuclides were performed using gamma ray spectrum analysis software, Genie 2000.

3. Radiological Health Risk Parameters

The radiological health risks indices are standard parameters used in radiation studies to assess and estimate the effects of radiation exposure on the health of people and the environment. These indices are useful in estimating the radiological effects of samples that contains radionuclides (^{238}U , ^{232}Th and ^{40}K). Some radiation health risks parameters associated with the studied soil samples are discussed below:

3.1 Radium Equivalent Activity Index (Ra_{eq})

The radium equivalent is an index used to describe the gamma output from different mixtures of Uranium (i.e. ^{226}Ra), ^{232}Th and ^{40}K in a material. From the activity concentrations of ^{226}Ra , ^{232}Th and ^{40}K , the radium equivalent concentration was calculated using the equation below [16].

$$Ra_{eq} = A_{Ra} + 1.43A_{Th} + 0.077A_K \quad (1)$$

Where A_{Ra} , A_{Th} and A_K are the activity concentrations of ^{238}U , ^{232}Th and ^{40}K (Bq/kg), respectively. An assumption was made in defining radium equivalent activity, that 370 Bq/kg of ^{226}Ra , 259Bq/kg of ^{232}Th and 4810 Bq/kg of ^{40}K yields the same gamma dose rate [17].

3.2 Absorbed Dose Rate (D)

If a radionuclide activity concentration is known then its exposure dose rate in air at 1m above the ground can be calculated using equation 2 [18].

$$D = 0.429A_{Ra} + 0.666A_{Th} + 0.042A_K \quad (2)$$

Where D is the dose rate in nGyh⁻¹ while A_{Ra}, A_{Th} and A_K are the specific activity concentrations in Bq kg⁻¹ of ²³⁸U, ²³²Th and ⁴⁰K respectively in the soil sample and 0.429, 0.666 and 0.042 (nGyh⁻¹ per Bq kg⁻¹) are the concentration-to-dose conversion factors.

The absorbed dose values are often converted to effective dose equivalent from which the excess life cancer risk is calculated, since the absorbed dose rate itself does not show possible biological effects.

3.3 External Hazard Index (H_{ex})

The concept of external hazard index (H_{ex}) was used to assess the potential health risk associated with humans and gamma radiation emitted by radionuclides. The index was estimated using equation 3 [19, 20, 21].

$$H_{ex} = \frac{A_{Ra}}{370} + \frac{A_{Th}}{259} + \frac{A_K}{4810} \quad (3)$$

Where A_{Ra}, A_{Th} and A_K are the average activity concentrations of ²²⁶Ra, ²³²Th and ⁴⁰K, respectively in Bq/kg as earlier defined. The value of this index must be less than unity for the radiation hazard to be insignificant. The maximum value of H_{ex} equal to unity corresponds to the upper limit of Ra_{eq} (i.e 370 Bq/Kg) [21].

3.4 Internal hazard index (H_{in})

Radon and its short-lived products are radiologically hazardous when inhaled. Internal exposure to radon and its daughter products is very hazardous and can lead to respiratory diseases like asthma and cancer. The internal hazard index (H_{in}) can be used to determine the internal exposure of living cell to radon and its products [10]. These indices were determined using equation 4 [19, 20, 21].

$$H_{in} = \frac{A_{Ra}}{185} + \frac{A_{Th}}{259} + \frac{A_K}{4810} \quad (4)$$

Where A_{Ra}, A_{Th} and A_K, are the average activity concentrations of ²²⁶Ra, ²³²Th and ⁴⁰K, respectively in Bq/kg. H_{in} should be less than unity for the radiation hazard to be insignificant.

3.5 Annual Effective Dose Equivalent

The concentrations of terrestrial gamma radiation from ²³⁸U, ²³²Th and ⁴⁰K in environmental matrix can be used to deduce the annual effective dose equivalent (AEDE) received by an adult in both indoor and outdoor situations from absorbed dose rate in the air, using their respective average conversion coefficients and occupancy factor. The conversion factor value was estimated to be 0.7SvGy⁻¹ for gamma ray exposure in the environment in both indoor and outdoor situations

[22], while the occupancy factor were 0.2 for outdoor considering that people on regular basis spent 20% of their time outdoors. For indoor measurement, the occupancy factor for building materials was estimated to be approximately 0.8. Hence to estimate the Annual Effective Dose Equivalent (AEDE), equations 5 and 6 can be used for outdoor and indoor situations respectively.

$$AEDE(Outdoor) \left(\frac{\mu Sv}{y} \right) = D_{air} \left(\frac{nGy}{h} \right) \times 8760 \left(\frac{h}{y} \right) \times 0.2 \times 0.7 \left(\frac{Sv}{Gy} \right) \times 10^{-3} \quad (5)$$

$$AEDE(Indoor) \left(\frac{\mu Sv}{y} \right) = D_{air} \left(\frac{nGy}{h} \right) \times 8760 \left(\frac{h}{y} \right) \times 0.8 \times 0.7 \left(\frac{Sv}{Gy} \right) \times 10^{-3} \quad (6)$$

The world AEDE for both indoor and outdoor terrestrial gamma radiation is 0.460mSv^{-1} . This index measures the risk of stochastic and deterministic effects in the irradiated individuals exposed [23].

3.6 Annual Gonadal Equivalent Dose (AGED)

Gonads, bone marrow and the bone surface cells are the most sensitive parts of human body to radiation, hence are regarded as organs of interest according to UNSCEAR [24]. It had been known that an increase in AGED will affect the bone marrow and destroy the red blood cells which are then replaced by white blood cells. This situation results in a health condition known as blood cancer (leukemia). AGED for members of the public can be calculated with given activity concentrations of ^{226}Ra , ^{232}Th and ^{40}K (in Bq/Kg) using the relationship below [25]:

$$AGED \left(\frac{mSv}{y} \right) = 3.09A_{Ra} + 4.18A_{Th} + 0.314A_K \quad (7)$$

Where, A_{Ra} , A_{Th} and A_K represent the radioactivity concentration of ^{226}Ra , ^{232}Th and ^{40}K (in Bq/Kg) in soil samples respectively.

3.7 Excess Lifetime Cancer Risk (ELCR)

The ELCR value describes the number of cancers expected in a given number of people on exposure to a carcinogen at a given dose. It is associated with the probability of developing cancer over a lifetime at a given exposure level. An increase in the ELCR causes a proportionate increase in the rate at which an individual can get cancer of the breast, prostate or even blood. Excess Lifetime Cancer Risk (ELCR) is given according to Taskin *et al.*, [26] as:

$$ELCR = AEDE \times \text{Average Duration of Life (DL)} \times \text{Risk Factor (RF)} \quad (8)$$

Where; AEDE is the Annual Effective Dose Equivalent, DL is the average duration of life / life expectancy (estimated as 70 years), and RF is the Risk Factor (Sv^{-1}), i.e., fatal cancer risk per Sievert. For stochastic effects, International Commission on Radiological Protection (ICRP) uses RF as 0.05Sv^{-1} for public [26], with the UNSCEAR standard for ELCR being 0.29×10^{-3} . Substituting the values of DL and RF in equation 8 gives equation 9 for ELCR.

$$ELCR = AEDE \times 70 \times 0.05 \quad (9)$$

4. Results and Discussion

In this study, the activity concentrations of naturally occurring radionuclide ^{40}K , ^{232}Th and ^{238}U in agricultural soil samples were measured using gamma ray spectroscopy method and the result presented in Table 1. These values were used in the assessment of radiological health risks in

agricultural soil samples within the bitumen belt of Ondo state, Nigeria. The activity concentration of ^{40}K ranges from 8.73Bqkg^{-1} (Omotosho) to 93.83Bqkg^{-1} (Araromi Sea-Side) with mean value of $35.85 \pm 23.75\text{Bqkg}^{-1}$. The activity concentrations of ^{238}U in the soil samples have its lowest value as 2.81Bqkg^{-1} (Okitipupa) and highest value as 9.55Bqkg^{-1} (Igbotako), with a mean value of $5.01 \pm 2.08\text{Bqkg}^{-1}$. While the activity concentration of ^{232}Th ranges from 0.35Bqkg^{-1} (Araromi Sea-Side) to 10.49Bqkg^{-1} (Agbabu) with a mean value of $3.37 \pm 2.98\text{Bqkg}^{-1}$. The activity concentrations of ^{40}K , ^{232}Th and ^{238}U obtained in this work were found to fall within the range of that obtained by Gbadamosi *et al.*, [12] in soils around tar-sand deposit area of Ogun state, Nigeria, but lower than that obtained by Olawale *et al*; [27] at Gbeleju-Loda and Ode-Irele in Ondo State Nigeria.

Table 1: Specific activity concentration of radionuclides in soil samples within the study area

Sample Location	Sample Code	Activity Concentration (Bq/Kg)		
		^{40}K	^{238}U	^{232}Th
Okitipupa	OKT	37.65	2.81	2.53
Iletitun	ILT	58.91	4.83	1.80
Omotosho	OMO	8.73	3.21	2.73
Igbotako	IGTK	13.7	9.55	8.83
Ode-Aye	AYE	47.59	8.20	2.83
Ode-Irele	ODIR	45.73	3.22	2.89
Iyasan	IYS	39.9	3.98	1.20
Akotogbo	AKTG	17.52	4.55	2.00
Loda	LOD	23.63	6.96	3.62
Agbabu	AGB	13.9	4.77	10.49
Igbekebo	IKB	46.72	5.73	1.42
Igbobini	IGBN	18.23	4.01	1.79
Araromi Sea-Side	ARR-SS	93.83	3.28	0.35
Mean		35.85 ± 23.75	5.01 ± 2.08	3.37 ± 2.98

The values of activity concentrations from the current study were compared with similar work on soil samples investigated from other countries by Alharbi [28]. The mean activity concentration value of ^{238}U was lower than reported values for soil of Turkey (55.42Bqkg^{-1}), Jordan (57.7Bqkg^{-1}), Bangladesh (30.93Bqkg^{-1}), Najaf-Iraq (77.33Bqkg^{-1}) and Northern India (56.02Bqkg^{-1}). It was also found that the mean value of the activity concentration of ^{232}Th was lower than reported values for soil of Turkey (22.86Bqkg^{-1}), Najaf-Iraq (9.36Bqkg^{-1}), Baghdad Iraq (21.74Bqkg^{-1}), Yemen (36.26Bqkg^{-1}), Jordan (18.1Bqkg^{-1}), Iran (43.4Bqkg^{-1}) and Saudi-Arabia (12.3Bqkg^{-1}) while the activity concentration of ^{40}K in soil samples of the present study area was lower than reported values for soil of Saudi-Arabia (535.0Bqkg^{-1}), Iran (555.1Bqkg^{-1}) and Bangladesh (467.8Bqkg^{-1}).

All the radiological health risk parameters calculated from the activity concentrations in this work are presented in Table 2. The values of radium equivalent calculated ranged from 7.786 to 23.232 (Bq/Kg) with a mean value of 12.44 ± 4.88 (Bq/Kg), which are by far lower than the world average value of 370 (Bq/Kg) [10].

Table 2: Estimated Radiological Health Risk Parameters

Sample Location	D (nGyh ⁻¹)	R _{eq} (Bq/Kg)	AEDE (mSvy ⁻¹)	AGED (mSvy ⁻¹)	ELCR x 10 ⁻³	H _{in}	H _{ex}
Okitipupa	4.47	9.327	0.006	31.08	0.019	0.033	0.025
Iletitun	5.75	11.940	0.007	40.95	0.025	0.045	0.032
Omotosho	3.56	7.786	0.004	24.07	0.015	0.030	0.021
Igbotako	10.55	23.232	0.013	70.72	0.045	0.089	0.063
Ode-Aye	7.40	15.911	0.009	52.11	0.032	0.065	0.043
Ode-Irele	5.23	10.874	0.006	36.39	0.022	0.038	0.029
Iyasan	4.18	8.768	0.005	29.84	0.018	0.034	0.024
Akotogbo	4.02	8.759	0.005	27.92	0.017	0.036	0.024
Loda	6.39	13.956	0.008	44.06	0.027	0.057	0.038
Agbabu	9.62	20.841	0.012	62.95	0.041	0.069	0.056
Igbekebo	5.37	11.358	0.007	38.31	0.023	0.046	0.031
Igbobini	3.68	7.97	0.005	25.60	0.016	0.032	0.022
Araromi Sea-Side	5.58	11.01	0.007	41.06	0.024	0.039	0.030
Mean	5.83±2.19	12.44 ±4.88	0.007±0.00	40.39±14.26	0.025±0.00	0.047±0.02	0.033±0.01
World Average	84.00	370	1.00	300.0	0.29	1.00	1.00

The calculated external hazard index (H_{ex}) varies between 0.021 (Omotosho) to 0.063 (Igbotako) with a mean value of 0.034 ± 0.01 . Similarly, the internal hazard index (H_{in}) ranged between 0.030 (Omotosho) to 0.089 (Igbotako) with a mean value of 0.047 ± 0.02 . All the soil samples had both their external (H_{ex}) and internal (H_{in}) hazard indices to be less than the safe limit (i.e less than 1)

The absorbed dose rate (D) ranged from 3.56 nGy⁻¹ (Omotosho) to 10.55 nGy⁻¹ (Igbotako) with a mean value of 5.83 ± 2.19 nGy⁻¹ which is lower than that obtained by Ademola and Ademonehin [29] in soil samples around a bituminous deposit in Ondo State, Nigeria, by Lu *et al.*, [30] in soil around Baqiao coal-fired power plant in China, by Alias *et al.*, [23] in forest, flat and slope areas of an oil palm plantation located at Jengka, Pahang, Malaysia.




Figure 2: Comparison of ELCR with the World Average

The estimated excess life cancer risk (ELCR) as presented in figure 2, varied from 0.015×10^{-3} (Omotosho) to 0.045×10^{-3} (Igbotako) with a mean value of 0.025×10^{-3} . The ELCR mean value was also found to be far lower than the safe limit of 0.29×10^{-3} . In fact, these values can be said to be near zero, an indication that the study area may be safe for a very long time, however inhabitants might likely develop cancer over time. Similarly Annual Gonadal Equivalent Dose (AGED) estimated in the current study ranged from 24.07 mSvy^{-1} (Omotosho) to 70.72 mSvy^{-1} (Igbotako) with a mean value of $40.39 \pm 14.26 \text{ mSvy}^{-1}$. The values obtained for AGED in the samples were lower than the recommended safe limit of 300 mSvy^{-1} as seen in figure 3. This implies that the gonadal values may pose no threat to the bone marrow and the bone surface cells of residents in the study area.




Figure 3: Comparison of AGED with the World Average

5. Conclusion

Assessment of radiological health risk in agricultural soil samples within bitumen belt of Ondo State, Nigeria have been carried out. From the analysis of the results, it was established that the mean value of air absorbed dose and annual effective dose equivalent are lower than the UNSCEAR standard values. Both external and internal radiation hazard indices were less than 1, while the calculated values of annual gonadal equivalent dose and excess life cancer risk were less than the recommended safe limits of $300\text{mSv}\cdot\text{y}^{-1}$ and 0.29×10^{-3} respectively. The result showed that the bitumen belt area of Ondo State, Nigeria has low level of radionuclide content hence no immediate radiological risk to the population.

References

1. Fasasi, M.K., Oyawale, A.A., Mokobia, C.E., Tchokossa, P., Ajayi, T.R. & Balogun, F.A. (2003). Natural radioactivity of the tar-sand deposits of Ondo state, south-western Nigeria. *Nuclear instruments and methods in physics research section A: Accelerators, Spectrometers, Detectors and Associated Equipment*. 505 (1-2): 449-453
2. Eganosi, E. A. (2021). Incidences of bitumen contamination of water sources in some communities of Ondo State, Nigeria. *Malaysian Journal of Civil Engineering*. 33(1): 27-33
3. Adegoke, O.S. (2000). Historical perspective of Bitumen/Tar sand development in Southwestern Nigeria. *Proceedings of the 1st International Summit on Bitumen in Nigeria*. November, 14-16.: 131-140.
4. Isinkaye, M.O., Jibiri, N.N., Bamidele, S.I. & Najam, L.A. (2018). Evaluation of radiological hazards due to natural radioactivity in bituminous soils from tar-sand belt of Southwest Nigeria using HpGe-Detector. *International Journal of Radiation Research*. 16(3): 351-362
5. Babatunde, B.B., Sikoki, F.D., Avwiri, G.O. & Chad-Umoreh, Y.E. (2019). Review of the status of radioactivity profile in the oil and gas producing areas of the Niger Delta region of Nigeria. *Journal of Environmental Radioactivity*. 202: 66-73
6. Ogedengbe, K. and Akinbile, C.O. (2009). Environmental Impact of Bitumen on Soil, Water and Plant in Lodasa area, Ode-Irele, Ondo state, Nigeria. *Journal of Science and Technology*. 29(1):
7. Kang, T.W., Park, W.P., Han, Y.U., Bong, K.M. & Kim, K. (2020). Natural and artificial radioactivity in volcanic ash soils of Jeju Island, Republic of Korea, and assessment of the radiation hazards: importance of soil properties. *Journal of Radioanalytical and Nuclear Chemistry*. 323(3): 1113–1124.
8. Ramola, R.C., Choubey, V.M., Prasad, G., Gusain, G.S., Tosheva, Z. & Kies, A. (2011). Radionuclide analysis in the soil of Kumaun Himalaya, India, using gamma ray spectrometry. *Current Science*. 906–914
9. Ajayi, O.S., Dike, C.G. & Balogun, K.O. (2018). Elemental and radioactivity analysis of rocks and soils of some selected sites in southwestern Nigeria. *Environ Forensics*. 19(2): 87–98.
10. UNSCEAR, (2000): United Nations Sources Committee on the Effects of Atomic Radiation. Report to the General Assemble, Annex B, exposure from natural radiation sources. United Nations, New York, 44-89
11. Abbasi, A. & Mirekhtyari, F. (2018). Risk assessment due to various terrestrial radionuclides concentrations scenarios. *International Journal of Radiation Biology*. 95(2):179-185
12. Gbadamosi, M.R., Afolabi, T.A., Banjoko, O.O., Ogunneye, A.L., Abudu, K.A., Ogunbanjo, O.O. & Jegede, D.O. (2018a). Spatial distribution and lifetime cancer risk due to naturally

occurring radionuclides in soils around tar-sand deposit area of Ogun State, southwest Nigeria. *Chemosphere*. 193:1036–1048.

13. Bohdan, K., Karel, Z. & Jorges, S. (1999). Bitumen in the late variscan hydrothermal vein-type uranium deposit in Czech Republic: Sources, radiation-induced alteration, and relation to mineralization. *Economic Geology*. 94: 1093-1114
14. Alexandre, P. & Kyser, T.K. (2006). Geochemistry of uraniferous bitumen in the southwest Athabasca Basin, Saskatchewan, Canada. *Economic Geology*. 101: 1605 – 1612.
15. Girigisu S, Ibeanu, I.G.E., Adeyemo, D.J., Onoja, R.A., Bappah, I.A. & Okoh, S. (2013). Assessment of radiological levels in soils from Bagega artisanal gold mining exercises at Bagega Zamfara State, Nigeria. *Archives of Applied Science Research*. 5(3):204-210
16. Otwoma, D., Patel, J.P., Bartold, S. & Mustapha, A.O. (2013). Estimation of annual effective dose and radiation hazard due to natural radionuclides in Mount Homa, southwest Kenya. *Radiation Protection Dosimetry*. 155(4): 497-504.
17. Diab, H.M., Nouh, S.A., Hamdy, A. & El-fiki, S.A. (2008). Evaluation of natural radioactivity in a cultivated area of a fertilizer factory. *Journal of Nuclear and Radiation Physics*. (3):53-62
18. Agbalagba, O.E. (2017). Assessment of excess life time cancer risk from gamma radiation level in Effurun and Warri city of Delta state, Nigeria. *Journal of Taibah University for Science*. 11: 367– 380.
19. Al Harabi, W.R., Alzahrani, J.H. & Abbady, A.G.E. (2011). Assessment of radiation hazard indices from granite rocks of the southeast Arabian Shield, Kingdom of Saudi Arabia. *Australian Journal of Basic and Applied Sciences*. 5(6): 672-682
20. Asgharizadeh, F., Abbasi, A., Hochaghani, O. & Gooya, E.S. (2012). Natural radioactivity in granite stones used as building materials in Iran. *Radiat Prot Dosimetry*. 149(3): 321-326.
21. Beretka, J. & Mathew, P.J. (1985). Natural radioactivity of Australian building materials, industrial wastes and by products. *Health Physics*. 48(1): 87-95.
22. Tsukuda, H., Hidenao, H., Hisamastu, S. & Yamasaki, S. (2002). Transfer of ¹³⁷Cs and Stable Cs from Paddy Soils to Polished Rice in Aomori, Japan. *Journal of Environmental Radioactivity*. 59 (3): 351-363.
23. Alias, M., Hamzah, Z., Saat, A., Omar, M. & Wood, A. (2008). Assessment of Absorbed Dose and Radiation Hazard Index from Natural Radioactivity. *The Malaysian Journal of Analytical Sciences*. 12 (1): 195-204.
24. UNSCEAR, (1988): United Nations Scientific Committee on the Effects and Risks of Ionizing Radiation, Source, Effects and Risks of Ionizing Radiation. Report to the General Assembly, United Nations, New York.
25. Avwiri, G. O., Olatubosun, S. A. & Ononugbu, C. P. (2014). Evaluation of Radiation Hazard Indices for Selected Dumpsites in Port Harcourt, Rivers State, Nigeria. *International Journal of Science and Technology*. 3 (10): 663 – 673
26. Taskin, H. Karavus, M., Ay, P., Topuzoglu, A., Hindiroglu, S. & Karaham, G. (2009): Radionuclide concentrations in soil and lifetime cancer risk due to gamma radioactivity in Kirklareli, Turkey. *Journal of Environmental Radioactivity*. 100(1): 49-53
27. Olawale, I.A., Adefisoye, O.P. & Catherine, B.S. (2018). Estimation of Natural Radionuclide Concentrations and Heavy Metal Contents due to Bitumen Exploitation at Gbeleju-Loda, Ode-Irele, Ondo State, Southwestern Nigeria. *Asian Journal of Physical and Chemical Sciences*. 6(4): 1-10
28. Alharbi, W.R. (2013). Natural radioactivity and dose assessment for brands of chemical and organic fertilizers used in Saudi Arabia. *Journal of Modern Physic*. 4: 344-348

29. Ademola, J.A. & Ademonehin, S. (2010). Radioactivity concentrations and dose assessment for bitumen and soil samples around a bituminous deposit in Ondo State, Nigeria. *Radioprotection*. 45(3): 359 – 368
30. Lu, X., Zhao, C., Chen, C. & Liu, W. (2012). Radioactivity level of soil around Baqiao coal-fired power plant in China. *Radiation Physics and Chemistry*. 81: 1827-183

UNDER PEER REVIEW

UNDER PEER REVIEW