

Review Article

Acquired von Willebrand Syndrome: A Review

ABSTRACT

Acquired von Willebrand Disease (AvWD) is an uncommon bleeding disorder resulting from the development of decreased quantities of circulating functional von Willebrand factor (vWF). The condition may occur due to a number of pathogenic mechanisms, and is invariably associated with an underlying disorder such as congenital heart disease, neoplasms, and autoimmunity or with the use of certain drugs. While its manifestations are similar to that of inherited von Willebrand disease (vWD) in terms of clinical signs and laboratory results, it differs in that the patient lacks a personal or family history of bleeding and presents later in life. Management of the disorder is based on treatment of the underlying condition, which usually results in resolution of the bleeding diathesis. Acute hemorrhage usually requires vWF concentrates and in some cases immunosuppression. Other options involve use of Intravenous Immunoglobulins (IVIg) or plasmapheresis. Recently, immunotherapies and novel agents such as Emicizumab, Lenalidomide and Rtiuximab have been tried as off-label options.

Keywords: *Acquired von Willebrand syndrome, Clinical Presentation, Diagnosis, Hemorrhagic Diathesis, Management, Pathophysiology.*

INTRODUCTION

The von Willebrand factor (vWF) is a glycoprotein **multimer which is involved** in the process of hemostasis: it accumulates at the site of endothelial injury and functions as a bridge between adhered platelets, as well as serving as a carrier molecule for circulating factor VIII (FVIII), increasing its plasma half-life by preventing its breakdown by activated Protein C.¹⁻³ Acquired von Willebrand syndrome (AvWS) is a rare bleeding diathesis resulting from a quantitative defect in the levels of circulating vWF.⁴ The global prevalence of the disease is estimated to be between 0.04% and 0.13%, which is thought to be an under-representation as these patients are often misdiagnosed with other disorders.^{5,6} Studies report that AvWS is increasing in prevalence both due to increased recognition of the disorder as a cause of bleeding, as well as an increase in underlying precipitants.^{4,7}

AvWS is most commonly seen in the older adult population.⁸ Clinical manifestations and lab work-up demonstrates features similar to inherited von Willebrand disease (vWD) but without a past or family history of bleeding.⁴ Patients often present with excessive bleeding following trauma/surgical procedures, or unprovoked mucocutaneous oozing; Hemorrhage associated with AvWS is not usually fatal, but may require acute management.^{3, 9} Although AvWS was first recognized in 1968 (in a patient with systemic lupus erythematosus (SLE)), it has received more attention in recent years due to the discovery of its association with various underlying disorders.^{1,10} The bleeding diathesis is usually triggered by a causal condition: it is commonly associated with the presence of lympho- and myeloproliferative disorders, solid neoplasms, cardiovascular and autoimmune diseases, as well as the use of certain drugs.^{9,11}

While the pathophysiology is multifactorial, *the synthesis and release of vWF is usually normal* (except in cases with hypothyroidism) with most cases occurring due to increased *clearance* of vWF from the blood stream.¹² Accelerated clearance may be attributable to the presence of specific and non-specific immunoglobulins, vWF adsorption onto neoplastic cells, or destruction of Large multimers of vWF (LvWF) as a result of shear stress or proteolytic degradation, however, multiple mechanisms may exist concurrently in a single patient.¹² The diagnosis of AvWS is frequently confounded by the requirement for extensive testing to differentiate it from other causes of bleeding, especially the inherited and more common vWD, while management is often complicated due to the underlying precipitant or its therapy, which usually requires complete treatment before AvWS resolves.^{4,8}

This review article explores contemporary knowledge on the pathophysiology and clinical presentation, as well as the diagnosis and management of patients with AvWS.

PATHOPHYSIOLOGY

Physiology:

The vWF is synthesized as dimeric subunits, which undergo multimerization to become a large precursor protein in the endoplasmic reticulum and Golgi apparatus of endothelial cells (stored in and secreted from Weibel-Palade bodies) and in megakaryocytes (alpha granules).^{13,14} This glycoprotein undergoes several processing steps post-translation before being secreted into the circulation, and its release is regulated by several stimuli, including thrombin, histamine, and epinephrine.^{14, 15} Its molecule is composed of several domains, including a signal peptide, propeptide, D domains, A1-A3 domains, B domains, and a C-terminal domain. The propeptide is

cleaved during processing, and the mature vWF molecule is composed of identical subunits linked by disulfide bonds.^{13, 16} It is a multimeric molecule with a size as large as 20,000 kDa, while its subunits may be as small as 250 kDa.^{17, 18}

This factor has several important functions in hemostasis, including platelet adhesion, aggregation and stabilizing FVIII in the blood.^{2, 18} Platelet adhesion occurs when vWF binds to the glycoprotein Ib/IX/V receptor complex on the surface of platelets. This interaction is critical for platelet plug formation and the initiation of hemostasis.¹⁹ In addition, vWF also binds to collagen, which is exposed when the endothelial lining is damaged, which helps to anchor platelets at the site of injury and promotes platelet activation and aggregation.^{19, 20} Higher multimers are more efficient at binding platelets and are more haemostatic.²¹ This factor also plays a crucial role in stabilizing FVIII in the blood.¹⁶ FVIII is a co-factor for the intrinsic pathway of coagulation and is rapidly degraded in the absence of vWF by activated Protein C.¹⁻³ The vWF binds to and protects FVIII from degradation, thereby increasing its half-life in plasma.^{16,22,23}

The activity of vWF is regulated by several mechanisms, including proteolysis, cleavage, and clearance.²⁴ Proteolysis occurs when vWF multimers are cleaved by the metalloprotease ADAMTS13 (as well as other metalloproteinases, plasmin and leukocyte-derived proteases), which cleaves vWF between the A2 and A3 domains. This cleavage is important for regulating vWF activity, as larger vWF multimers are more active than smaller ones.^{25, 26} Cleavage can also occur through the action of thrombin, which cleaves vWF at the N-terminus of the A1 domain. This cleavage reduces the ability of vWF to bind to platelets and collagen.⁴ Clearance of vWF from the circulation occurs via receptor-mediated endocytosis, where endothelial cells and hepatocytes take up vWF.^{4, 14}

Pathology: AvWS was previously thought to be an uncommon cause of bleeding diatheses, however, the determination of its association with structural heart disease (valvular abnormalities, septal defects) in recent times established that the disease appears to be more common than previously proposed.^{5, 6,27} It is now understood to be invariably associated with an underlying disorder. These include common causes such as cardiovascular disorders, lymphoproliferative diseases (LPDs) including chronic lymphocytic leukemia and plasma cell dyscrasias such as monoclonal gammopathy of undetermined significance (MGUS), multiple myeloma and Waldenström macroglobulinemia. Less frequent provoking disorders include myeloproliferative neoplasms (MPNs) such as essential thrombocythaemia, polycythemia rubra vera, chronic myeloid leukemia, primary myelofibrosis, solid tumors (such as Wilms tumours), autoimmune disease (for e.g., systemic lupus erythematosus) and certain drugs (sodium valproate, griseofulvin, ciprofloxacin, hydroxyethyl starch).^{4,28-30}

AvWS has a complex, multifactorial pathophysiology which commonly revolves around the increased clearance of vWF from plasma secondary to shear stress, proteolysis, development of autoantibodies or cell surface adsorption, consistently occurring secondary to a provoking disorder.^{31,32} Less commonly, such as in cases with hypothyroidism, patients develop AvWS due to a decrease in production of vWF (the factor itself is qualitatively normal). Hypothyroidism may also be associated with impaired release of vWF into circulation.³³

Shear stress can result in the degradation of LvWF due to mechanical destruction in patients with valvular heart disease, left ventricular assist devices (LVADs), extracorporeal membrane oxygenation (ECMO) or areas of vascular stenosis.³² Moreover, LvWF undergoes a

tumbling motion as it moves through circulation which applies force on its structure causing it to elongate and unfold making it susceptible to degradation by enzymes.³⁴

LvWF can also undergo increased rates of degradation secondary to proteolysis by the enzyme ADAMTS13 (a disintegrin and metalloproteinase with a thrombospondin type 1 motif, member 13), especially in cases with MPNs.^{4, 35} Additionally, patients with LPDs (particularly plasma cell dyscrasias) and those with autoimmune diseases may develop autoantibodies against domains (both functional and non-functional) within the structure of vWF. These immune complexes are then removed from blood circulation by the reticulo-endothelial system.^{28,36} Direct inhibitors of vWF's interaction with circulating platelets may also occur, however, these are a rare occurrence.⁴

Lastly, adsorption of LvWF on malignant cells in cases with LPDs (Waldenström macroglobulinemia, non-Hodgkin lymphoma, Multiple Myeloma), MPNs and solid tumors (such as Wilms tumours) results in increased clearance from blood circulation, which results in the development of AvWS, especially in patients with high cell loads.^{37,38} Furthermore, certain tumour cells, such as the clonal plasma cells found in MGUS, exhibit an anomalous expression of glycoprotein Ib (GPIb) which allows vWF to bind to it, resulting in decreased circulating levels. This is also true for MPNs such as essential thrombocythaemia as well as reactive thrombocytosis, as GPIb is also expressed on platelets, and hence patients with thrombocytosis may have low circulating vWF levels.^{4,39-41} The frequency of association of different underlying conditions with the development of AvWS, and the likely precipitating mechanisms, is displayed in **Table 1**.

Table-1: The frequency and likely mechanism of Acquired von Willebrand Disease

Disorder	Frequency	Mechanism
Lymphoproliferative disorders		
Chronic lymphocytic leukaemia	Anecdotal ⁴²	Likely immune-mediated ⁴²
Multiple myeloma	7 - 15% ^{43,44}	Autoantibody, adsorption ⁴³⁻⁴⁵
Lymphoplasmacytic lymphoma	13% ⁴⁶	Increased degradation ⁴⁶
Waldenström macroglobulinemia	Anecdotal ²⁸	Increased degradation ²⁸
Monoclonal gammopathy of undetermined significance	Anecdotal ⁴⁷	Increased clearance, increased degradation ⁴⁸
Solid tumours		
Wilms tumor	8% ^{49,50}	Likely adsorption ⁵⁰
Cardiovascular disorders		
Congenital valvular stenosis	12% ⁵¹	Shear stress ⁵¹
Congenital heart disease	21% ⁵²	Likely shear stress ⁵³
Left ventricular assist devices	100% ⁵⁴	Shear stress, increased degradation ⁵⁴
Extra-corporeal membrane oxygenation	100% ⁵⁵	Likely shear stress ^{53,55}
Myeloproliferative neoplasms		
Essential Thrombocythaemia	20 55% ^{56,57}	Adsorption, increased degradation ⁵⁶
Polycythaemia rubra vera	17 49% ^{57,58}	Adsorption, increased degradation ⁵⁶⁻⁵⁸
Chronic myeloid leukaemia	Anecdotal ⁵⁹	Unknown
Primary myelofibrosis	15 33% ^{58,60}	Adsorption, increased degradation ⁵⁶⁻⁵⁸
Other Disorders		
Autoimmune disease	2% ⁶¹	Immune-mediated ⁶¹
Hypothyroidism	32% ⁶²	Low production ⁶²

CLINICAL PRESENTATION

Patients with AvWS usually present in late adult life, with a median age at diagnosis of around 60 years, and typically do not have a past or family history of a bleeding diathesis.⁶³ The clinical presentation can vary widely depending on the underlying cause of the condition, as well

as its severity. Bleeding is usually mucosal, as in vWD, and involves the nasal, oral, uterine, vaginal and/or gastrointestinal mucosae. Patients may also present with easy bruisability, menorrhagia, or prolonged bleeding after injury or surgery. They may present with symptoms and signs related to the underlying medical condition, such as an autoimmune disease, LPDs, MPNs, or cardiovascular disease.^{4,8,64} The ISTH registry reports that patients with AvWS secondary to LPDs have a greater tendency to have more severe episodes of hemorrhage compared to those who suffer from other underlying conditions; however, this may simply be attributable to the higher incidence with which AvWS occurs in LPDs.⁶¹ It should be noted here that death secondary to hemorrhage precipitated by AvWS is a rare occurrence.⁶¹

The association of vWD with the formation of angiodysplasias is well described, and these lesions also form in AvWS. It is estimated to occur in 12% of all cases of vWD.⁶⁵ The most common sites involved are the caecum and ascending colon, but patients may have multiple lesions affecting different parts of the gastrointestinal tract. Angiodysplasias are delicate and mild insults can lead to intractable bleeding.⁶⁶ While the exact mechanism of formation is unclear, angiodysplasias are thought to form due to impaired platelet adhesion and aggregation caused by vWF deficiency, resulting in blood vessel wall weakness and dilatation. In addition, an imbalance between circulating vWF and factors involved in angiogenesis may result in an overgrowth of blood vessels that are prone to rupture and bleeding.⁶⁶

DIAGNOSIS

The diagnosis of AvWS is suspected in adult patients with a new-onset bleeding diathesis, in the absence of a past or family history of bleeding disorders.⁴⁻⁸ A number of general and specific investigations which help in establishing a diagnosis, and are discussed as follows:

General Investigations:

- a. **Complete blood counts:** These counts, including the platelet count, are usually normal, unless the patient has an underlying platelet disorder such as essential thrombocythaemia.⁸
- b. **Prothrombin time:** Prothrombin Time (PT) is usually normal, unless deranged by an underlying condition.⁴
- c. **Activated partial thromboplastin time:** Patients who have low circulating FVIII levels have prolonged activated Partial Thromboplastin Time (aPTT), but this only occurs when FVIII levels are below 30%.⁶⁷
- d. **Bleeding time:** Bleeding time is prolonged in patients with altered vWF due to poor platelet aggregation.^{5,68}

Specific Investigations:

- a. **Platelet function analyser-100 closure time:** Platelet Function Analyser-100 Closure Time (PFA-100 CT) is also prolonged, while a normal PFA-100 CT carries a 99.3% sensitivity in predicting the absence of a bleeding diathesis secondary to vWF dysfunction but cannot differentiate between vWD and AvWS.⁶⁹

- b. **vWF: Antigen levels:** No cut-off levels have been described for vWF: Antigen (vWF:Ag), and those for vWD are used.^{3,4} Levels of vWF:Ag <30 IU/dL are considered indicative of vWD, while levels between 30 – 50 IU/dL are suspicious of it.⁷⁰ The vWF:Ag levels in plasma may be normal or decreased in patients with AvWS, depending on etiology.^{4,8} However, it should be noted that a number of factors such as race and blood group O individuals, among others, can affect the levels of circulating vWF:Ag, and may not be adequately reflective of the presence of AvWS.^{71,72}
- c. **vWF: Ristocetin Cofactor activity:** The vWF: Ristocetin Cofactor (vWF:RCo) activity is markedly reduced in patients with AvWS, and the vWF:RCo/vWF:Ag ratio is typically below 0.7.⁷³
- d. **vWF: Collagen Binding activity:** The vWF: Collagen Binding (vWF:CB) activity is also very low, but this test is only employed when the aforementioned tests do not yield conclusive answers.⁷⁴
- e. **vWF multimer electrophoresis:** This assay will help to detect a quantitative decrease in LvWF, which can be seen in patients with AvWS, especially when the primary pathology causes shear destruction of LvWF.^{51-55,75}
- f. **Plasma vWF propeptide:** Plasma vWF propeptide (vWFpp) indicates the degree of biosynthesis of vWF and a vWFpp/vWF:Ag ratio of >2 indicates that there is an increased clearance of vWF, which can occur in AvWS.⁷⁶
- g. **FVIII activity levels:** FVIII levels are low in AvWS, as less vWF is available to protect FVIII from degradation.⁴⁻⁸

While the aforementioned tests are useful in establishing whether there is a pathology in the function of vWF, none of them help distinguish AvWS from vWD. Diagnosis may be further

complicated by the use of antithrombotic/anticoagulation therapy for underlying disorders. Diagnosing AvWS may be based on the algorithm displayed in **Figure 1**.




Figure-1: Diagnosing acquired von Willebrand syndrome and differentiating it from inherited von Willebrand disease.

Testing for autoantibodies against vWF have a limited role in the diagnostic evaluation of AvWS, as such cases account for only a minority of patients.^{37,77} However, since such patients are known to have a higher tendency to have more severe episodes of bleeding (as autoantibodies bind to and inactivate LvWF preferentially), they should be tested in each patient of AvWS, as their presence provides valuable information with regards to clinical course/prognosis.⁷⁸ Such

testing is based on mixing studies wherein patient plasma is mixed with normal plasma and incubated at body temperature followed by the estimation of vWF:RCo and vWF:CB levels.⁷⁹ The presence of an inhibitor will result in low activity of these two compounds. However, it should be noted that this test is limited by the inability to detect autoantibodies that increase the clearance of vWF from plasma without decreasing its activity.^{4,79,80} More advanced techniques such as Enzyme-Linked Immuno-Sorbent Assays (ELISA) may be utilized to detect such antibodies, however, such methods are not routinely employed.⁸⁰

In short, establishing a concrete diagnosis of AvWS is problematic as it is difficult to differentiate vWD from AvWS because of similar clinical and laboratory pictures, and a combination of high clinical suspicion and suggestive laboratory workup should be considered before labeling a patient with AvWS, particularly when a patient presents with mild symptoms and at a young age. Furthermore, differentiating AvWS from vWD is of paramount importance as the management modalities are quite different.

MANAGEMENT

Management of AvWS involves identifying and treating the precipitating condition, controlling bleeding once it occurs, and preventing future hemorrhagic episodes:

Treating underlying condition/Inhibitor eradication: Identification and management of the underlying precipitant is the only process by which a durable cure can be achieved, such as with immunosuppression, chemotherapy or withdrawing offending drug etc..¹⁻⁸ It is pertinent to note here that patients with hypothyroidism may develop a resolution of AvWS following adequate treatment with thyroxine.¹⁻⁸

Managing bleeding symptoms: Hemorrhages may range from mild to severe. The management of bleeding symptoms may involve the administration of specific drugs to promote clotting, and/or replacement of plasma vWF:

- a. **Desmopressin:** This is an intravenously/subcutaneously/intranasally administered synthetic analog of 1-Deamino-8-D-Arginine Vasopressin (DDAVP) which works by rapidly releasing endothelial vWF.^{38,77} It is given at a dose of 0.3 mg/kg every 12 to 24 hours (maximum three doses due to tachyphylaxis) to manage active bleeding, as well as for prevention, in AvWS.^{38,81} Patients receiving DDAVP while suffering from MGUS have demonstrated a significant increase in levels of plasma vWF after administration, but the effect was short-lived. This may be due to the presence of circulating anti-vWF antibodies, especially in cases precipitated by LPDs, which results in rapid deactivation and clearance of DDAVP-released vWF.⁸² The effect was also transient in certain MPNs due to the enzymatic degradation of released vWF, or its adsorption of the neoplastic cell surface.⁴³ Previously, the ISTH registry declared that approximately a third of patients treated with DDAVP had an adequate response, but this greatly varied with the underlying disorder: patients with cardiovascular disease and MPNs had a lower response rate compared to those who suffered from autoimmune or LPDs.⁶¹
- b. **vWF concentrates:** These concentrates are derived from donor plasma and also usually contain other clotting factors such as FVIII, which can be employed in the treatment of AvWS. The ISTH registry recorded that the frequency of a positive response to vWF/FVIII concentrates was 40%, when vWF is administered at dosages ranging from 30-100 units/kg.⁶¹ However, the half-life of factors infused in this manner, including

vWF, is shorter in cases with AvWS compared to patients with vWD, thus increased doses need to be employed to attain therapeutic plasma levels.^{43,82,83}

- c. **Antifibrinolytic agents:** Drugs such as aminocaproic acid and tranexamic acid act by inhibiting the activation of plasminogen, thus prolonging the half-life of the fibrin meshwork essential for a stable clot.⁸⁴ Both these medications can be given orally, locally or intravenously, and serve as useful adjuncts to the treatments mentioned previously, both for prophylaxis and for patients with acute bleeding.^{4,84} However, as a single agent these drugs can only be used for minor bleeding episodes. Aminocaproic acid is given at a dose of 50-60 mg/kg every 4-6 hours, while tranexamic acid is dosed at 20-25 mg/kg every 8-12 hours.^{4,31}
- d. **Recombinant Activated Factor VII:** Recombinant Activated Factor VII (rFVIIa) works by bypassing the FVIII/IX coagulation pathway via two mechanisms: i) at low concentrations, rFVIIa overwhelms the inhibitory action of zymogen factor VII which increases thrombin generation, an action which requires the presence of tissue factor, or ii) at high concentrations, rFVIIa binds to activated platelets activating factor X in sufficient quantities to facilitate a spike in thrombin generation, which is tissue factor independent.^{85,86} It is administered at a dose of 90 µg/kg, usually in about three doses and has been used effectively in patients with AvWS who have failed to respond to DDAVP and vWF/FVIII concentrates.^{4,7}

Preventing Future Episodes:

- a. **Intravenous Immunoglobulins:** Intra-Venous Immunoglobulins (IVIg) may be effective in cases of AvWS via a variety of mechanisms including: i) neutralizing autoantibodies and inhibiting their secretion, ii) neutralizing circulating immune complexes and iii)

blocking Fc receptors and preventing internalization of vWF by endothelial cells.⁴ IVIg is administered at a dose of 1 g/kg/day for two days or 0.4 g/kg/day for five days in patients with AvWS.^{4,87} Response is usually noted between one to two days after administration which is seen as an increase in circulating vWF and FVIII:C levels. The response may last for up to three to four weeks, necessitating repeat dosing after three weeks of administration to maintain efficacy.^{4,87} However, it is appropriate to note here improvement in plasma levels of these factors lags behind administration of IVIg making it of limited utility in patients with an acute hemorrhage, necessitating the administration of fast-acting agents, such as those mentioned above.^{4,87}

IVIg is reported to produce a successful response in one-third patients suffering from AvWS according to the ISTH registry, and has been seen to be chiefly useful in patients suffering from solid tumors, autoimmune diseases and LPDs, particularly in cases with AvWS secondary to MGUS.⁶¹ A trial comparing IVIg to DDAVP and VWF/FVIII concentrates in patients suffering from AvWS secondary to MGUS demonstrated that IVIg produced more sustained increments in levels of circulating vWF and FVIII.⁸⁸ However, this benefit was not seen in patients suffering from MGUS where the antibody class involved was IgM. Furthermore, prophylactic infusions of IVIg every three weeks was associated with a significant reduction in gastrointestinal bleeding in patients with AvWS secondary to IgG-MGUS.⁸⁸

- b. **Plasmapheresis:** Plasmapheresis can be employed in cases where autoantibodies are suspected to be responsible for reduced levels of circulating vWF in patients with AvWS, and is noted to be particularly useful in patients suffering from IgM-MGUS, who typically are non-responsive to other treatments such as IVIg.^{4,89}

Novel Therapies: While traditional therapies such as IVIg and plasmapheresis can be effective in the management of AvWS, novel therapies are also being explored for its management with varying degrees of response:

- a. **Recombinant vWF:** Recombinant vWF (rvWF) represents a potential treatment modality that can replace traditional plasma-derived vWF concentrates in managing patients with AvWS, however, experience in this setting is limited. It is associated with a longer half-life compared to plasma-derived vWF and contains LvWF. The standard dose is 80 IU/kg intravenously.⁹⁰
- b. **Rituximab:** This is a monoclonal anti-CD20 antibody that targets B-lymphocytes, which play a role in the production of anti-vWF autoantibodies. It has been seen to be effective at doses of 150 mg to 375 mg/m² weekly for three to four doses in some patients with AvWS, resulting in a substantial improvement in coagulation profiles.^{7,91}
- c. **Lenalidomide:** Lenalidomide treatment has been seen to be associated with an increase in plasma vWF antigen (vWF:Ag) and vWF ristocetin cofactor (vWF:RCo) levels in patients with AvWS secondary to MGUS, which may be sustained for up to two years post-treatment. Furthermore, this effect may be ascribed not just to its action on plasma cells but also due to decreased plasma clearance of vWF.^{92,93}
- d. **Emicizumab:** This is a bispecific, FVIII-mimetic antibody that mimics the in-vivo function of FVIII, which is deficient in hemophilia A. A shorter half-life of FVIII in patients with AvWS compounds bleeding tendencies of such patients and Emicizumab has been shown to improve bleeding symptoms in patients with acquired hemophilia A associated with vWD, and its use may be extrapolated to patients with AvWS in future.⁹⁴

Adjunctive Management: Patients may develop bleeding severe enough to warrant transfusions of red cell concentrates.⁹⁵ Furthermore, drugs that interfere with coagulation such as aspirin, non-steroidal anti-inflammatory drugs (NSAIDs), direct oral anticoagulants (DOACs), warfarin and parental anticoagulants should be avoided as they increase the risk of bleeding.⁹⁶ Patients should be managed by a multidisciplinary team to manage AvWS as well as its underlying cause for effective management. This may involve collaboration between hematologists, oncologists, rheumatologists, and other healthcare professionals. Moreover, patients should remain on regular monitoring and follow-ups to monitor their bleeding symptoms and adjust treatment as needed.⁴ Additionally, the treatment modalities mentioned above should be used with caution in patients who are at risk for the development of thrombosis, especially rapidly acting agents like DDAVP.^{97,98} Bleeding secondary to gastrointestinal angiodysplasia and patients of AvWS due to undergo surgery represent a unique challenge to manage: a number of different modalities, in addition to conventional methods (like DDAVP and vWF concentrates), such as thalidomide, rituximab, octreotide, and the use of statins have been described in literature.^{95,99,100}

CONCLUSION

AvWS is becoming an increasingly common bleeding diathesis, with a multifactorial etiology, and is almost universally linked with an underlying disease. The disorder itself is usually associated with mild to moderate symptomatology, but can rarely be life-threatening. Patients typically present in late adulthood with laboratory findings suggestive of vWD but without a past or family history of a hemorrhagic diathesis. This makes establishing a diagnosis quite challenging as there is no specific gold-standard investigation; diagnoses are made on clinical suspicion and should be established in consultation with clinical hematologists and

hematopathologists. Screening for this bleeding disorder in patients with neoplastic or autoimmune disease presenting with a hemorrhagic diathesis is advisable. Treatment involves control of acute hemorrhage with short-term hemostatic measures such as DDAVP, factor concentrates and anti-fibrinolytics, intermediate-term control can be obtained with modalities like IVIg and plasmapheresis and long-term strategies such as chemo-immunotherapy for underlying Lymphoma or Waldenstrom's. Novel agents such as immunotherapies require further investigation before they can be prescribed more frequently.

REFERENCES

1. Kozlov S, Okhota S, Avtaeva Y, Melnikov I, Matroze E, Gabbasov Z. Von Willebrand factor in diagnostics and treatment of cardiovascular disease: Recent advances and prospects. *Front Cardiovasc Med.* 2022 Dec 2;9:1038030. doi: 10.3389/fcvm.2022.1038030.
2. Cortes GA, Moore MJ, El-Nakeep S. Physiology, Von Willebrand Factor. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2022 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK559062/>.
3. Sabih A, Babiker HM. Von Willebrand Disease. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2022 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK459222/>.
4. Franchini M, Mannucci PM. Acquired von Willebrand syndrome: focused for hematologists. *Haematologica.* 2020 Aug;105(8):2032-2037. doi: 10.3324/haematol.2020.255117.
5. Kumar S, Pruthi RK, Nichols WL. Acquired von Willebrand disease. *Mayo Clin Proc.* 2002 Feb;77(2):181-7. doi: 10.4065/77.2.181.
6. Federici AB. Acquired von Willebrand syndrome: an underdiagnosed and misdiagnosed bleeding complication in patients with lymphoproliferative and myeloproliferative disorders. *Semin Hematol.* 2006 Jan;43(1 Suppl 1):S48-58. doi: 10.1053/j.seminhematol.2005.11.003.
7. Leebeek FW. New Developments in Diagnosis and Management of Acquired Hemophilia and Acquired von Willebrand Syndrome. *Hemasphere.* 2021 Jun 1;5(6):e586. doi: 10.1097/HS9.0000000000000586.
8. Mital A. Acquired von Willebrand Syndrome. *Adv Clin Exp Med.* 2016 Nov-Dec;25(6):1337-1344. doi: 10.17219/acem/64942.

9. Charlebois J, Rivard GÉ, St-Louis J. Management of acquired von Willebrand syndrome. *Transfus Apher Sci.* 2018 Dec;57(6):721-723. doi: 10.1016/j.transci.2018.10.012.
10. Simone JV, Cornet JA, Abildgaard CF. Acquired von Willebrand's syndrome in systemic lupus erythematosus. *Blood.* 1968 Jun;31(6):806-12.
11. Langer AL, Connell NT. Acquired von Willebrand Syndrome. *Hematol Oncol Clin North Am.* 2021 Dec;35(6):1103-1116. doi: 10.1016/j.hoc.2021.07.005.
12. Shetty S, Kasatkar P, Ghosh K. Pathophysiology of acquired von Willebrand disease: a concise review. *Eur J Haematol.* 2011 Aug;87(2):99-106. doi: 10.1111/j.1600-0609.2011.01636.x.
13. Haberichter SL. von Willebrand factor propeptide: biology and clinical utility. *Blood.* 2015 Oct 8;126(15):1753-61. doi: 10.1182/blood-2015-04-512731.
14. Mojzisch A, Brehm MA. The Manifold Cellular Functions of von Willebrand Factor. *Cells.* 2021 Sep 8;10(9):2351. doi: 10.3390/cells10092351.
15. Michels A, Swystun LL, Mewburn J, Albáñez S, Lillicrap D. Investigating von Willebrand Factor Pathophysiology Using a Flow Chamber Model of von Willebrand Factor-platelet String Formation. *J Vis Exp.* 2017 Aug 14;(126):55917. doi: 10.3791/55917.
16. Peyvandi F, Garagiola I, Baronciani L. Role of von Willebrand factor in the haemostasis. *Blood Transfus.* 2011 May;9 Suppl 2(Suppl 2):s3-8. doi: 10.2450/2011.002S.
17. Furlan M. Von Willebrand factor: molecular size and functional activity. *Ann Hematol.* 1996 Jun;72(6):341-8. doi: 10.1007/s002770050184.
18. Sadler JE. Biochemistry and genetics of von Willebrand factor. *Annu Rev Biochem.* 1998;67:395-424. doi: 10.1146/annurev.biochem.67.1.395.
19. Bryckaert M, Rosa JP, Denis CV, Lenting PJ. Of von Willebrand factor and platelets. *Cell Mol Life Sci.* 2015 Jan;72(2):307-26. doi: 10.1007/s00018-014-1743-8.
20. Ruggeri ZM, Mendolicchio GL. Interaction of von Willebrand factor with platelets and the vessel wall. *Hamostaseologie.* 2015;35(3):211-24. doi: 10.5482/HAMO-14-12-0081.
21. Bharati KP, Prashanth UR. Von Willebrand disease: an overview. *Indian J Pharm Sci.* 2011 Jan;73(1):7-16. doi: 10.4103/0250-474X.89751.
22. Federici AB. The factor VIII/von Willebrand factor complex: basic and clinical issues. *Haematologica.* 2003 Jun;88(6):EREPO2.
23. Fuller JR, Knockenhauer KE, Leksa NC, Peters RT, Batchelor JD. Molecular determinants of the factor VIII/von Willebrand factor complex revealed by BIVV001 cryo-electron microscopy. *Blood.* 2021 May 27;137(21):2970-2980. doi: 10.1182/blood.2020009197.
24. Schwameis M, Schörghofer C, Assinger A, Steiner MM, Jilma B. VWF excess and ADAMTS13 deficiency: a unifying pathomechanism linking inflammation to thrombosis in DIC, malaria, and TTP. *Thromb Haemost.* 2015 Apr;113(4):708-18. doi: 10.1160/TH14-09-0731.
25. Raife TJ, Cao W, Atkinson BS, Bedell B, Montgomery RR, Lentz SR, Johnson GF, Zheng XL. Leukocyte proteases cleave von Willebrand factor at or near the ADAMTS13 cleavage site. *Blood.* 2009 Aug 20;114(8):1666-74. doi: 10.1182/blood-2009-01-195461.

26. Chen J, Chung DW. Inflammation, von Willebrand factor, and ADAMTS13. *Blood*. 2018 Jul 12;132(2):141-147. doi: 10.1182/blood-2018-02-769000.
27. Drenko P, Přenosilová P, Lavičková A. Acquired von Willebrand syndrome (AvWS). *Vnitr Lek*. 2022 Fall;68(E-4):16-19.
28. Wolfe Z, Lash B. Acquired von Willebrand Syndrome in IgM Monoclonal Gammopathy as the Presentation of Lymphoplasmacytic Lymphoma. *Case Rep Hematol*. 2017;2017:9862620. doi: 10.1155/2017/9862620.
29. Gkalea V, Fotiou D, Dimopoulos MA, Kastritis E. Monoclonal Gammopathy of Thrombotic Significance. *Cancers (Basel)*. 2023 Jan 12;15(2):480. doi: 10.3390/cancers15020480.
30. Dicke C, Schneppenheim S, Holstein K, Spath B, Bokemeyer C, Dittmer R, Budde U, Langer F. Distinct mechanisms account for acquired von Willebrand syndrome in plasma cell dyscrasias. *Ann Hematol*. 2016 May;95(6):945-57. doi: 10.1007/s00277-016-2650-x.
31. Tiede A. Diagnosis and treatment of acquired von Willebrand syndrome. *Thromb Res*. 2012 Dec;130 Suppl 2:S2-6. doi: 10.1016/S0049-3848(13)70003-3.
32. Mehta R, Athar M, Girgis S, Hassan A, Becker RC. Acquired Von Willebrand Syndrome (AVWS) in cardiovascular disease: a state of the art review for clinicians. *J Thromb Thrombolysis*. 2019 Jul;48(1):14-26. doi: 10.1007/s11239-019-01849-2.
33. Manfredi E, van Zaane B, Gerdes VE, Brandjes DP, Squizzato A. Hypothyroidism and acquired von Willebrand's syndrome: a systematic review. *Haemophilia*. 2008 May;14(3):423-33. doi: 10.1111/j.1365-2516.2007.01642.x.
34. Zhussupbekov M, Méndez Rojano R, Wu WT, Massoudi M, Antaki JF. A Continuum Model for the Unfolding of von Willebrand Factor. *Ann Biomed Eng*. 2021 Sep;49(9):2646-2658. doi: 10.1007/s10439-021-02845-5.
35. Zheng XL. ADAMTS13 and von Willebrand factor in thrombotic thrombocytopenic purpura. *Annu Rev Med*. 2015;66:211-25. doi: 10.1146/annurev-med-061813-013241.
36. Owari M, Harada-Shirado K, Togawa R, Fukatsu M, Sato Y, Fukuchi K, Endo M, Takahashi H, Kimura S, Osaki T, Souri M, Ichinose A, Shibata Y, Hashimoto Y, Ikezoe T. Acquired von Willebrand Syndrome in a Patient with Multiple Comorbidities, Including MALT Lymphoma with IgA Monoclonal Gammopathy and Hyperviscosity Syndrome. *Intern Med*. 2023 Feb 15;62(4):605-611. doi: 10.2169/internalmedicine.9815-22.
37. van Genderen PJ, Michiels JJ. Acquired von Willebrand disease. *Baillieres Clin Haematol*. 1998 Jun;11(2):319-30. doi: 10.1016/s0950-3536(98)80051-6.
38. Mohri H. Acquired von Willebrand syndrome: its pathophysiology, laboratory features and management. *J Thromb Thrombolysis*. 2003 Jun;15(3):141-9. doi: 10.1023/B:THRO.0000011369.70824.e6.
39. Tefferi A, Hanson CA, Kurtin PJ, Katzmann JA, Dalton RJ, Nichols WL. Acquired von Willebrand's disease due to aberrant expression of platelet glycoprotein Ib by marginal zone lymphoma cells. *Br J Haematol*. 1997 Mar;96(4):850-3. doi: 10.1046/j.1365-2141.1997.d01-2088.x.
40. Mayne ES, Tait M, Jacobson BF, Pillay E, Louw SJ. Combination of acquired von Willebrand syndrome (AVWS) and Glanzmann thrombasthenia in monoclonal

- gammopathy of uncertain significance (MGUS), a case report. *Thromb J*. 2018 Nov 26;16:30. doi: 10.1186/s12959-018-0184-2.
41. Yee MEM, Batsuli G, Chonat S, Park S. Thrombocytosis with acquired von Willebrand disease in an adolescent with sickle cell disease. *Clin Case Rep*. 2020 Nov 20;9(1):457-460. doi: 10.1002/ccr3.3556.
 42. Innocenti I, Morelli F, Autore F, Tomasso A, Corbingi A, Bellesi S, Za T, De Stefano V, Laurenti L. Remission of acquired von Willebrand syndrome in a patient with chronic lymphocytic leukemia treated with venetoclax. *Leuk Lymphoma*. 2019 Dec;60(12):3078-3080. doi: 10.1080/10428194.2019.1612063.
 43. Mohri H, Motomura S, Kanamori H, Matsuzaki M, Watanabe S, Maruta A, Kodama F, Okubo T. Clinical significance of inhibitors in acquired von Willebrand syndrome. *Blood*. 1998 May 15;91(10):3623-9. Erratum in: *Blood* 1999 Jan 1;93(1):413.
 44. Djunic I, Elezovic I, Ilic V, Tomin D, Suvajdzic-Vukovic N, Bila J, Antic D, Vidovic A, Milosevic-Jovcic N. Acquired von Willebrand syndrome in multiple myeloma. *Hematology*. 2011 Jul;16(4):209-12. doi: 10.1179/102453311X12953015767617.
 45. Mohri H, Tanabe J, Ohtsuka M, Yoshida M, Motomura S, Nishida S, Fujimura Y, Okubo T. Acquired von Willebrand disease associated with multiple myeloma; characterization of an inhibitor to von Willebrand factor. *Blood Coagul Fibrinolysis*. 1995 Sep;6(6):561-6. doi: 10.1097/00001721-199509000-00010.
 46. Hivert B, Caron C, Petit S, Charpy C, Fankam-Siaka C, Lecocq S, Zawadzki C, Susen S, Rusu M, Duhamel A, Tournilhac O, Goudemand J, Morel P. Clinical and prognostic implications of low or high level of von Willebrand factor in patients with Waldenstrom macroglobulinemia. *Blood*. 2012 Oct 18;120(16):3214-21. doi: 10.1182/blood-2011-11-388256.
 47. Vanderschueren S, Mylle M, Dierickx D, Delforge M, Verhamme P, Peerlinck K, Meersseman W, Knockaert DC. Monoclonal gammopathy of undetermined significance: significant beyond hematology. *Mayo Clin Proc*. 2009 Sep;84(9):842-5. doi: 10.1016/S0025-6196(11)60495-3.
 48. Galletta E, Galvanin F, Bertomoro A, Daidone V, Casonato A. Acquired von Willebrand syndrome in patients with monoclonal gammopathy of undetermined significance investigated using a mechanistic approach. *Blood Transfus*. 2023 Jan;21(1):74-82. doi: 10.2450/2021.0121-21.
 49. Coppes MJ, Zandvoort SW, Sparling CR, Poon AO, Weitzman S, Blanchette VS. Acquired von Willebrand disease in Wilms' tumor patients. *J Clin Oncol*. 1992 Mar;10(3):422-7. doi: 10.1200/JCO.1992.10.3.422.
 50. Fosbury E, Szychot E, Slater O, Mathias M, Sibson K. An 11-year experience of acquired von Willebrand syndrome in children diagnosed with Wilms tumour in a tertiary referral centre. *Pediatr Blood Cancer*. 2017 Mar;64(3). doi: 10.1002/pbc.26246.
 51. Binnetoğlu FK, Babaoğlu K, Filiz ŞG, Zengin E, Altun G, Kılıç SÇ, Sarper N. Acquired von Willebrand syndrome in children with aortic and pulmonary stenosis. *Cardiovasc J Afr*. 2016 Jul/Aug;27(4):222-227. doi: 10.5830/CVJA-2015-093.

52. Waldow HC, Westhoff-Bleck M, Widera C, Templin C, von Depka M. Acquired von Willebrand syndrome in adult patients with congenital heart disease. *Int J Cardiol.* 2014 Oct 20;176(3):739-45. doi: 10.1016/j.ijcard.2014.07.104.
53. Nascimbene A, Neelamegham S, Frazier OH, Moake JL, Dong JF. Acquired von Willebrand syndrome associated with left ventricular assist device. *Blood.* 2016 Jun 23;127(25):3133-41. doi: 10.1182/blood-2015-10-636480.
54. Meyer AL, Malehsa D, Budde U, Bara C, Haverich A, Strueber M. Acquired von Willebrand syndrome in patients with a centrifugal or axial continuous flow left ventricular assist device. *JACC Heart Fail.* 2014 Apr;2(2):141-5. doi: 10.1016/j.jchf.2013.10.008.
55. Kalbhenn J, Schmidt R, Nakamura L, Schelling J, Rosenfelder S, Zieger B. Early diagnosis of acquired von Willebrand Syndrome (AVWS) is elementary for clinical practice in patients treated with ECMO therapy. *J Atheroscler Thromb.* 2015;22(3):265-71. doi: 10.5551/jat.27268.
56. Mital A, Prejzner W, Bieniaszewska M, Hellmann A. Prevalence of acquired von Willebrand syndrome during essential thrombocythemia: a retrospective analysis of 170 consecutive patients. *Pol Arch Med Wewn.* 2015;125(12):914-20. doi: 10.20452/pamw.3211.
57. Rottenstreich A, Kleinstern G, Krichevsky S, Varon D, Lavie D, Kalish Y. Factors related to the development of acquired von Willebrand syndrome in patients with essential thrombocythemia and polycythemia vera. *Eur J Intern Med.* 2017 Jun;41:49-54. doi: 10.1016/j.ejim.2016.11.011.
58. Song IC, Kang S, Lee MW, Ryu H, Lee HJ, Yun HJ, Jo DY. Acquired von willebrand syndrome in patients with philadelphia-negative myeloproliferative neoplasm. *Blood Res.* 2023 Mar 31;58(1):42-50. doi: 10.5045/br.2023.2022218.
59. Jain A. A rare case of chronic myeloid leukemia with acquired von Willebrand disease presenting as subdural hematoma. *J Cancer Res Ther.* 2015 Oct-Dec;11(4):1022. doi: 10.4103/0973-1482.146136.
60. Mital A, Prejzner W, Hellmann A. Acquired von Willebrand Syndrome During the Course of Myelofibrosis: Analysis of 32 Cases. *Adv Clin Exp Med.* 2015 Nov-Dec;24(6):1001-6. doi: 10.17219/acem/52361.
61. Federici AB, Rand JH, Bucciarelli P, Budde U, van Genderen PJ, Mohri H, Meyer D, Rodeghiero F, Sadler JE; Subcommittee on von Willebrand Factor. Acquired von Willebrand syndrome: data from an international registry. *Thromb Haemost.* 2000 Aug;84(2):345-9. Erratum in: *Thromb Haemost* 2000 Oct;84(4):739.
62. Stuijver DJ, Piantanida E, van Zaane B, Galli L, Romualdi E, Tanda ML, Meijers JC, Büller HR, Gerdes VE, Squizzato A. Acquired von Willebrand syndrome in patients with overt hypothyroidism: a prospective cohort study. *Haemophilia.* 2014 May;20(3):326-32. doi: 10.1111/hae.12275.
63. Kumar S, Pruthi RK, Nichols WL. Acquired von Willebrand's syndrome: a single institution experience. *Am J Hematol.* 2003 Apr;72(4):243-7. doi: 10.1002/ajh.10298.
64. Sandrock-Lang K, Glonnegger H, Zieger B. Acquired von Willebrand Syndrome in Children. *Hamostaseologie.* 2022 Apr;42(2):117-122. doi: 10.1055/a-1790-6156.

65. Fressinaud E, Meyer D. International survey of patients with von Willebrand disease and angiodysplasia. *Thromb Haemost.* 1993 Sep 1;70(3):546.
66. Selvam S, James P. Angiodysplasia in von Willebrand Disease: Understanding the Clinical and Basic Science. *Semin Thromb Hemost.* 2017 Sep;43(6):572-580. doi: 10.1055/s-0037-1599145.
67. Little JW, Falace DA, Miller CS, Rhodus NL. Partial Thromboplastin Time, Acquired Bleeding and Hypercoagulable Disorders. In: Little and Falace's Dental Management of the Medically Compromised Patient. Mosby; 2013;409-436. Available from: <https://www.https://www.sciencedirect.com/science/article/pii/B9780323080286000245/>.
68. Nitu-Whalley IC, Lee CA. Acquired von Willebrand syndrome--report of 10 cases and review of the literature. *Haemophilia.* 1999 Sep;5(5):318-26. doi: 10.1046/j.1365-2516.1999.00340.x.
69. Favalaro EJ. Utility of the platelet function analyser (PFA-100/200) for exclusion or detection of von Willebrand disease: A study 22 years in the making. *Thromb Res.* 2020 Apr;188:17-24. doi: 10.1016/j.thromres.2020.01.029.
70. O'Donnell JS. Low VWF: insights into pathogenesis, diagnosis, and clinical management. *Blood Adv.* 2020 Jul 14;4(13):3191-3199. doi: 10.1182/bloodadvances.2020002038.
71. Bellissimo DB, Christopherson PA, Flood VH, Gill JC, Friedman KD, Haberichter SL, Shapiro AD, Abshire TC, Leissingner C, Hoots WK, Lusher JM, Ragni MV, Montgomery RR. VWF mutations and new sequence variations identified in healthy controls are more frequent in the African-American population. *Blood.* 2012 Mar 1;119(9):2135-40. doi: 10.1182/blood-2011-10-384610.
72. Gill JC, Endres-Brooks J, Bauer PJ, Marks WJ Jr, Montgomery RR. The effect of ABO blood group on the diagnosis of von Willebrand disease. *Blood.* 1987 Jun;69(6):1691-5.
73. Flood VH. Perils, problems, and progress in laboratory diagnosis of von Willebrand disease. *Semin Thromb Hemost.* 2014 Feb;40(1):41-8. doi: 10.1055/s-0033-1363166.
74. Kallas A, Talpsep T. The von Willebrand factor collagen-binding activity assay: clinical application. *Ann Hematol.* 2001 Aug;80(8):466-71. doi: 10.1007/s002770100329.
75. Thomazini CM, Soares RPS, da Rocha TRF, Sachetto ATA, Santoro ML. Optimization of von Willebrand factor multimer analysis in vertical mini-gel electrophoresis systems: A rapid procedure. *Thromb Res.* 2019 Mar;175:76-83. doi: 10.1016/j.thromres.2019.01.018.
76. Lee A, Sinclair G, Valentine K, James P, Poon MC. Acquired von Willebrand syndrome: von Willebrand factor propeptide to von Willebrand factor antigen ratio predicts remission status. *Blood.* 2014 Jul 31;124(5):e1-3. doi: 10.1182/blood-2014-02-557132.
77. Federici AB, Budde U, Castaman G, Rand JH, Tiede A. Current diagnostic and therapeutic approaches to patients with acquired von Willebrand syndrome: a 2013 update. *Semin Thromb Hemost.* 2013 Mar;39(2):191-201. doi: 10.1055/s-0033-1334867.
78. van Genderen PJ, Vink T, Michiels JJ, van 't Veer MB, Sixma JJ, van Vliet HH. Acquired von Willebrand disease caused by an autoantibody selectively inhibiting the binding of von Willebrand factor to collagen. *Blood.* 1994 Nov 15;84(10):3378-84.

79. Rinder MR, Richard RE, Rinder HM. Acquired von Willebrand's disease: a concise review. *Am J Hematol.* 1997 Feb;54(2):139-45. doi: 10.1002/(sici)1096-8652(199702)54:2<139::aid-ajh7>3.0.co;2-y.
80. Franchi F, Biguzzi E, Stufano F, Siboni SM, Baronciani L, Peyvandi F. A two-step approach (Enzyme-linked immunosorbent assay and confirmation assay) to detect antibodies against von Willebrand factor in patients with Acquired von Willebrand Syndrome. *Thromb Res.* 2014 Dec;134(6):1316-22. doi: 10.1016/j.thromres.2014.09.020.
81. Tuohy E, Litt E, Alikhan R. Treatment of patients with von Willebrand disease. *J Blood Med.* 2011;2:49-57. doi: 10.2147/JBM.S9890. Epub 2011 Apr 20. Erratum in: *J Blood Med.* 2013 May 20;4:57-8.
82. Abou-Ismaïl MY, Rodgers GM, Bray PF, Lim MY. Acquired von Willebrand syndrome in monoclonal gammopathy - A scoping review on hemostatic management. *Res Pract Thromb Haemost.* 2021 Feb 17;5(2):356-365. doi: 10.1002/rth2.12481.
83. Dane KE, Lindsley JP, Kickler T, Streiff MB, Moliterno A, Yui J, Naik R, Chaturvedi S. Continuous-infusion von Willebrand factor concentrate is effective for the management of acquired von Willebrand disease. *Blood Adv.* 2021 Jul 27;5(14):2813-2816. doi: 10.1182/bloodadvances.2021004843.
84. Levy JH, Koster A, Quinones QJ, Milling TJ, Key NS. Antifibrinolytic Therapy and Perioperative Considerations. *Anesthesiology.* 2018 Mar;128(3):657-670. doi: 10.1097/ALN.0000000000001997.
85. Monroe DM, Hoffman M, Oliver JA, Roberts HR. A possible mechanism of action of activated factor VII independent of tissue factor. *Blood Coagul Fibrinolysis.* 1998 Mar;9 Suppl 1:S15-20.
86. Giansily-Blaizot M, Schved JF. Recombinant human factor VIIa (rFVIIa) in hemophilia: mode of action and evidence to date. *Ther Adv Hematol.* 2017 Dec;8(12):345-352. doi: 10.1177/2040620717737701.
87. Federici AB, Rossi V, Sacchi E, Franchini M. Are intravenous immunoglobulins really inappropriate in acquired von Willebrand syndrome? *Blood Transfus.* 2012 Jul;10(3):402-3; author reply 404-5. doi: 10.2450/2011.0091-11.
88. Federici AB, Stabile F, Castaman G, Canciani MT, Mannucci PM. Treatment of acquired von Willebrand syndrome in patients with monoclonal gammopathy of uncertain significance: comparison of three different therapeutic approaches. *Blood.* 1998 Oct 15;92(8):2707-11.
89. Tiede A, Rand JH, Budde U, Ganser A, Federici AB. How I treat the acquired von Willebrand syndrome. *Blood.* 2011 Jun 23;117(25):6777-85. doi: 10.1182/blood-2010-11-297580.
90. Erratum: Gill JC, Castaman G, Windyga J, et al. Hemostatic efficacy, safety, and pharmacokinetics of a recombinant von Willebrand factor in severe von Willebrand disease. *Blood.* 2015;126(17):2038-2046. *Blood.* 2016 Jun 2;127(22):2777. doi: 10.1182/blood-2016-04-711259. Erratum for: *Blood.* 2015 Oct 22;126(17):2038-46.
91. Kurahashi H, Kawabata Y, Michishita Y, Kitabayashi A, Kobayashi T, Kitadate A, Takahashi N. [Durable remission attained with rituximab therapy in a patient with acquired von Willebrand syndrome associated with CD20-positive lymphoproliferative

- disorder]. *Rinsho Ketsueki*. 2018;59(4):420-425. Japanese. doi: 10.11406/rinketsu.59.420.
92. Lavin M, Brophy TM, Rawley O, O'Sullivan JM, Hayden PJ, Browne PV, Ryan K, O'Connell N, O'Donnell JS. Lenalidomide as a novel treatment for refractory acquired von Willebrand syndrome associated with monoclonal gammopathy. *J Thromb Haemost*. 2016 Jun;14(6):1200-5. doi: 10.1111/jth.13317.
 93. Green A, Shen YP, Nelson AT, Sarode R, Ibrahim IF, Cao J, Afraz S, Yates SG. Successful use of lenalidomide to treat refractory acquired von Willebrand disease associated with monoclonal gammopathy. *Ann Hematol*. 2022 Dec;101(12):2627-2631. doi: 10.1007/s00277-022-04991-9.
 94. Thomas VM, Abou-Ismaïl MY, Lim MY. Off-label use of emicizumab in persons with acquired haemophilia A and von Willebrand disease: A scoping review of the literature. *Haemophilia*. 2022 Jan;28(1):4-17. doi: 10.1111/hae.14450.
 95. Biguzzi E, Siboni SM, Peyvandi F. How I treat gastrointestinal bleeding in congenital and acquired von Willebrand disease. *Blood*. 2020 Sep 3;136(10):1125-1133. doi: 10.1182/blood.2019003702.
 96. Yee J, Kaide CG. Emergency Reversal of Anticoagulation. *West J Emerg Med*. 2019 Aug 6;20(5):770-783. doi: 10.5811/westjem.2018.5.38235.
 97. McCarty TS, Shah AD. Desmopressin. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK554582/>.
 98. Waheed W, Aljerdi S, Decker B, Cushman M, Hamill RW. Cerebral venous thrombosis associated with thyrotoxicosis, the use of desmopressin and elevated factor VIII/von Willebrand factor. *BMJ Case Rep*. 2016 Aug 8;2016:bcr2016216584. doi: 10.1136/bcr-2016-216584.
 99. Tan HH, Ge ZZ, Chen HM, Gao YJ. Successful treatment with thalidomide for a patient with recurrent gastrointestinal bleeding due to angiodysplasia diagnosed by capsule endoscopy. *J Dig Dis*. 2013 Mar;14(3):153-5. doi: 10.1111/1751-2980.12009.
 100. Franchini M, Seidizadeh O, Mannucci PM. Prophylactic management of patients with von Willebrand disease. *Ther Adv Hematol*. 2021 Dec 22;12:20406207211064064. doi: 10.1177/20406207211064064.