

## Association of natal and maternal risk factors with neonatal Apgar score in Mosul.

### Abstract:

**Background:** The Apgar score is a recognized and practical way to report a newborn infant's condition right after birth and their reaction to resuscitation, if necessary; nonetheless, it has been improperly used to forecast specific negative neurologic outcomes.

**Aim of the study:** To identify the most frequent maternal risk factors associated with low Apgar scores among newborns and evaluate the effect of neonatal sex, type of delivery, mothers' age on Apgar score at Al-Batool Teaching Hospital in Mosul.

**Patients and Methods:** This study used a cross-sectional Hospital based study design to achieve the study objectives. The data was obtained and collected from 93 maternal and neonatal record files for the mothers who delivered at Al-Batool Teaching Hospital for the period from 1<sup>st</sup> May 31<sup>st</sup> June 2023 with a simple random sampling method was used. All maternity data for women and full-term newborns, regardless of the delivery method, meet the inclusion criteria for this study. All maternal files involving home births, premature births, and referrals from other facilities following deliveries were disqualified. The information was gathered and examined by researchers.

**Results:** The current study involved 93 full term neonates; 58 were males and 35 were females with male: female ratio of 1.65:1. Of them 35 (37.6%) neonates delivered by normal vaginal delivery while the remaining 58 (62.4%) delivered through cesarean section. The mean Apgar score at 1 min was  $5.311 \pm 2.274$  while at 5 min. was  $7.462 \pm 1.891$  and the mean maternal age was  $27.57 \pm 5.942$  years. The commonest risk factors found was obstructed labour with mean Apgar score of  $3.381 \pm 1.687$  at 1 min and  $6.142 \pm 1.768$  at 5 min, followed by previous 2 CS and meconium stained liquor. Neonates with no maternal risks had mean Apgar score of  $8.285 \pm 0.755$  at 1 min and  $9.571 \pm 0.534$  at 5 min. The statistical difference between Apgar score at 1 and 5 min was significant for each of males and females while the difference between males and females

was not significant at each of 1 and 5 min. The mean Apgar score at 5 min was significantly higher than that at 1 min for those with maternal risk the difference between those with maternal risk and those with no maternal risk was statistically significant for both at 1 min and 5 min. Furthermore, mean Apgar score at 5 min was significantly higher than that at 1 min for normal vaginal delivery and cesarean section while the difference between Apgar score in normal vaginal delivery and cesarean section at each of 1 and 5 min was not significant. The Apgar scores for at both 1 and 5 min were very weakly and insignificantly correlated with the mothers' ages.

**Conclusions:** Mothers were most at risk for obstructed labor, and the mean Apgar scores of their newborns were the lowest. At both 1 and 5 minutes, there was a statistically significant difference between the groups with and without maternal risk. The difference in the Apgar score at 1 and 5 minutes for male and female newborns and between those who delivered vaginally normally and via cesarean section was not statistically significant.

**Keywords:** Apgar score, Maternal risk, Neonates

**Introduction:**

Dr. Virginia Apgar created a score in the early 1950s to evaluate the newborn's physical status and whether resuscitation is necessary <sup>(1)</sup>. The score was initially assessed at 1<sup>st</sup> minute post delivery, but it was later discovered that the Apgar score at 5 minutes was a stronger indicator of neonatal survival <sup>(2)</sup>. The five factors that make up the Apgar score—heart rate, respiratory effort, muscular tone, reflex irritability, and color—are each assigned a number between 0 and 2. As a result, total scores might be anywhere from 0 to 10, with higher values indicating greater physical health. Among term newborns (born after 37 weeks), a low Apgar score (usually defined as less than 4 or less than 7) is linked to a higher risk of neonatal death <sup>(3)</sup>. The risk of newborn death was recently revealed to be higher among term infants with 5-minute Apgar scores in the normal range (7 to 10) than among those with scores of 7 or 8 <sup>(4)</sup>.

The Apgar score is a recognized and practical way to report a newborn infant's condition right after birth and their reaction to resuscitation, if necessary; nonetheless, it has been improperly used to forecast specific negative neurologic outcomes <sup>(5)</sup>.

The Neonatal Resuscitation Program recommendations specify that the examination should be repeated every 5 minutes for up to 20 minutes if the Apgar score is less than 7 at 5 minutes. However, a score given during resuscitation is not identical to one given to a baby that is breathing on its own <sup>(6)</sup>.

Since many of the factors that affect the Apgar score are changed by resuscitation, there is no established standard for reporting an Apgar score in newborns receiving resuscitation after birth. Although the idea of an aided score that takes resuscitative efforts into account has been floated, the predictive validity has not been

investigated. An enhanced Apgar score report form is recommended to accurately describe these infants, offer accurate documentation, and collect reliable data <sup>(7)</sup>. Numerous elements, such as maternal sedation or anesthesia, congenital defects, gestational age, trauma, and inter-observer variability, might affect the Apgar score. Additionally, before the score is impacted, the metabolic disturbance needs to be considerable <sup>(8)</sup>.

The purposes of this study were to identify the most frequent maternal risk factors associated with low Apgar scores among newborns and evaluate the effect of neonatal sex, type of delivery, mothers' age on Apgar score at Al-Batool Teaching Hospital in Mosul.

## **Patients and Methods:**

### **Study design:**

This study used a cross-sectional Hospital based study design to achieve the study objectives.

### **Study setting:**

The data was collected from 93 maternal and neonatal record files for the mothers who delivered at Al-Batool Teaching Hospital for the period from 1<sup>st</sup> May 31<sup>st</sup> June 2023 with a simple random sampling method was used. All maternity data for women and full-term newborns, regardless of the delivery method, meet the inclusion criteria for this study. Excluded were any maternal files including home births, premature births, or referrals from other facilities following births.

### **Data collection and analysis:**

A document evaluation checklist created by the researchers in English was used to gather data. The data gathering tool's authenticity and dependability were

guaranteed. Obstetric and pediatric specialists with more than five years of experience were asked to evaluate the data collecting tool's clarity, relevance, and simplicity in order to ensure validity. Prior to the actual trial, the instrument was piloted on 10% of the maternal records to make sure that all crucial reliability-related characteristics were addressed. The Statistical Package for the Social Sciences (SPSS) version 26 was used to analyze the data. The maternal factors linked with the Apgar score at 1 and 5 minutes were investigated using the student t-test for independent and dependent two means, with a significance level of 0.05.

### Results:

The current study involved 93 full term neonates; 58 were males and 35 were females with male: female ratio of 1.65:1.


Of them 35 (37.6%) neonates delivered by normal vaginal delivery while the remaining 58 (62.4%) delivered through cesarean section. The mean Apgar score at 1 min was  $5.311 \pm 2.274$  while at 5 min. was  $7.462 \pm 1.891$  and the mean maternal age was  $27.57 \pm 5.942$  years as shown in table (1).

**Table (1): Study characteristics.**

Study characteristics		Frequency	Percentage
Sex	Male	58	62.4
	Female	35	37.6
Types of delivery	NVD	35	37.6
	CS	58	62.4
Apgar score		Mean	Standard deviation
At 1 min		5.311	2.274
At 5 min		7.462	1.891
Maternal age		27.57	5.942

The distribution of the mother according to the maternal risk showed in figure (1), it illustrated that 7 mothers had no risk factors. The commonest risk factors found

was obstructed labour in 21 mothers followed by previous 2 CS in 12 mothers, and meconium stained liquor in 11 mothers.



**Figure (1): The distribution of the mother according to the maternal risk.**

Mean of Apgar at 1 and 5 min according to maternal risk demonstrated in table (2). The table showed that neonates with no maternal risks had mean Apgar score of  $8.285 \pm 0.755$  at 1 min and  $9.571 \pm 0.534$  at 5 min. The risk of obstructed labour was the commonest among the mothers and their neonates had the lowest mean Apgar score of  $3.381 \pm 1.687$  at 1 min and  $6.142 \pm 1.768$  at 5 min.

**Table (2): Mean of Apgar at 1 and 5 min according to maternal risk.**

Maternal variables	Frequency	Apgar At 1 min	Apgar At 5 min
No risk	7	8.285±0.755	9.571±0.534
Obstructed labour	21	3.381±1.687	6.142±1.768
Prev. 2C/S	12	5.750±1.815	7.416±1.781
Meconium stained liquor	11	4.545±2.659	6.818±2.522
PET	8	5.375±2.133	7.375±2.065
Prev. 3C/S	8	5.875±2.416	7.500±1.511
breech	4	6.500±1.732	8.250±1.258
Decrease FM	4	6.000±1.414	8.000±0.816
Polyhydramnios	4	5.000±1.414	7.750±0.500
DM	4	4.250±2.217	6.250±2.362
Rupture membrane >24h	4	5.250±1.258	7.750±0.957
Prev. 1C/S	4	7.25±1.50	8.50±1.00
Prev. 5 C/S	3	4.33±3.055	6.33±2.886
Post date	3	6.333±3.055	9.000±1.732
Maternal anemia	2	5.50±2.121	8.50±0.707
UTI	1	8.0±0.0	9.0±0.0
Antepartum hemorrhage	1	3.0±0.0	9.0±0.0
Hepatitis B	1	8.0±0.0	9.0±0.0
Twin pregnancy	1	4.0±0.0	5.0±0.0
Prev. 4C/S	1	7.0±0.0	10.0±0.0
Cord prolapsed	1	4.0±0.0	8.0±0.0

The comparisons of Apgar score between 1 and 5 min and between sexes demonstrated in table (3) and revealed that mean Apgar score at 1 min was  $5.362 \pm 2.157$  among males and  $5.228 \pm 2.486$  among females while at 5 min was  $7.431 \pm 1.758$  among males and  $7.514 \pm 2.119$  among females. The statistical difference between Apgar score at 1 and 5 min was significant for each of males and females while the difference between males and females was not significant at each of 1 and 5 min.

**Table (3): The comparisons of Apgar score between 1 and 5 min and between sexes.**

Sex	Apgar at 1 min.	Apgar at 5 min.	p-value *
	Mean $\pm$ SD	Mean $\pm$ SD	
<b>Males (n=58)</b>	5.362 $\pm$ 2.157	7.431 $\pm$ 1.758	<b>0.000</b>
<b>Females (n=35)</b>	5.228 $\pm$ 2.486	7.514 $\pm$ 2.119	<b>0.000</b>
<b>p-value **</b>	0.785	0.838	

\* Paired t-test has been used ; \*\*t-test for independent two means

The comparisons of Apgar score between 1 and 5 min and between maternal risks was demonstrated in table (4). The table elicited that mean Apgar score at 5 min was significantly higher than that at 1 min for those with maternal risk (7.290  $\pm$  1.858 vs 5.069  $\pm$  2.184) and those with no maternal risk (9.571 $\pm$  0.534 vs 8.285 $\pm$  0.755). Moreover, the difference between those with maternal risk and those with no maternal risk was statistically significant for both at 1 min (p=0.000) and (p=0.002) and 5 min.

**Table (4): The comparisons of Apgar score between 1 and 5 min and between maternal risks.**

Maternal risk	Apgar at 1 min.	Apgar at 5 min.	p-value *
	Mean $\pm$ SD	Mean $\pm$ SD	
<b>With maternal risk (n=86)</b>	5.069 $\pm$ 2.184	7.290 $\pm$ 1.858	<b>0.000</b>
<b>With no maternal risk (n=7)</b>	8.285 $\pm$ 0.755	9.571 $\pm$ 0.534	<b>0.000</b>
<b>p-value **</b>	<b>0.000</b>	<b>0.002</b>	

\* Paired t-test has been used ; \*\*t-test for independent two means

The comparisons of Apgar score between 1 and 5 min and between types of delivery demonstrated in table (5) and revealed that mean Apgar score at 5 min was significantly higher than that at 1 min for normal vaginal delivery

(7.314±2.068 vs 4.885±2.620) and cesarean section (7.551±1.788 vs 5.569±2.018) while the difference between Apgar score in normal vaginal delivery and cesarean section at each of 1 and 5 min was not significant.

**Table (5): The comparisons of Apgar score between 1 and 5 min and between types of delivery.**

Types of delivery	Apgar at 1 min.	Apgar at 5 min.	p-value *
	Mean ± SD	Mean ± SD	
NVD (n=35)	4.885±2.620	7.314±2.068	<b>0.000</b>
CS (n=58)	5.569±2.018	7.551±1.788	<b>0.000</b>
p-value **	0.162	0.560	

\* Paired t-test has been used ; \*\*t-test for independent two means

The correlation of mothers' ages with Apgar score demonstrated in table (6). The Apgar scores for at both 1 and 5 min were very weakly and insignificantly correlated with the mothers' ages.

**Table (6): The correlation of mothers' ages with Apgar score.**

Pearson's R	<i>r</i>	Asymp. Std. Error <sup>a</sup>	Approx. T <sup>b</sup>	p-value
Apgar at 1 min	0.080	0.101	0.766	0.446 <sup>c</sup>
Apgar at 5 min	0.122	0.097	1.176	0.243 <sup>c</sup>
N of Valid Cases	93			

a. Not assuming the null hypothesis.

b. Using the asymptotic standard error assuming the null hypothesis.

c. Based on normal approximation.

## Discussion:

The mean maternal age for the mothers of neonates included in the present study was 27.57 years with standard deviation of 5.942 years. The correlations of the maternal age with Apgar scores at 1 and 5 min were statistically not significant. A comparable result showed in a study conducted by Lungameni *et al.*,<sup>(9)</sup> in which the mean mothers' age was 29 years and the study showed no significant correlation between the initial Apgar score and age ( $p>0.897$ ). Also Getachew *et al.*,<sup>(10)</sup> study reported that about 75.6% of the mothers were aged 21–34 years. Similarly, in Iran<sup>(11)</sup> and Malaysia<sup>(12)</sup>, the studies reported no connection between low Apgar scores and maternal age groups. Straube *et al.*, 2010<sup>(13)</sup> showed that the maternal age had an influence on Apgar score of newborns. Additionally, Ibrahim *et al.*,<sup>(14)</sup> study from Ethiopia found an association between maternal age with Apgar score.

The current study examined the question whether full term neonates show similar sex-related differences in their 1 and 5 min Apgar scores. The current study found that there were significant improvements at 5 min in comparison to 1 min but there were no significant difference between males and females at both times of Apgar score. In contrary, Nagy *et al.*,<sup>(15)</sup> study reported that the lower Apgar scores was associated more with full-term, male, in comparing to female newborns. Moreover, a large population-based study conducted by Lima *et al.*,<sup>(16)</sup> found that male sex was independently associated with low Apgar score with subsequent neonatal death. The relationship between the male sex and lower Apgar scores thus, may be directly or indirectly related to the sex related gestational differences, instead of a direct sex–Apgar score link.

In the present study, the neonates of mothers with no maternal risks showed normal Apgar scores while those of mothers with different risk showed low Apgar

scores especially at 1 min. Different risks were assessed in this study and the mothers with obstructed labour had lowest Apgar scores of their neonates. Mothers with previous 2 and 3CS, PET, breech, rupture membrane >24h, decrease fetal movement, polyhydramnios, post date, cord prolapsed, Antepartum hemorrhage, and maternal anemia showed Apgar score <7 at 1 min and improved at 5min while those with twin pregnancy, previous 5 CS, DM, and meconium stained liquor, the Apgar score at 5 min still below 7. Furthermore, the current work showed significant differences between those with and without maternal risk at 1 and 5 min in addition to significant difference between Apgar score at 1 and 5 min for each of mothers groups. Similar finding regarding the prolonged obstructed labour was detected by Altman *et al.*, in Sweden <sup>(17)</sup>, by Salustiano *et al.*, in Brazil <sup>(18)</sup>, and by Gudayu in Ethiopia <sup>(19)</sup>. The reason for low Apgar score could be due to fetal distress resulted from prolonged duration of labor. Studies carried out in Ethiopia found that 18.5% of newborns were delivered using meconium-tainted amniotic fluid. The high proportion of newborns delivered through meconium-stained amniotic fluid with either grade 2 (3.1%) or grade 3 (96.9%) meconium can be used to explain the significant association between the meconium grades and the immediate Apgar score found in the study (p0.000). These outcomes were in line with research by Sori <sup>(20)</sup>, who noted that liquor was grade 2 or grade 3 meconium stained in 74.8% of the cases. Other studies from Gonder, Northwest Ethiopia <sup>(21)</sup> and India <sup>(22)</sup> also showed low Apgar scores in these patients. This may be because infants born to women who had amniotic fluid stained with meconium are more likely to aspirate it, filling the lung's tiny airways and alveoli. This may result in restricted lung mobility and airway blockage. A low Apgar score indicating birth asphyxia could then develop as a result of inadequate gas exchange. The two primary causes of cesarean sections linked with an immediate low Apgar score in China were observed to be fetal distress and eclampsia <sup>(23)</sup>. Comparing neonates

whose mothers experienced antepartum hemorrhage to those whose mothers did not, it was discovered that the likelihood of having a poor Apgar score increased by around 3.96 times, a finding that was also made in studies carried out in Addis Ababa, Ethiopia <sup>(24)</sup> and Tigray, Ethiopia <sup>(25)</sup>. Because of the effects of antepartum bleeding, this theory may be plausible.

Lungameni *et al.*, <sup>(9)</sup> study found that diabetes mellitus ( $p>0.298$ ), HIV ( $p>0.442$ ), pregnancy induced hypertension ( $p>0.156$ ), and UTI ( $p>0.140$ ) were not associated with immediate Apgar score. On the opposite, in a study carried out in Turkey <sup>(26)</sup>, an association was found between diabetes mellitus and low Apgar score.

According to Adebami <sup>(27)</sup>, low HB and intrauterine anemia put pregnant women in Nigeria at risk for giving birth to infants with low Apgar scores. According to other studies, mothers who neglected healthy eating habits or had iron deficiency anemia in the second trimester may have caused low nutrients, failing to meet the fetal development's nutritional needs, which resulted in the newborn's low Apgar score.

The outcomes of Lungameni *et al.*'s <sup>(9)</sup> study also show a correlation between poor initial Apgar scores and the following prenatal problems. Similar findings were reported in Brazil by Santos and colleagues <sup>(28)</sup> who found that inadequate vaginal examination could extend the second stage of labor and result in CPD, which is linked to a low Apgar score. Tewesa *et al.* <sup>(29)</sup> did not discover any correlation between APH, CPD, and a poor Apgar score.

According to Sahanak's <sup>(30)</sup> research, newborns whose mothers underwent general anesthesia for a CS had poor Apgar scores. This is because the concentration of anesthetic drugs in the mother's blood might lead to cardiovascular depression, which interferes with placental perfusion and fetoplacental exchange and may lower the Apgar score <sup>(31)</sup>.

While variations between mean Apgar scores at 1 and 5 minutes were significant for both normal vaginal birth and CS, the mean difference between babies from vaginal delivery and cesarean delivery was not significant at either 1 or 5 minutes in this study. Similar result was reported by the study of Rahmanian *et al.*, 2014<sup>(32)</sup> who concluded that difference between mean Apgar score was not significant at 5 minutes of birth from CS and NVD. Moreover, Lungameni *et al.*,<sup>(9)</sup> found significant connection between the types of deliveries and the instant low Apgar score ( $p < 0.000$ ) whether CS or NVD. The relationship between NVD and Apgar score  $< 7$  may be due to actions as, well as, complications throughout childbirth, such as labor duration and breech presentation, which may had impact on choosing method of the delivery<sup>(33-35)</sup>. Opposite to the present study, a study conducted by Paudyal<sup>(36)</sup>, significant difference was found between the mean Apgar score in 1<sup>st</sup> and 5<sup>th</sup> minutes after birth while there is no significant difference between Apgar score among newborns in vaginal delivery and cesarean delivery in 1<sup>st</sup> and 5<sup>th</sup> minutes after birth.

In the present study Apgar score for normal vaginal delivery and CS were below 7 at 1 min and increased to be above 7 at 5 min. Tavares *et al.*,<sup>(37)</sup> study found that along with the factors linked to Apgar score  $< 7$  at 5<sup>th</sup> min, the NVD had a association of threefold times. A further study conducted by Lai *et al.*, 2017<sup>(38)</sup> stated that delivery by cesarean was linked with low to moderate Apgar score than NVD. However, the verification concerning the relationship between mode of delivery and Apgar score is contentious as declared by Khalid *et al.*, 2018<sup>(39)</sup> and Rahmanian *et al.*, 2014<sup>(32)</sup>. A study prepared by Kilsztajn *et al.*, 2007<sup>(40)</sup> found that neonates born through CS had higher Apgar score compared to NVD. Likewise, study done by Fajar *et al.*, 2017<sup>(41)</sup> recommended that neonatal outcomes were better in breech presentation than NVD and found significant

relationship between Apgar score and mode delivery. But some researchers proposed that mode of delivery had no impact on Apgar score as concluded by Burt *et al.*,<sup>(42)</sup> study.

### **Conclusions:**

Mothers were most at risk for obstructed labor, and the mean Apgar scores of their newborns were the lowest. At both 1 and 5 minutes, a significant statistical difference was observed between the groups with and without maternal risk. The difference in the Apgar score that calculated at 1 and 5 minutes for male as, well as, female newborns and between those who delivered vaginally normally and via cesarean section was not statistically significant.

### **References:**

1. Apgar V. A proposal for a new method of evaluation of the newborn infant. *Curr Res Anesth Analg.* 1953; 32: 260-267.
2. Drage JS, Kennedy C, Schwarz BK. The Apgar score as an index of neonatal mortality: a report from the Collaborative Study of Cerebral Palsy. *Obstet Gynecol.* 1964; 24: 222-230.
3. Iliodromiti S, Mackay DF, Smith GCS, Pell JP, Nelson SM. Apgar score and the risk of cause-specific infant mortality: a population-based cohort study. *Lancet.* 2014; 384: 1749-1755.
4. Razaz N, Cnattingius S, Joseph KS. Association between Apgar scores of 7 to 9 and neonatal mortality and morbidity: population based cohort study of term infants in Sweden. *BMJ.* 2019; 365: 11656.
5. Kasdorf E, Laptook A, Azzopardi D, Jacobs S, Perlman JM. Improving infant outcome with a 10 min Apgar of 0. *Arch Dis Child Fetal Neonatal Ed.* 2015 Mar; 100(2): F102-105.

6. American Academy of Pediatrics and American Heart Association. Textbook of Neonatal Resuscitation. 6th edition. Elk Grove Village, IL: American Academy of Pediatrics and American Heart Association; 2011.
7. Lopriore E, vanBurk F, Walther F, Arnout J. Correct use of the Apgar score for resuscitated and intubated newborn babies: questionnaire study. *BMJ*. 2004; 329: 143–144.
8. American College of Obstetrics and Gynecology, Task Force on Neonatal Encephalopathy; American Academy of Pediatrics. Neonatal Encephalopathy and Neurologic Outcome, 2<sup>nd</sup> edition. Washington, DC: American College of Obstetricians and Gynecologists; 2014.
9. Lungameni J, Nghitanwa EM, Uusiku L, & Karera A. Maternal factors associated with immediate low Apgar score in newborn babies at an intermediate hospital in Northern Namibia. *Journal of Public Health in Africa*. 2022; 13(3). <https://doi.org/10.4081/jphia.2022.2045>
10. Getachew B, Etefa T, Asefa A, Terefe B, & Dereje D. Determinants of Low Fifth Minute Apgar Score among Newborn Delivered in Jimma University Medical Center, Southwest Ethiopia. *International Journal of Pediatrics*. 2020. <https://doi.org/10.1155/2020/9896127>
11. Tewesa E, Chirwa E, Majamanda MD, et al. Associative Factors for Birth Asphyxia at Queen Elizabeth Central Hospital, Malawi. *Journal of Biosciences and Medicines*. 2017; 5.
12. Nayeri F, Shariat M, Dalili H, et al. Perinatal risk factors for neonatal asphyxia in Vali-e-Asr hospital, Tehran-Iran. *Iranian Journal of Reproductive Medicine*. 2012: 137-140.
13. Straube S, Voigt M, Jorch G, Hallier E, Briese V and Borchardt U (2010) Investigation of the association of Apgar score with maternal socio-economic and biological factors: an analysis of German perinatal statistics.

- Archives of gynecology and obstetrics 282(2): 135–141. DOI: <https://doi.org/10.1007/s00404-009-1217-7>
14. Ibrahim NA, Muhye A, Abdulie S. Prevalence of Birth Asphyxia and Associated Factors among Neonates Delivered in Dilchora referral Hospital, in Dire Dawa Ethiopia. *Clinics Mother Child Health*. 2017; 14: 4.
  15. Nagy E, Orvos H, Bakki J, Pal A. Sex-differences in Apgar scores for full-term neonates. *Acta Pædiatrica/Acta Pædiatrica*. 2009; 98: 897–900. DOI:10.1111/j.1651-2227.2009.01238.x
  16. Lima JC, Mingarelli AM, Segri NJ, Zavala AA, Takano OA. Population-based study on infant mortality. *Cien Saude Colet*. 2017; 22: 931–939, doi: 10.1590/1413-81232017223.12742016. <https://doi.org/10.1590/1413-81232017223.12742016>
  17. Altman M, Sandström A, Petersson G, Frisell T, Cnattingius S, Stephansson O. Prolonged second stage of labor is associated with low Apgar score. *European Journal of Epidemiology*. 2015; 30(11): 1209–1215. doi: 10.1007/s10654-015-0043-4.
  18. Salustiano EMA, Campos JADB, Ibidi SM, Ruano R, Zugaib M. Baixos índices de Apgar no quinto minuto de vida em população de baixo risco: fatores maternos e obstetricos e resultados pos-natais. *Revista da Associação Médica Brasileira*. 2012; 58(5): 587–593. doi: 10.1590/S0104-42302012000500017.
  19. Gudayu TW. Proportion and factors associated with low fifth minute Apgar score among singleton newborn babies in Gondar University Referral Hospital; North West Ethiopia. *African Health Sciences*. 2017; 17(1): 1–6. doi: 10.4314/ahs.v17i1.2.

20. Sori DA and Belete A Wolde. Meconium Stained Amniotic Fluid: Factors affecting Maternal and Perinatal Outcomes at Jimma University Specialized Teaching Hospital, South West Ethiopia. *Gynecol Obstet*. 2016; 394.
21. Wosenu L, Worku AG, Teshome DF, Gelagay AA. Determinants of birth asphyxia among live birth newborns in University of Gondar referral hospital, northwest Ethiopia: a case-control study. *PLoS ONE*. 2018; 13(9): e0203763.
22. Gane B, Vishnu Bhat B, Rao R, Nandakumar S, Adhisivam B, Joy R, *et al*. Antenatal and intrapartum risk factors for perinatal asphyxia: A case control study. *Curr Pediatr Res*. 2013; 17: 119–122.
23. Li F, Wu T, Lei X, *et al*. The Apgar score and Infant mortality. *PLoS One*. 2013; 8: e69072.
24. Mulugeta T, Sebsibe G, Fenta FA, Sibhat M. Risk factors of perinatal asphyxia among newborns delivered at public hospitals in Addis Ababa, Ethiopia: case–control study. *Pediatr Health Med Ther*. 2020; 11: 297.
25. Tasew H, Zemicheal M, Teklay G, Mariye T, Ayele E. Risk factors of birth asphyxia among newborns in public hospitals of Central Zone, Tigray, Ethiopia 2018. *BMC Res Notes*. 2018; 11(1): 1–7.
26. Kebapcilar L, Kebapcilar AG, Ilhan TT, *et al*. Is the Mean Platelet Volume a Predictive Marker of a Low Apgar Score and Insulin Resistance in Gestational Diabetes Mellitus? A Retrospective Case- Control Study. *Journal of Clinical and Diagnostic Research*. 2016; 10: OC06-OC10.
27. Adebami OJ. Maternal and Foetal determinants of mortality in babies with birth asphyxia at Osogbo, Southwestern Nigeria. *Global Advanced Research Journal of Medicine and Medical Science*. 2015; 270-276.

- 28.Santos NCP, Vogt ES, Duarte DE, *et al.* Factors associated with low Apgar in newborns in birth center. *Rev Bras Enferm.* 2019; 72: 297-304.
- 29.Tewesa E, Chirwa E, Majamanda MD, *et al.* A. Associative Factors for Birth Asphyxia at Queen Elizabeth Central Hospital, Malawi. *Journal of Biosciences and Medicines.* 2017; 5.
- 30.Sahanak S. Comparison of Apgar score in Neonates: spinal versus General Anaesthesia for Elective Caesarean section. *Journal of Evolution Medical and Dental Sciences.* 2014; 539-544.
- 31.Bakhsha F, Yousefi Z, Aryai M, *et al.* Comparison of Apgar score in newborn by vaginal delivery and spinal anesthesia and its relationship with contributing factors. *J Bas Res Med Sci.* 2015; 10-15.
- 32.Rahmanian K, Jahromi AS, Rahmanian V, Ghasvari M and Abari PF. Association of APGAR score with delivery mode in the non-distress newborns. *Online journal of biological sciences.* 2014; 14(1): 21-25. DOI: <https://doi.org/10.3844/ojbssp.2014.21.25>
- 33.Gudayu TW. Proportion and factors associated with low fifth minute Apgar score among singleton newborn babies in Gondar University referral hospital, North West Ethiopia. *Afr Health Sci.* 2017; 17(1): 1–6. doi: 10.4314/ahs.v17i1.2.
- 34.Linder N, Linder I, Fridman E, Kouadio F, Lubin D, Merlob P, *et al.* Birth trauma – risk factors and short-term neonatal outcome. *J Matern Fetal Neonatal Med.* 2013; 26(15): 1491–1495. doi: 10.3109/14767058.2013.789850.
- 35.Duarte G, da Cunha SP, Shimazaki WS, Marcolino AC, Nogueira AA, Berezowski AT, *et al.* Tocotraumatismo com repercussão neonatal. *Rev Bras Ginecol Obstet.* 1997; 19(1): 7–11.

36. Paudyal L. Comparison of APGAR Scores of Newborns with Mode of Delivery and Its Associated Factors. *Int. J. Soc. Sc. Manage.* 2020; 7(3): 176-182. DOI: 10.3126/ijssm.v7i3.29961
37. Tavares VB, Affonso G, Da Rocha ES, Garcia Rodrigues LF, Moraes C, Coelho S, Araújo SS & Gomes C. Factors associated with 5-min APGAR score, death and survival in neonatal intensive care: A case-control study. *BMC Pediatrics.* 2022; 22. <https://doi.org/10.1186/s12887-022-03592-9>
38. Lai S, Flatley C, Kumar S. Perinatal risk factors for low and moderate five-minute Apgar scores at term. *Eur J Obstet Gynecol Reprod Biol.* 2017; 210: 251-256. DOI: <https://doi.org/10.1016/j.ejogrb.2017.01.008>
39. Khalid MA, Ghani R, Khalid MF, Malik MS, Waqas A. Association of delivery procedure with APGAR scores among neonates born to healthy Pakistani mothers: a pilot study. *F1000Research.* 2018; 7(346): 346. DOI: <https://doi.org/10.12688/f1000research.13784.1>
40. Kilsztajn S, Lopes EDS, Carmo MSND and Reyes AMDA. APGAR score associated with mode of delivery in Sao Paulo State, Brazil. *Cad.Saude Publica.* 2007; 23: 1886-1892. PMID: 17653406.
41. Fajar JK, Andalas M and Harapan H. Comparison of Apgar scores in breech presentations between vaginal and cesarean delivery. *Ci ji yi xue za zhi = Tzu-chi medical journal.* 2017; 29(1): 24-29. DOI: [https://doi.org/10.4103/tcmj.tcmj\\_5\\_17](https://doi.org/10.4103/tcmj.tcmj_5_17)
42. Burt, Richard, Vaughan T and Daling J. Evaluating the risks of cesarean section: Low Apgar score in repeat C-section and vaginal deliveries. *American journal of public health.* 1988; 78: 1312-1314. DOI: <https://doi.org/10.2105/AJPH.78.10.1312>