

# Influence of nutrient management through use of bio-organic manures on bio-chemical attributes of Aonla (*Emblca officinalis* Gaertn.) cv. Na-10

## ABSTRACT

The present investigation was conducted to evaluate influence of nutrient management using bio-organic manure on bio-chemicals in Aonla. The research was carried at Main Experiment Station, Horticulture, Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya (U.P.) during the year 2021. It was laid out in a Randomized Block Design with 10 treatments namely: T<sub>1</sub> – Control, T<sub>2</sub>- RDF 100% (1kg.N: 0.5kg.P:1kg.K per tree), T<sub>3</sub>- FYM (10kg./tree) + RDF 100%, T<sub>4</sub>- Poultry Manure (7.5kg./tree) + RDF 100%, T<sub>5</sub>- FYM (10kg./tree) + RDF 50% + Azospirillum (10ml./tree), T<sub>6</sub>- Poultry Manure (7.5kg/tree) + RDF 50% + Azospirillum (10ml./tree), T<sub>7</sub> -FYM (10kg./tree) + RDF 50% + PSB (10ml./tree), T<sub>8</sub>- Poultry Manure + RDF 50% + Azospirillum (10ml./tree), T<sub>9</sub> -FYM (10kg/tree) + RDF 50% + Azospirillum (10ml./tree) + PSB (10ml./tree) and T<sub>10</sub>- Poultry Manure (7.5kg/tree) + RDF 50% + Azospirillum (10ml./tree) + PSB (10ml./tree). The experiment was replicated thrice. Results showed treatment T<sub>10</sub>- Poultry Manure (7.5kg/tree) + RDF 50% + Azospirillum (10ml/tree) + PSB (10ml./tree) outperformed the rest with maximum TSS (11.83<sup>o</sup> Brix), Ascorbic acid content ( 599.98 mg/ 100g fruit pulp), reducing sugar (3.28%), non-reducing sugar (3.00%), total sugar (6.29%) and minimum acidity (1.32%). The treatment combination is therefore recommended for application to Aonla trees in Eastern Uttar Pradesh in order to obtain high yields with better quality fruits.

**Keywords:** Aonla, *Azospirillum*, Nutrient management, Poultry Manure

## INTRODUCTION

The Indian gooseberry or aonla (*Emblca officinalis* Gaerten) belongs to family “Euphorbiaceae” with the chromosome number 2n=28. Aonla also is known in different names in different region like Amla, Amolphal, Amalakamu, Dhatri, Nelli, Usirika and Maryobalan (**Sharma and Nagaich (2022)**). It is native to Indo- China, particularly in central and southern India. In India Aonla cultivation is done mainly in northwest Himalayas to eastern Himalaya. The domestication of Aonla was first started in Varanasi, Uttar Pradesh with the initiative of Maharaja of Kashi. Banarasi, a superior genotype was selected from the wild Aonla trees are available in large number in the nearby Vindhyan hills. Authentic information regarding its cultivation dates back to 1881-82 in the Pratapgarh district of Uttar Pradesh (**Singh et al., 2019**). It occupies an area of 100 thousand hectares with a production of 1206 thousand MT (**Anonymous, 2021-22**).

In Uttar Pradesh, Aonla was more cultivated in nearby belt of Pratapgarh followed by Ayodhya district. Area under Aonla orchard in Pratapgarh district is about 1300 hectares. Whereas, the area in Sadder block of district of Pratapgarh is approximately 3250 hectares (**Rai et al., 2017**). Aonla is a

subtropical plant and prefers dry subtropical climate but it can be successfully cultivated in wide range of soils and climatic conditions. Owing to its hardy nature, suitability to various wastelands, high productivity, nutritive and therapeutic value Aonla has become an important fruit. Aonla is a medium sized, much-branched tree occupying height of 10-20 m. Inflorescence is racemose type, flower minute, unisexual with short pedicel. Fruit depressed round, globose or oblate, indented at the base. It is richest sources of vitamin C (400-1300mg./100g fruit pulp) among the fruits next to Barbados cherry. (Tripathi *et al.*, 2022). Sustained nutrient management is the passport to enter into the 21<sup>st</sup> century. In view of these situations, the application of nutrients from the combinations of inorganic, organic and biological sources not only fulfills the nutrient requirement of the crop but also improve the soil health. Hence integrated nutrient management practices will help to increase the productivity of the crop and enrich the soil (Sharma *et al.*, 2018). Organic manures supply plant nutrients and micronutrients. They improve soil physical properties like soil structure, infiltration, porosity, water holding capacity, bulk density etc. Organic manures act as buffering agents and supplies food for beneficial living organism (Aal *et al.*, 2020). [Include the objective please!!!!](#)

## **MATERIALS AND METHODS**

The present investigation was carried out at Main Experiment Station, Horticulture, Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya (U.P.) during the year 2021-2022. The experiment was laid out in Randomized Block Design with 10 treatments namely: T<sub>1</sub>Control, T<sub>2</sub> RDF 100% (1kg.N: 0.5kg.P:1kg.K per tree), T<sub>3</sub> FYM (10kg./tree) + RDF 100%, T<sub>4</sub> Poultry Manure (7.5kg./tree) + RDF 100%, T<sub>5</sub> FYM (10kg./tree) + RDF50% + *Azospirillum* (10ml./tree), T<sub>6</sub> Poultry Manure (7.5kg./tree) + RDF 50% + *Azospirillum* (10ml./tree), T<sub>7</sub> FYM (10kg./tree) + RDF 50% + PSB (10ml./tree), T<sub>8</sub> Poultry Manure (7.5kg./tree) + RDF 50% + PSB (10ml./tree), T<sub>9</sub> FYM (10kg./tree) + RDF 50% + *Azospirillum* (10ml./tree) + PSB (10ml./tree), and T<sub>10</sub> Poultry Manure (7.5kg./tree) + RDF 50% + *Azospirillum* + (10ml./tree) + PSB (10ml./tree) replicated thrice. The prescribed schedule for the Aonla plantation was followed for the usual cultural operations, plant protection measures, and basal application of manures and fertilizers. Data was collect on Total Soluble Solids, ascorbic acid, acidity, reducing sugar, non-reducing sugar and total sugar including numerous other fruit characteristics. TSS was calculated using a hand refractometer and displayed in °Brix. By using the standard procedure outlined in AOAC (1990), several chemical parameters, including total sugars, ascorbic acid, and acidity were measured. The data obtained during experimentation were statistically analyzed as per the method given by Panse and Sukhatme (1985).

**Comment [H1]:** Include the following information for the site:

1. Soil type and its properties
2. Weather conditions
3. Age of the Aonla trees
4. Organoleptic test results

## RESULTS AND DISCUSSION

### Total Soluble Solid (<sup>0</sup>brix)

Results (Table 1) showed significant difference on total soluble solids of fruits influenced by various treatments. Treatment T<sub>10</sub>- Poultry Manure (7.5kg/tree) + RDF 50% + *Azospirillum* (10ml/tree) +PSB (10ml/tree) recorded maximum total soluble solids value (11.82<sup>0</sup>Brix) followed by T<sub>9</sub>- FYM (10kg/tree) + RDF 50% + *Azospirillum* (10ml/tree) +PSB (10ml/tree). The lowest total soluble solid (8.82 <sup>0</sup>brix) was observed in the control treatment. It is assumed that TSS content increased with *Azospirillum* and FYM application which contributed to the quick metabolic transformation of starch and pectin into soluble compounds and rapid translocation of sugars from leaves to the developing fruits leading to conversion of complex polysaccharides into simple sugars with reference to Kumar *et al.* (2018) who reported that application of 100% recommended dose of fertilizers (RDF) along with vermicompost + poultry manure + *Azospirillum* + PSB recorded maximum fruit TSS. The findings are also in conformity with Babiskar *et al.* (2011), Sharma *et al.* (2022) in kiwifruit and Jaiswal *et al.* (2023) in guava.

Table 1: Shows TSS, ascorbic acid, acids, reducing and non sugars and total sugars in Aonla fruit

	Treatments	TSS ( <sup>0</sup> Brix)	Ascorbic acid (mg/100g fruit pulp)	Acidity (%)	Reducing sugar (%)	Non-reducing sugar (%)	Total sugars (%)
T <sub>1</sub>	Control	8.82	488.16	2.01	2.10	1.9	4.00
T <sub>2</sub>	RDF 100 % (1kg N : 0.5kg P : 1kg K per tree)	9.40	499.58	1.97	2.39	2.17	4.57
T <sub>3</sub>	FYM (10kg/tree) + RDF 100%	9.48	509.51	1.85	2.65	2.25	4.91
T <sub>4</sub>	Poultry manure (7.5kg/tree) + RDF 100%	9.54	512.16	1.79	2.76	2.30	5.06
T <sub>5</sub>	FYM (10kg/tree) + RDF50% + <i>Azospirillum</i> (10ml/tree)	9.69	529.50	1.73	2.88	2.39	5.27
T <sub>6</sub>	Poultry manure (7.5kg/tree) + RDF 50% + <i>Azospirillum</i> (10ml/tree)	10.25	536.90	1.66	2.92	2.52	5.45

T <sub>7</sub>	FYM (10kg/tree) + RDF 50% + PSB (10ml/tree)	10.50	555.37	1.55	3.10	2.62	5.72
T <sub>8</sub>	Poultry manure + RDF 50% + PSB (10ml/tree)	10.58	569.63	1.49	3.16	2.88	6.04
T <sub>9</sub>	FYM + RDF 50% + <i>Azospirillum</i> (10ml/tree) + PSB (10ml/tree)	10.66	596.01	1.40	3.25	2.92	6.18
T <sub>10</sub>	Poultry manure (7.5kg/tree) + RDF 50% + <i>Azospirillum</i> (10ml/tree) + PSB (10ml/tree)	11.83	599.98	1.32	3.28	3.00	6.29
	SEm ±	0.47	1.35	0.01	0.03	0.04	0.05
	CD at 5%	1.41	4.02	0.04	0.08	0.12	0.14

**Comment [H2]:** The statistics is not coming out clear on whether significant or not.

### Ascorbic acid (mg./100g fruit pulp) and Acidity (%)

The highest ascorbic acid value 599.98 in Table 1 was recorded in treatment T<sub>10</sub> Poultry Manure (7kg/tree) + RDF 50% + *Azospirillum* (10ml/tree) + PSB (10ml/tree), followed by treatment T<sub>9</sub>- FYM (10kg/tree) + RDF 50% + *Azospirillum* (10ml/tree) +PSB (10ml/tree). However the lowest ascorbic acid 488.16 was observed in the control with treatment The results indicated that the application of FYM, *Azospirillum* and PSB had significant influence on the content of ascorbic acid in the fruit. These results are in conformity with the findings of Yadav *et al.* (2007), Bohane *et al.* (2014), Kour *et al.* (2019), Sharma *et al.* (2022). Treatments T<sub>10</sub>- Poultry Manure (7.5 kg/tree) + RDF 50% + *Azospirillum* (10ml/tree) + PSB (10ml/tree) and T<sub>9</sub> FYM (10kg/tree) + RDF 50% + *Azospirillum* (10ml/tree) reduced acidity percent and obtained minimum fruit acidity value (01.32%) followed by treatment T<sub>9</sub>. The decrease in acidity of fruits may be attributed to their conversion into sugars. Similar findings were also reported by Bohane *et al.* (2014), Jamra *et al.* (2018), Athani (2009), Yadav *et al.* (2007) Sharma *et al.* (2022) and Jaiswal *et al.* (2023) in guava.

### Reducing and Non-reducing Sugar (%)

Results (Table 1) have shown significant maximum reducing sugar (3.28 %) with application of Poultry Manure (7.5 kg./tree) + RDF 50% + *Azospirillum* (10ml./tree) +PSB (10ml./tree) followed by FYM (10kg/tree) + RDF 50% + *Azospirillum* (10ml./tree). However minimum reducing sugar

(2.10%) was recorded in control treatment T<sub>1</sub>. An increase in reducing sugars contents with *Azospirillum* and Farm Yard Manure application could be attributed to the quick metabolic transformation of starch and pectin into soluble compounds and rapid translocation of sugars from leaves to the developing fruits as well as the conversion of complex polysaccharides into simple sugars. Similar results were noted by Jamra *et al.* (2018), Kour *et al.* (2019) in Aonla, Verma *et al.* (2014) in phalsa, Rai *et al.* (2009) in Pear and Chawla *et al.* (2020). The application of Poultry Manure (7.5 kg/tree) + RDF 50% *Azospirillum* + (10ml/tree) +PSB (10ml/tree) (Table 1) showed significant maximum non- reducing sugars (3.0% ) followed by FYM (10kg/tree) + RDF 50% + *Azospirillum* (10ml/tree) PSB (10ml/tree). While, minimum non-reducing sugar (1.9%) was observed in the control treatment T<sub>1</sub>. The same reason could be applied that the quick metabolic transformation of starch and pectin into soluble compounds and rapid translocation of sugars from leaves to the developing fruits. The high non-reducing sugar (2.52%) were also recorded with application of Poultry Manure (7.5 kg/tree) + RDF 50% + *Azospirillum* (10ml/tree) + PSB (10ml/tree). The research findings were supported by Verma *et al.* (2014) in Phalsa, Chawla *et al.* (2020), Jamra *et al.* (2018) and Kour *et al.* (2019) in Aonla.

#### **Total Sugar (%):**

In Table 1 the highest total sugars value (6.29%) were recorded in T<sub>10</sub> treatment that is Poultry Manure (7.5 kg/tree) + RDF 50% + *Azospirillum* (10ml/tree) +PSB (10ml/tree) followed by T<sub>9</sub> treatment FYM (10kg/tree) + RDF 50% + *Azospirillum* (10ml/tree) and T<sub>8</sub> were proved equally good with T<sub>10</sub>. However minimum total sugar percent (4.00%) was recorded in control treatment T<sub>1</sub>. The findings are corroborating with those reported by Singh *et al.* (2008), Sharma *et al.* (2022), Kour *et al.* (2019) in aonla fruits, Verma *et al.* (2014) and Yadav *et al.* (2008) in Phalsa.

#### **CONCLUSION:**

Based on the results of this present investigation, it can be concluded that treatment T<sub>10</sub> (Poultry Manure (7.5kg/ tree) + RDF 50% + *Azospirillum* (10ml/tree) + PSB (10ml/tree) produced the maximum TSS (11.83<sup>0</sup> Brix), Ascorbic acid content ( 599.98 mg/ 100g fruit pulp), reducing sugar (3.28%), non-reducing sugar (3.00%), total sugar (6.29%) and minimum acidity (1.32%) of aonla which was at par with T<sub>9</sub>-FYM (10kg/tree) + RDF 50% + *Azospirillum* (10ml./tree) + PSB (10ml./tree) can be recommended to aonla growers in eastern Uttar Pradesh for obtaining higher yield with better quality fruits.

## FUTURE SCOPE

Integrated nutrient management, developed on the principles of eco-friendly and efficient balanced fertilizer and based in optimization of nutrient supplies from all the available sources, inorganic and organic, which lowers production costs and increases productivity, were the greatest ways to cultivate Aonla.

## REFERENCE

**Aal, J.M.; Patel, K. M.; Patel, S.J.; and Sindha, D. J. (2020).** Effect of integrated nutrient management on fruit yield of aonla (*Emblica officinalis* Gaertn) cv. Gujarat Aonla-1, *International Journal of Current Microbiology and Applied Sciences*, 9(10): 417-423.

[Anonymous \(2021-22\), Indian Horticulture Database, National Horticulture Board, Department of Agriculture and Co-operation & Farmers Welfare, Government of India, Gurugram.](#)

**A.O.A.C. 1990.** Official methods of analysis of 16th edition. Association of official Agricultural Chemists, Washington D.C.

~~[Anonymous \(2021-22\), Indian Horticulture Database, National Horticulture Board, Department of Agriculture and Co-operation & Farmers Welfare, Government of India, Gurugram.](#)~~

**Athani, S.I.; Ustad, A.I.; Prabhuraj, H.S.; Swamy, G.S.K.; Patil, P.B. and Kotikal, Y.K. (2009).** Influence of vermicompost on growth, fruit yield and quality of guava cv. Sardar. *Acta Horticulture*, 735(2): 381-85.

**Bhabiskar, M.N.; Bharad, S.G.; Dod, V.N. and Barne, V.G. (2011).** Effect of integrated nutrient management on yield and quality of sapota. *Plant Archives*, 11(2):661-663.

**Bohane L, Tiwari R. (2014).** Effect of integrated nutrient management on physicochemical parameters of ber under Malwa plateau conditions. *Annals of Plant and Soil Research*, 16(4):346-348.

- Chawla, R., and Sadawarti, R. K. (2020).** Effect of bio-fertilizers and organic manures on growth, yield and fruit quality of fruit crops. *Plant Arch*, 20(1): 3767-3768.
- Jaiswal, A., Prasad, V. M., Bhadur, V., and Kumar, N. (2023).** Effect of organic manure and inorganic fertilizer on growth, yield and quality of guava (*Psidium guajava* L.), *The Pharma Innovation Journal*, 12(2): 2133-2136.
- Jamra, R., Kanwar, J., Dubey, R. and Tiwari, R. (2018).** Effect of integrated nutrient management practices on physical, biochemical properties and yield of aonla (*Embllica officinalis* Gaertn). *International Journal of Chemical Studies*, 6(2): 1256-1258.
- Kour, D., Wali, V. K., Bakshi, P., Bhat, D. J., Sharma, B. C., Sharma, V., and Sinha, B. K. (2019).** Effect of Organic and Inorganic fertilizers along with Azotobacter on Growth, Yield and Quality of Aonla (*Embllica officinalis* gaertn.) Cv. Na-7. *International Journal of Current Microbiology and Applied Sciences*, 8(9): 1142-1151.
- Kumar, A., Prasad, V.M., Singh, D., Bahadur, V., David, A.A. and Beer, K. (2018).** Effect of bio-fertilizers, vermicompost and Trichoderma on yield and economics of strawberry (*Fragaria x annanasa* Duch.) cv. Sweet Charlie. *International Journal of Current Microbiology and Applied Sciences*, 7(6): 1534-1538.
- Panse, V.C. and Sukhatme, P.V. (1985).** Statistical Methods for Agricultural Workers, ICAR, New Delhi.
- Rai, J.; Singh, S.P. and Singh, A.K. (2017).** Economics of aonla production in district Pratapgarh (U.P.). *International Research Journal of Agricultural Economics and Statistics*, 8(2): 426-430.
- Rai, P.N., Singh, P.N. and Binayak, C. (2009).** Effect of different sources of organic manures on yield and quality of low-chill pear (*Pyrus pyrifolia*) cv. Gola. *Progressive Horticulture*, 41(1): 103-106.
- Sharma, A. and Nagaich, K. (2022).** Biofertilizer—An Effective Component of Integrated Nutrient Management in Aonla. *Annals of Biology*, 38(1): 83-87.
- Sharma, A. and Nagaich, K. (2022).** Biofertilizer—An Effective Component of Integrated Nutrient Management in Aonla. *Annals of Biology*, 38(1): 77-82.

- Sharma, B.D., Meena Anita, Jatav, M.K. and Balai, R.C. (2018).** Integrated nutrient management for horticultural crops in arid region. ICAR -CIAH/ Tech./ Pub. No.77, pp 42.
- Sharma, S., Rana, V. S., Rana, N., Sharma, U., Gudeta, K., Alharbi, K., and Bhat, S. A. (2022).** Effect of Organic Manures on Growth, Yield, Leaf Nutrient Uptake and Soil Properties of Kiwifruit (*Actinidia deliciosa* Chev.) cv. Allison. *Plant Archives*, 11(23): 3354.
- Singh, A.; Yadav, A.L., Yadav, D.K. and Mishra, S.K. (2008).** Effect of integrated nutrient management on yield and quality of aonla (*Emblica officinalis* Gaertn.) cv. NA-10. *Plant Archives*, 8(1): 473- 474.
- Singh, A. K., Singh, S. A. N. J. A. Y., Saroj, P. L., Mishra, D. S., Singh, P. P., and Singh, R. K. (2019).** Aonla (*Emblica officinalis*) in India: a review of its improvement, production and diversified uses. *Indian Journal of Agriculture Sciences*, 89(11): 1773-1781.
- Tripathi, V. K., and Nand, V. (2022).** Effect of foliar application of Boron, Zinc and NAA on fruit retention, yield and quality attributes of Aonla. *Progressive Horticulture*, 54(1): 76-81.
- Verma R.S.; Singh HK, Verma SS. (2014).** Effect of integrated nutrient management on plant growth, fruit yield and quality of phalsa (*Grewia subinaequalis* D.C.). *Asian Journal of Horticulture*, 9(1):48-52.
- Yadav, D.K.; Pathak, Sanjay and Yadav, A.L. (2008).** Effect of integrated nutrient management on physico-chemical analysis of Phalsa (*Grewia subinaequalis* D.C.). *Plant Archives*, 8(1): 461-463.
- Yadav, Rajesh; Singh, H.K. and Yadav; A.L. (2007).** Effect of integrated nutrient management on productivity and quality of aonla (*Emblica officinalis* Gaertn). *Plant Archives* 7(2): 881- 883.