

# Influence of nutrient management through use of bio-organic manures on bio-chemical attributes of Aonla (*Emblia officinalis* Gaertn.) cv. Na-10

## ABSTRACT

The present investigation was conducted to evaluate influence of nutrient management using bio-organic manure on bio-chemicals in Aonla. The research was carried at Main Experiment Station, Horticulture, Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya (U.P.) during the year 2021. It was laid out in a Randomized Block Design with 10 treatments namely: T<sub>1</sub> – Control, T<sub>2</sub>- RDF 100% (1kg.N: 0.5kg.P:1kg.K per tree), T<sub>3</sub>- FYM (10kg./tree) + RDF 100%, T<sub>4</sub>- Poultry Manure (7.5kg./tree) + RDF 100%, T<sub>5</sub>- FYM (10kg./tree) + RDF 50% + Azospirillum (10ml./tree), T<sub>6</sub>- Poultry Manure (7.5kg/tree) + RDF 50% + Azospirillum (10ml./tree), T<sub>7</sub> -FYM (10kg./tree) + RDF 50% + PSB (10ml./tree), T<sub>8</sub>- Poultry Manure + RDF 50% + Azospirillum (10ml./tree), T<sub>9</sub> –FYM (10kg/tree) + RDF 50% + Azospirillum (10ml./tree) + PSB (10ml./tree) and T<sub>10</sub>- Poultry Manure (7.5kg/tree) + RDF 50% + Azospirillum (10ml./tree) + PSB (10ml./tree). The experiment was replicated thrice. Results showed treatment T<sub>10</sub>- Poultry Manure (7.5kg/tree) + RDF 50% + Azospirillum (10ml./tree) + PSB (10ml./tree) outperformed the rest with maximum TSS (11.83<sup>0</sup> Brix), Ascorbic acid content ( 599.98 mg/100gfruitpulp), reducing sugar (3.28%), non-reducing sugar (3.00%), total sugar (6.29%) and minimum acidity (1.32%). The treatment combination is therefore recommended for application to Aonla trees in Eastern Uttar Pradesh in order to obtain high yields with better quality fruits.

**Keywords:** Aonla, *Azospirillum*, Nutrient management, Poultry Manure

## INTRODUCTION

The Indian gooseberry or aonla (*Emblia officinalis* Gaerten) belongs to family “Euphorbiaceae” with the chromosome number 2n=28. Aonla also is known in different names in different region like Amla, Amolphal, Amalakamu, Dhatri, Nelli, Usirika and Maryobalan (Sharma and Nagaich (2022)). It is native to Indo-China, particularly in central and southern India. In India Aonla cultivation is done mainly in north west Himalaya to eastern Himalaya. The domestication of Aonla was first started in Varanasi, Uttar Pradesh with the initiative of Maharaja of Kashi. Banarasi, a superior genotype was selected from the wild Aonla trees available in large number in the nearby Vindhyan hills. Authentic information regarding its cultivation dates back to 1881-82 in the Pratapgarh district of Uttar Pradesh (Singh et al., 2019). It occupies an area of 100 thousand hectares with a production of 1206 thousand MT (Anonymous, 2021-22).

In Uttar Pradesh, Aonla was more cultivated in nearby belt of Pratapgarh followed by Ayodhya district. Area under Aonla orchard in Pratapgarh district is about 1300 hectares. Whereas, the area in Sadder block of district of Pratapgarh is approximately 3250 hectares (Rai et al., 2017). Aonla is a subtropical plant and prefers dry subtropical climate but it

can be successfully cultivated in wider range of soils and climatic conditions. Owing to its hardy nature, suitability to various wastelands, high productivity, nutritive and therapeutic value Aonla has become an important fruit. Aonla is a medium sized, much-branched tree occupying height of 10-20 m. Inflorescence is racemose type, flower minute, unisexual with short pedicel. Fruit depressed round, globose or oblate, indented at the base. It is richest sources of vitamin C (400-1300mg./100g fruit pulp) among the fruits next to Barbados cherry. (Tripathi *et al.*, 2022). Sustained nutrient management is the passport to enter into the 21<sup>st</sup> century. In view of these situations, the application of nutrients from the combinations of inorganic, organic and biological sources not only fulfills the nutrient requirement of the crop but also improve the soil health. Hence integrated nutrient management practices will help to increase the productivity of the crop and enrich the soil (Sharma *et al.*, 2018). Organic manures supply plant nutrients and micronutrients. They improve soil physical properties like soil structure, infiltration, porosity, water holding capacity, bulk density etc. Organic manures act as buffering agents and supplies food for beneficial living organism (Aal *et al.*, 2020). The present investigation will be undertaken to generate the sufficient information with the following objectives.

1. To find out the effect of INM on physical attributes of aonla fruits,
2. To assess the effect of INM on chemical attributes of aonla fruits and
3. To work out the economic feasibility of the various treatment

## **MATERIALS AND METHODS**

The present investigation was carried out at Main Experiment Station, Horticulture, Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya (U.P.) during the year 2021-2022. The experiment was laid out in Randomized Block Design with 10 treatments namely: **T<sub>1</sub>** Control, **T<sub>2</sub>** RDF100% (1kg.N:0.5kg.P:1kg.K per tree), **T<sub>3</sub>** FYM (10kg./tree)+RDF100%, **T<sub>4</sub>** Poultry Manure (7.5kg./tree)+RDF100%, **T<sub>5</sub>** FYM (10kg./tree)+RDF50%+*Azospirillum* (10ml./tree), **T<sub>6</sub>** Poultry Manure (7.5kg/tree)+RDF50%+*Azospirillum* (10ml./tree), **T<sub>7</sub>** FYM (10kg./tree)+RDF50%+PSB (10ml./tree), **T<sub>8</sub>** Poultry Manure (7.5kg/tree)+RDF50%+PSB (10ml./tree), **T<sub>9</sub>** FYM (10kg/tree)+RDF50%+*Azospirillum* (10ml./tree)+PSB (10ml./tree), and **T<sub>10</sub>** Poultry Manure (7.5kg/tree)+RDF50%+*Azospirillum* (10ml./tree)+PSB (10ml./tree) replicated thrice. The Ayodhya district's climate is classified as semi-arid, with three distinct seasons: rainy or wet, winter, and summer or hot. The rainy season begins the last week of June and lasts until September or even into October, with 1200 mm of rain on average. The soil was identified as a

sandy loam with an average pH of 7.71 and an average proportion of fine sand (64.77%), silt (22.76%), and clay (14.95%). Thirty-six year old plants were used in the experiment.

The prescribed schedule for the Aonla plantation was followed for the usual cultural operations, plant protection measures, and basal application of manures and fertilizers. Data was collect on Total Soluble Solids, ascorbic acid, acidity, reducing sugar, non-reducing sugar and total sugar including numerous other fruit characteristics. TSS was calculated using a hand refractometer and displayed in <sup>0</sup>Brix. By using the standard procedure outlined in AOAC (1990), several chemical parameters including total sugars, ascorbic acid, and acidity were measured. Thedataobtainedduring experimentation werestatistically analyzedas perthe method given by **Panse and Sukhatme (1985)**.

## RESULTS AND DISCUSSION

### TotalSolubleSolid(<sup>0</sup>brix)

Results (Table 1) showed significant differenceontotalsolublesolidsoffruitsinfluencedby various treatments.

Treatment T<sub>10</sub>-Poultry

Manure(7.5kg/tree)+RDF50%+Azospirillum(10ml/tree)+PSB(10ml/tree)recordedmaximum total soluble solids value (11.82<sup>0</sup>Brix) followed by T<sub>9</sub>- FYM (10kg/tree) +RDF 50% +Azospirillum (10ml/tree) +PSB (10ml/tree).The lowesttotal solublesolid(8.82<sup>0</sup>brix)wasobservedinthe control treatment. It is assumed that TSScontentincreasedwith AzospirillumandFYMapplicationwhich contributed to the quick metabolic transformation of starch and pectin into solublecompoundsandrapidtranslocation of sugarsfromleavestothedevelopingfruits leading toconversionofcomplexpolysaccharidesintosimplesugars with reference to**Kumaretal.(2018)who** reported that application of 100% recommended dose of fertilizers (RDF) along withvermicompost+poultrymanure+Azospirillum+PSBrecordedmaximumfruitTSS.Thefindings arealsoin conformitywith**Babiskaretal.(2011)**, **Sharmaet al. (2022)** in kiwifruit and**Jaiswal et al. (2023)** in guava.

Table 1: Shows TSS, ascorbic acid, acids, reducing and non sugars and total sugars in Aonla fruit

	Treatments	TSS( <sup>0</sup> Brix)	Ascorbic acid(mg/100gfruit pulp)	Acidity(%)	Reducing sugar(%)	Non-reducing sugar(%)	Totalsugars (%)
T <sub>1</sub>	Control	8.82	488.16	2.01	2.10	1.9	4.00
T <sub>2</sub>	RDF100%(1kgN:0.5kgP:1kgKpertr	9.40	499.58	1.97	2.39	2.17	4.57

	ee)						
T <sub>3</sub>	FYM(10kg/tree)+RDF100%	9.48	509.51	1.85	2.65	2.25	4.91
T <sub>4</sub>	Poultry manure(7.5kg/tree)+RDF100%	9.54	512.16	1.79	2.76	2.30	5.06
T <sub>5</sub>	FYM(10kg/tree)+RDF50%+Azospirillum(10ml/tree)	9.69	529.50	1.73	2.88	2.39	5.27
T <sub>6</sub>	Poultrymanure(7.5kg/tree)+RDF50%+Azospirillum(10ml/tree)	10.25	536.90	1.66	2.92	2.52	5.45
T <sub>7</sub>	FYM(10kg/tree)+RDF50%+PSB(10ml/tree)	10.50	555.37	1.55	3.10	2.62	5.72
T <sub>8</sub>	Poultry manure+RDF50%+PSB(10ml/tree)	10.58	569.63	1.49	3.16	2.88	6.04
T <sub>9</sub>	FYM+RDF50%+Azospirillum(10ml/tree)+PSB(10ml/tree)	10.66	596.01	1.40	3.25	2.92	6.18
T <sub>10</sub>	Poultry manure (7.5kg/tree)+RDF50%+Azospirillum(10ml/tree)+PSB(10ml/tree)	11.83	599.98	1.32	3.28	3.00	6.29
	<b>SEm±</b>	0.47	1.35	<b>0.01</b>	0.03	0.04	0.05
	<b>CD at 5%</b>	1.41	4.02	<b>0.04</b>	0.08	0.12	0.14

### Ascorbic acid(mg./100fruitpulp)and Acidity(%)

The highest ascorbic acid value 599.98 in Table 1 was recorded in treatment T<sub>10</sub> Poultry Manure(7kg/tree)+RDF50%+Azospirillum (10ml/tree) + PSB (10ml/tree), followed by treatment T<sub>9</sub>- FYM(10kg/tree) + RDF 50% + Azospirillum (10ml/tree) +PSB (10ml/tree). However the lowest ascorbic acid 488.16 was observed in the control treatment. The results indicated that the application of FYM, Azospirillum and PSB had significant influence on the content of ascorbic acid in the fruit. These results are in conformity with the findings of Yadav *et al.* (2007), Bohane *et al.* (2014), Kour *et al.* (2019), Sharma *et al.* (2022). Treatments T<sub>10</sub>- Poultry Manure(7.5kg/tree)+RDF50%+Azospirillum(10ml/tree)+PSB(10ml/tree) and T<sub>9</sub> FYM (10kg/tree) +RDF50%+Azospirillum(10ml/tree) reduced acidity percent and obtained minimum fruit acidity value (01.32%) followed by treatment T<sub>9</sub>. The decrease in acidity of fruits may be attributed to their conversion into sugars. Similar findings were also reported by Bohane *et al.* (2014),

**Jamra et al. (2018), Athani (2009), Yadav et al. (2007) Sharma et al. (2022) and Jaiswal et al. (2023)** in guava.

### **Reducing and Non-reducing Sugar (%)**

Results (Table 1) have shown significant maximum reducing sugar (3.28%) with application of Poultry Manure (7.5 kg./tree) + RDF 50% + *Azospirillum* (10 ml./tree) + PSB (10 ml./tree) followed by FYM (10 kg/tree) + RDF 50% + *Azospirillum* (10 ml./tree). However minimum reducing sugar (2.10%) was recorded in control treatment T<sub>1</sub>. An increase in reducing sugars contents with *Azospirillum* and Farm Yard Manure application could be attributed to the quick metabolic transformation of starch and pectin into soluble compounds and rapid translocation of sugars from leaves to the developing fruits as well as the conversion of complex polysaccharides into simple sugars. Similar results were noted by **Jamra et al. (2018), Kour et al. (2019)** in Aonla, **Verma et al. (2014)** in phalsa, **Rai et al. (2009)** in Pear and **Chawla et al. (2020)**. The application of Poultry Manure (7.5 kg/tree) + RDF 50% *Azospirillum* + (10 ml/tree) + PSB (10 ml/tree) (Table 1) showed significant maximum non-reducing sugars (3.0%) followed by FYM (10 kg/tree) + RDF 50% + *Azospirillum* (10 ml/tree) PSB (10 ml/tree). While, minimum non-reducing sugar (1.9%) was observed in the control treatment T<sub>1</sub>. The same reason could be applied that the quick metabolic transformation of starch and pectin into soluble compounds and rapid translocation of sugars from leaves to the developing fruits. The high non-reducing sugar (2.52%) were also recorded with application of Poultry Manure (7.5 kg/tree) + RDF 50% + *Azospirillum* (10 ml/tree) + PSB (10 ml/tree). The research findings were supported by **Verma et al (2014)** in Phalsa, **Chawla et al. (2020), Jamra et al. (2018) and Kour et al. (2019)** in Aonla.

### **Total Sugar (%):**

In Table 1 the highest total sugars value (6.29%) were recorded in T<sub>10</sub> treatment that is Poultry Manure (7.5 kg/tree) + RDF 50% + *Azospirillum* (10 ml/tree) + PSB (10 ml/tree) followed by T<sub>9</sub> treatment FYM (10 kg/tree) + RDF 50% + *Azospirillum* (10 ml/tree) and T<sub>8</sub> were proved equally good with T<sub>10</sub>. However minimum total sugar percent (4.00%) was recorded in control treatment T<sub>1</sub>.

The findings are corroborating with those reported by Singh *et al.* (2008), Sharma *et al.* (2022), Kour *et al.* (2019) in aonla fruits, Verma *et al.* (2014) and Yadav *et al.* (2008) in Phalsa.

## CONCLUSION:

Based on the results of this present investigation, it can be concluded that treatment T<sub>10</sub> (Poultry Manure (7.5kg/ tree) + RDF 50% + *Azospirillum* (10ml/tree) + PSB (10ml/tree) produced the maximum TSS (11.83<sup>0</sup> Brix), Ascorbic acid content ( 599.98 mg/ 100gfruitpulp), reducing sugar (3.28%), non-reducing sugar (3.00%), total sugar (6.29%) and minimum acidity (1.32%) of aonla which was at par with T<sub>9</sub>-FYM (10kg/tree) + RDF 50% + *Azospirillum* (10ml./tree) + PSB (10ml./tree) can be recommended to aonla growers in eastern Uttar Pradesh for obtaining higher yield with better quality fruits.

## FUTURE SCOPE

Integrated nutrient management, developed on the principles of eco-friendly and efficient balanced fertilizer and based in optimization of nutrient supplies from all the available sources, inorganic and organic, which lowers production costs and increases productivity, were the greatest ways to cultivate Aonla.

## REFERENCE

- Aal, J.M.; Patel, K. M.; Patel, S.J.; and Sindha, D. J. (2020). Effect of integrated nutrient management on fruit yield of aonla (*Emblica officinalis* Gaertn) cv. Gujarat Aonla-1, *International Journal of Current Microbiology and Applied Sciences*, 9(10): 417-423.
- Anonymous(2021-22), Indian Horticulture Database, National Horticulture Board. Department of Agriculture and Co-operation & Farmers Welfare. Government of India, Gurugram.
- A.O.A.C. 1990. Official methods of analysis of 16th edition. Association of official Agricultural Chemists, Washington D.C.
- Athani, S.I.; Ustad, A.I.; Prabhuraj, H.S.; Swamy, G.S.K.; Patil, P.B. and Kotikal, Y.K. (2009). Influence of vermicompost on growth, fruit yield and quality of guava cv. Sardar. *Acta Horticulture*, 735(2): 381-85.

- Bhabiskar, M.N.; Bharad, S.G.; Dod, V.N. and Barne, V.G. (2011).** Effect of integrated nutrient management on yield and quality of sapota. *Plant Archives*, 11(2):661-663.
- Bohane L, Tiwari R. (2014).** Effect of integrated nutrient management on physicochemical parameters of ber under Malwa plateau conditions. *Annals of Plant and Soil Research*, 16(4):346-348.
- Chawla, R., and Sadawarti, R. K. (2020).** Effect of bio-fertilizers and organic manures on growth, yield and fruit quality of fruit crops. *Plant Arch*, 20(1): 3767-3768.
- Jaiswal, A., Prasad, V. M., Bhadur, V., and Kumar, N. (2023).** Effect of organic manure and inorganic fertilizer on growth, yield and quality of guava (*Psidium guajava* L.), *The Pharma Innovation Journal*, 12(2): 2133-2136.
- Jamra, R., Kanwar, J., Dubey, R. and Tiwari, R. (2018).** Effect of integrated nutrient management practices on physical, biochemical properties and yield of aonla (*Emblica officinalis* Gaertn). *International Journal of Chemical Studies*, 6(2): 1256-1258.
- Kour, D., Wali, V. K., Bakshi, P., Bhat, D. J., Sharma, B. C., Sharma, V., and Sinha, B. K. (2019).** Effect of Organic and Inorganic fertilizers along with Azotobacter on Growth, Yield and Quality of Aonla (*Emblica officinalis* Gaertn.) Cv. Na-7. *International Journal of Current Microbiology and Applied Sciences*, 8(9): 1142-1151.
- Kumar, A., Prasad, V.M., Singh, D., Bahadur, V., David, A.A. and Beer, K. (2018).** Effect of bio-fertilizers, vermicompost and Trichoderma on yield and economics of strawberry (*Fragaria x annanasa* Duch.) cv. Sweet Charlie. *International Journal of Current Microbiology and Applied Sciences*, 7(6): 1534-1538.
- Panse, V.C. and Sukhatme, P.V. (1985).** Statistical Methods for Agricultural Workers, ICAR, New Delhi.
- Rai, J.; Singh, S.P. and Singh, A.K. (2017).** Economics of aonla production in district Pratapgarh (U.P.). *International Research Journal of Agricultural Economics and Statistics*, 8(2): 426-430.

- Rai, P.N., Singh, P.N. and Binayak, C. (2009).** Effect of different sources of organic manures on yield and quality of low-chill pear (*Pyrus pyrifolia*) cv. Gola. *Progressive Horticulture*, 41(1): 103-106.
- Sharma, A. and Nagaich, K. (2022).** Biofertilizer–An Effective Component of Integrated Nutrient Management in Aonla. *Annals of Biology*, 38(1): 83-87.
- Sharma, A. and Nagaich, K. (2022).** Biofertilizer–An Effective Component of Integrated Nutrient Management in Aonla. *Annals of Biology*, 38(1): 77-82.
- Sharma, B.D., Meena Anita, Jatav, M.K. and Balai, R.C. (2018).** Integrated nutrient management for horticultural crops in arid region. ICAR -CIAH/ Tech./ Pub. No.77, pp 42.
- Sharma, S., Rana, V. S., Rana, N., Sharma, U., Gudeta, K., Alharbi, K., and Bhat, S. A. (2022).** Effect of Organic Manures on Growth, Yield, Leaf Nutrient Uptake and Soil Properties of Kiwifruit (*Actinidia deliciosa* Chev.) cv. Allison. *Plant Archives*, 11(23): 3354.
- Singh, A.; Yadav, A.L., Yadav, D.K. and Mishra, S.K. (2008).** Effect of integrated nutrient management on yield and quality of aonla (*Emblia officinalis*Gaertn.) cv. NA-10. *Plant Archives*, 8(1): 473- 474.
- Singh, A. K., Singh, S. A. N. J. A. Y., Saroj, P. L., Mishra, D. S., Singh, P. P., and Singh, R. K. (2019).** Aonla (*Emblia officinalis*) in India: a review of its improvement, production and diversified uses. *Indian Journal of Agriculture Sciences*, 89(11): 1773-1781.
- Tripathi, V. K., and Nand, V. (2022).** Effect of foliar application of Boron, Zinc and NAA on fruit retention, yield and quality attributes of Aonla. *Progressive Horticulture*, 54(1): 76-81.
- Verma R.S.; Singh HK, Verma SS. (2014).** Effect of integrated nutrient management on plant growth, fruit yield and quality of phalsa (*Grewia subinaequalis* D.C.). *Asian Journal of Horticulture*, 9(1):48-52.
- Yadav, D.K.; Pathak, Sanjay and Yadav, A.L. (2008).** Effect of integrated nutrient management on physico-chemical analysis of Phalsa (*Grewia subinaequalis* D.C.). *Plant Archives*, 8(1): 461-463.

**Yadav, Rajesh; Singh, H.K. and Yadav; A.L. (2007).** Effect of integrated nutrient management on productivity and quality of aonla (*Emblica officinalis* Gaertn). *Plant Archives* 7(2): 881- 883.