

## Effect of Organic and Inorganic fertilizer on Growth and Yield attributes of Okra (*Abelmoschus esculentus* L.)

### ABSTRACT

Research was conducted by using effective management application of Flyash, Azospirillum and inorganic fertilizer with objective to enhance the growth and yield of okra crop. The field experiment was carried out at the research farm of soil science and agricultural chemistry, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj during Zaid season 2022. The experiment was laid out in Randomized Block Design (RBD) with 9 treatments replicated three times. Data were collected on plant height, number of leaves, number of branches, fruits per plants and yield. Results showed significant differences on all parameters such that the best plant height with the most leaves, high number of branches, greater number of fruits per plant and highest yield was registered in treatment with the combination of Flyash 25 t ha<sup>-1</sup>, Azospirillum 2 kg ha<sup>-1</sup> with RDF 100: 60: 50 kg ha<sup>-1</sup> indicating that the synergy by Azospirillum, Flyash and fertilizers improved growth and yield. The results are based on one season experiment, therefore further investigation is recommended to substantiate the findings.

**Keywords:** Azospirillum, Flyash, growth, inorganic fertilizers and yield parameters

### 1. INTRODUCTION

Okra or ladies finger (*Abelmoschus esculentus* L.) requires well draining soil that is rich in organic matter and nutrients such as nitrogen, phosphorus, and potassium. Soil affects the water holding capacity and pH level, which can impact the plant's growth and yield. Additionally, soil health is crucial for the prevention of diseases and pests that can infect the okra plant. Therefore, it is important to maintain soil fertility and quality through proper management practices such as crop rotation, composting, and appropriate irrigation. Soil is a critical component in the successful cultivation of okra and must be managed carefully to ensure optimal plant growth and yield (Kumar *et al.*, 2017).

Flyash has great potentiality in agriculture due to its efficiency in modification of soil health and crop performance. Flyash is the end residue from combustion of

Formatted: Font: 12 pt

pulverized bituminous or sub-bituminous coal (lignite) in the furnace of thermal power plants and consists of mineral constituents of coal which is not fully burnt. The high concentration of elements (K, Na, Zn, Ca, Mg and Fe) in flyash increases the yield of many agricultural crops. But compared to other sectors, the use of flyash in agriculture is limited. Flyash having both the soil amending and nutrient-enriching [properties](#) is helpful in improving crop growth and yield in low fertility acid lateritic soils. Many researchers have demonstrated that flyash increased the crop yield and improved the physical and chemical characteristics of the soil ([Basu et al., 2019](#)).

Azospirillum works by colonizing the roots of the okra plant and fixing atmospheric nitrogen, which is then used by the plant for its growth and development. [The results](#) in improved plant vigor, increased shoot and root growth, and increased flower and fruit production ([Shree et al., 2014](#)).

## **2. MATERIALS AND METHOD**

The field experiment was carried out at the research farm of soil science and agricultural chemistry, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj during Zaid season 2022. The maximum temperature of the location ranges between 46<sup>o</sup>C-48<sup>o</sup>C and seldom falls below 4<sup>o</sup>C-5<sup>o</sup>C. The relative humidity ranges between 20-94%. [The](#) average rainfall of this area is around 1100mm annually. The experiment was laid out in Randomized Block Design (RBD) with 9 treatments [replicated three times](#). Detailed combination of treatments [is](#) given in Table 1. [Data](#) were [collected on](#) plant height, number of leaves, number of branches, fruits per plants and yield.

[How was the data analyzed?](#)

[What was the nutrient content of Flyash after analysis?](#)

[What was the nutrient content of Azospirillum after analysis?](#)

**Comment [H1]:** What was the status of the soil before the experiment? This information is paramount.

**Table 1.:** Treatment Combination

Treatment	Treatment Combination
T <sub>1</sub>	Absolute control
T <sub>2</sub>	[NPK @ 0% + FLYASH @ 50% + AZOSPIRILLUM @ 25%]
T <sub>3</sub>	[NPK @ 0% + FLYASH @ 100% + AZOSPIRILLUM @ 50%]
T <sub>4</sub>	[NPK @ 50% + FLYASH @ 0% + AZOSPIRILLUM @ 0%]
T <sub>5</sub>	[NPK @ 50% + FLYASH @ 50% + AZOSPIRILLUM @ 25%]
T <sub>6</sub>	[NPK @ 50%+FLYASH @ 100% + AZOSPIRILLUM @ 50%]
T <sub>7</sub>	[NPK @ 100% + FLYASH @ 0%+ AZOSPIRILLUM @ 0%]
T <sub>8</sub>	[NPK @ 100% + FLYASH @ 50% + AZOSPIRILLUM @ 25%]
T <sub>9</sub>	[NPK @ 100% + FLYASH @ 100% + AZOSPIRILLUM @ 50%]

### 3. RESULTS AND DISCUSSION

#### 3.1. Plant Height at 25, 50 and 75 DAS

The effect of nutrients [from](#) NPK, Flyash and Azospirillum on plant height was found significant at 25, 50 and 75 DAS. The maximum plant height 25.91, 92.12 and 114.89 cm [were](#) recorded in T<sub>9</sub> (NPK @ 100% + FA @ 100% + Azospirillum @ 50%) and minimum plant height 20.91, 82 and 106.25 cm [were](#) recorded in T<sub>1</sub> (Absolute Control) at 25, 50 and 75 DAS respectively. The application of NPK, Flyash and Azospirillum in combination can result in a synergistic effect on plant growth and development. NPK provides essential nutrients to plants. This leads to improved plant growth and development, resulting in an increase in plant height at different growth stages [as reported by](#) Kumar *et al.*, (2017). Increase in plant height due to

increase in NPK, Flyash and Azospirillum [that provide](#) adequate nutrients which in turn [enhanced the](#) vigorous vegetative growth of plants and subsequently increase the plant height through cell elongation, cell division and photosynthesis of plant cell. Similar findings were reported by Singh *et al.*, [\(2018\)](#).

### **3.2 Number of Leaves per Plant at 25, 50 and 75 DAS**

[Results in Table 2 showed significant difference on number of leaves due to the](#) effect of nutrients [from](#) NPK, Flyash and Azospirillum at 25, 50 and 75 DAS. [The](#) maximum number of leaves 14.02, 27.52 and 39.31 [were](#) recorded in T<sub>9</sub> (NPK @100% + FA @100% + Azospirillum @50%) and minimum number of leaves 8.12, 21.17 and 28.76 [were](#) recorded in T<sub>1</sub> (absolute control) at 25, 50 and 75 DAS respectively. The combination of organic and inorganic fertilizers [can have a](#) synergistic effect on plant growth and development. Flyash provides nutrients while Azospirillum enhances [their](#) uptake and produces plant growth regulators. the combination of these two inputs can promote vegetative growth and increase the number of leaves in okra plants [as reported by](#) Verma *et al.*, [\(2013\)](#).

### **3.3 Number of branches per plant at 25, 50 and 75 DAS**

[Results on number of branches per plant revealed significant difference such that the](#) maximum number of branches 16.02, 20.75 and 27.27 [at 25, 50 and 75 DAS were](#) registered in T<sub>9</sub> and minimum number of branches 9.11, 15.02 and 19.13 [were](#) recorded in T<sub>1</sub> [respectively](#). This could be due to the effect brought about by Flyash, Azospirillum, and inorganic fertilizers. [The nutrients](#) positively influences [the](#) number of branches in okra plants by improving nutrient availability, enhancing microbial activity, and promoting overall plant vigor (Singh *et al.*, [2018](#)).

### **3.4 Number of fruits per plant**

[Results in Table 2 showed significant difference on number of fruits per plant.](#) [Treatment \(T<sub>9</sub>\) registered highest number of fruits per plant \(19.87\) while the lowest](#) number of fruits (14.21) was recorded in the control treatment. [This is attributed to](#)



<b>SE±</b>	?	?	?	?	?	?	?	?	?	?	?
<b>CV%</b>	?	?	?	?	?	?	?	?	?	?	?
<b>P=0.05</b>	?	?	?	?	?	?	?	?	?	?	?

**Comment [H2]:** Fill in the statistics or else it will be difficult to defend the significance reported in the write-up.

#### 4. CONCLUSION

It is evident to conclude that the application of Flyash and Azospirillum with inorganic fertilizers in treatment T<sub>9</sub> (NPK @100% + FA @100% + Azospirillum @50%) was found best in increasing growth and yield of okra. Since the results are based on one season experiment, further trail is needed to substantiate the results.

#### REFERENCES

Basu M, Pande M, Bhadoria PBS, Mahapatra SC. Potential flyash utilization in agriculture: A global review. *Progress in Natural Science*. 2019;19(10):1173-1186.

Das B, Sahoo S, Mishra P. Effect of nitrogen on growth, yield and quality of okra (*Abelmoschus esculentus* L.). *International Journal of Agriculture, Environment and Biotechnology*. 2014;7(2):285-291.

Ginindza MM, Aderibigbe TO, Jaiyeoba IA. Effect of potassium fertilizer application on okra (*Abelmoschus esculentus* L. moench) growth, yield and quality in Switzerland. *International Journal of Agriculture and Biology*. 2015;17(5):973-978.

Kumar V, Yadav A, Singh A, Singh D, Singh R. Effect of integrated nutrient management on growth, yield and quality of okra (*Abelmoschus esculentus* L.). *Journal of Pharmacognosy and Phytochemistry*. 2017; 6(5):3525-3527.

Meena VD, Dotaniya ML, Coumar V, Rajendiran S, Sharma P. Effect of phosphorus on crop growth, yield, and nutrient content in okra (*Abelmoschus esculentus*) under field conditions. *Journal of Plant Nutrition*. 2017;40(11):1573-1580.

Shree M, Kumar V, Tripathi RD. Azospirillum sp. inoculation improves growth and yield of okra (*Abelmoschus esculentus*) in a farmer's field. African Journal of Microbiology Research. 2014;8(45): 4001-4008.

Singh R, Kumar R, Sharma S. Impact of organic and inorganic sources of nutrient on yield and quality of okra (*Abelmoschus esculentus* L. moench). International Journal of Chemical Studies. 2018;6(2):251-256.

Verma A, Kaur S, Kumar A, Singh G. Effect of organic manures and inorganic fertilizers on growth, yield, and quality of okra (*Abelmoschus esculentus* L. moench). Journal of Plant Nutrition. 2013;36(7):1073-1089.

Yadav RL, Yadav KS, Kumar S, Yadav S, Yadav P. Influence of integrated nutrient management on yield and quality of okra (*Abelmoschus esculentus* L.) in inceptisols. Indian Journal of Agricultural Sciences. 2020;90(4):51-54.