

## Effect of foliar spray of micronutrients and plant growth regulators on flowering, fruit pomology and oil yield

---

### ABSTRACT

Olive (*Olea europaea*) is a small evergreen tree native to the Mediterranean region, known for its distinctive and valuable fruit, the olive. The olive tree has been cultivated for thousands of years and holds great cultural and economic significance in various parts of the world. The use of micronutrients sprays in olive cultivation is a common practice to address nutrient deficiencies, enhance plant health and increase flowering and improve fruit quality. These sprays contain essential minerals like iron, zinc, manganese, and boron, which are readily absorbed by the olive trees through their leaves. Micronutrients sprays help to correct imbalances, improve nutrient uptake, and boost overall growth and fruit development. They are particularly effective in regions with poor soil conditions or where specific micronutrient deficiencies are prevalent, ensuring healthy olive trees and optimal yield of high-quality fruits. Therefore, the present investigation was carried out at the Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture Technology and Sciences, Prayagraj, Uttar Pradesh during the olive harvesting 2021-2022 with a view to determine the effect of different micronutrient applications olive for its growth quality. On the basis of the present investigation, it is concluded that among the various treatments applied to enhance the Flowering, Fruit set, fruit Yield, Fruit quality and Oil quality of Olive cultivars, treatment T<sub>10</sub> (Boron @ 150 ppm + Zinc @ 150 ppm + NAA@ 150 ppm + GA<sub>3</sub> @ 150 ppm) was found to be superior among others, followed by treatment T<sub>5</sub> (Boron @ 100ppm + Zinc @ 100 ppm + NAA @ 100ppm + GA<sub>3</sub> @ 100ppm) and treatment T<sub>0</sub> (control) was found to be inferior. It is also concluded that among all cultivars, Arbequina cultivar was found best. Thus, treatment T<sub>10</sub> (Boron @ 150 ppm + Zinc @ 150 ppm + NAA@ 150 ppm + GA<sub>3</sub> @ 150 ppm) with Arbequina cultivar is best recommended for overall flowering and fruiting characters of olive like, days to flowering, number of flowers per panicle, fruit drop percentage, fruit diameter, fruit length, fruit weight, fruit yield per tree, fruit yield per hectare, oil content, total suspended solids (TSS) of fruit.

**Keywords:** *Olive, micronutrients, Plant growth regulators, Boron, Zinc, NAA, GA<sub>3</sub>.*

---

### INTRODUCTION

Olive is native to the Mediterranean region; botanical name of Olive is *Olea europaea* and it belongs to the Oleaceae family. It is an evergreen tree with a height ranging from 12-15 meters. Olive is characterized by its silvery-green leaves, tiny whitish flowers and small oval shaped fruits. Olive trees are evergreen in nature and they can tolerate adverse edaphic conditions including high temperature and draught; these abilities of olive tree make it one of the hardy crops and bear fruits even during the tough environmental challenges. Olive trees are alternate bearers; they produce important yield for one year and lower yield in the succeeding year. Olive is of major agricultural importance in the Mediterranean region as the source of olive oil (Bertrand *et al.* 2002). Olive is one among the oldest cultivated crops known, it holds a significant historical, cultural, medicinal and economic importance in numerous countries. Olive is most widely cultivated in Mediterranean countries namely, Spain, Greece, Italy, Turkey, Portugal, and Tunisia. In India olive cultivation is pertained to Parts of Rajasthan, Himachal Pradesh, Uttar Pradesh, and Delhi. In the world, the olive area and production is 10,839,026 ha and 18,083,800 tons respectively (F.A.O., 2021). Olives are primarily used for oil extraction apart from this oil is also consumed as snacks, pickles. Although, olive tree has been designated as a drought tolerant yet, it requires sufficient soil moisture during certain stages of growth. Unfortunately, majority of olive plantations were undertaken on hill slopes, in the drought prone areas of mid hills and valley areas of the state.

Furthermore, these areas are completely devoid of irrigation facility. An erratic trend of monsoon and winter rains has become more conspicuous in the last decade which further aggravated the problem of poor growth and bearing of olive trees. Acute water stress during autumn coupled with scanty or insufficient and irregular rainfall distribution. Boron induces pollen tube growth resulted from its role on tryptophan synthesis as an auxin precursor biosynthesis. The main function of boron is related to cell wall strength and development, cell division, sugar transport and hormones development, RNA metabolism, respiration, indole acetic acid (IAA) metabolism and as part of the cell membranes. Lewis (1980) speculated that B may be required in stigma and styles to physiologically inactivate callus present in pollen tube walls that would otherwise elicit phytoalexin production to inhibit pollen tube growth. The boron requirement is much higher for reproductive growth period than for vegetative growth and increases flower production and retention, pollen tube elongation and germination, and seed and fruit development. Several investigators studied the effect of zinc and/or boron on fruit set, productivity and fruit quality in many plant species. Talaie *et al.* (2001) showed that foliar spray of B and Zn decreased fruit drop and increased fruit quality in the 'Zard' olive. Hassan *et al.* (2010) found that boric acid treatments increased pollen germination than control and increased percentage of retained fruits in 'Picual' olive. Abd El-Migeed *et al.* (2015) on 'Picual' olive reported that boric acid spray at 300 mg/l increased fruit length. Osman (1999) on olive found that boron treatments either as foliar or soil applications increased percentage of retained fruits. Khayyat *et al.* (2015) reported that boric acid at 1500 mg/l on 'Shahany' date palm increased pulp weight, pulp/seed ratio; fruit length and diameter. Plant growth regulators or Phytohormones play a very important role in modification or regulation of the physiological processes in plants. Among Phytohormones NAA and GA3 play a key role in Flowering, and Fruit set of Olive. Plant growth regulators are applied to plants through various methods but, Plant growth regulators when applied through Foliar spray gives desired results within a short span of time since, they are easily absorbed by plant tissues when applied directly on different plant parts. NAA plays a major role in strengthening the root system by new root initiation in the plants. It enables proper functioning of roots which transport water, nutrients and other needed elements for vegetative and floral growth of the tree. NAA application in olive is mostly done for the purpose of fruit thinning in order to avoid overcrowding of fruits on the branch. Application of NAA reduces excess number of fruits on the branch thus promoting the growth of evenly spaced fruits on branch and directly plays a major role in increasing the overall fruit quality of Olive Arnon *et al.* (2009). Gibberellic acid plays a major role when applied on Olive trees through Foliar spray. It promotes cell enlargement and mesocarp development of Olives, which is a desirable character in increasing the fruit size and fruit quality of Olive. GA3 also helps in reduction of fruit drop and increases the fruit retention in Olive trees. Since GA3 directly regulates elongation, enlargement and growth of cells thus increasing the fruit length and diameter of the olive fruits. Ramezani *et al.* (2010).

## **MATERIAL AND METHODS**

The present investigation was done to understand the plant growth, fruit yield and quality of olive using different sprays of micronutrients. The details of the materials used, and the methods adopted in the investigation, which was carried out at Horticultural Research Farm, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences (SHUATS), Prayagraj (India) during the *Winter* season of 2022-23. The design used in study was randomized block design

(RBD) each treatment was replicated thrice. The data were statistically analysed (by the method suggested by Fisher and Yates (1963)). The treatments comprised of T<sub>0</sub> (Control), T<sub>1</sub> (Boron @ 100 ppm), T<sub>2</sub> (Zinc @ 100 ppm), T<sub>3</sub> (NAA @100 ppm), T<sub>4</sub> (GA<sub>3</sub> @ 100 ppm), T<sub>5</sub> (Boron@ 100 ppm+ Zinc @ 100 ppm + NAA @100 ppm +GA<sub>3</sub> @100 ppm), T<sub>6</sub> (Boron @ 150 ppm), T<sub>7</sub> (Zinc @ 150 ppm), T<sub>8</sub> (NAA @150 ppm), T<sub>9</sub> (GA<sub>3</sub> @ 150 ppm) and T<sub>10</sub> (Boron @ 150 ppm+ Zinc @ 150 ppm + NAA @150 ppm + GA<sub>3</sub> @ 150 ppm). Spraying was done prior to fruit harvest at time of formation of fruit. Observations were recorded at different stages of growth periods for characters like days to flowering, number of flowers per panicles, days to flowering to fruiting, fruit length, fruit diameter, fruit weight. Chemical parameters like oil content, TSS etc were also calculated. Fruit weight was calculated after harvest using electronic balance. TSS was measured using refractometer.

## RESULTS AND DISCUSSION

### A) Flowering attributes: Days to flowering, number of flowers per panicles

Days to flowering showed that there were significant differences among the treatments. The minimum days to flowering (13.33) was recorded under the treatment T<sub>10</sub> which is at par with T<sub>5</sub> (15.33) followed by the treatment T<sub>9</sub> (18.33) T<sub>8</sub> (19.33) respectively. The maximum days to flowering (28.67) was recorded under the treatment T<sub>0</sub> (Control) followed by the treatments T<sub>1</sub> (26.67) and T<sub>2</sub> (26.00) respectively. The maximum number of flowers per panicle (26.00) was recorded under the treatment T<sub>10</sub> which is at par with treatment T<sub>5</sub> (24.33) which is followed by the treatment T<sub>9</sub> (22.00) flowers respectively. The minimum number of flowers per panicle (9.33) was recorded in the treatment T<sub>0</sub> (Control) and followed by the treatments T<sub>1</sub> (13.33) followed by T<sub>2</sub> (15.67) flowers respectively. The maximum flowers per panicle was observed in T<sub>10</sub>. Significant improvement in flowering and fruiting components such as the number of flowers per shoot, fruit set (%), fruit drop (%) and fruit retention (%) as influenced by increasing levels of NAA and boron treatments was observed. Combined form of micronutrients and GA<sub>3</sub> plays a vital role in prompting proper growth, increasing flower production and retention, pollen tube elongation and germination, and seed and fruit development by Regulating plant's hormone levels. Not only it regulates the flowering but also plays an important role in controlling the flower drops. The similar experiment was also conducted by Aftab, 1994 and found the use of zinc, boron and growth regulators on flowering, fruiting, and maturity of litchi. The obtained results are close with that obtained by Shaheen (1995), Osman (1999), and Hassan (2000) on Olive. The minimum days to flowering in T<sub>10</sub> might be since Boron plays an essential role in plant's life cycle and very essential for normal growth of plants. [Fageria *et al.* (2007), McLaughlin *et al.* (1999)]. While Significant improvement in flowering and fruiting components such as the number of flowers per shoot, fruit set (%), fruit drop (%) and fruit retention (%) as influenced by increasing levels of NAA and boron treatments was observed during the present investigation. Similar results were also seen in the experiment conducted by Badal and Tripathi (2021) on olive (*Psidium guajava* L. cv L-49).

### B) Fruiting attributes

#### 1. Percent of fruit drop, Fruit weight, fruit length, fruit diameter and fruit volume.

Fruit drop Percentage showed that there were significant differences among the treatments. Minimum fruit drop percentage (19.67) was recorded under the treatment T<sub>10</sub> followed by T<sub>5</sub> (21.33), T<sub>9</sub> (25.67) which was at par with T<sub>8</sub> (27.00). Maximum fruit drop percentage (37.00) was recorded under the treatment T<sub>0</sub> (Control) followed by T<sub>3</sub> (34.33) is followed by T<sub>1</sub> (34.67). The least percent of fruit drop

were observed in the treatment T<sub>10</sub>. Environmental factors are key responsible for the fruit drop in most of the species. Application of GA<sub>3</sub> with the combination of micronutrients have shown greater responses against the fruit drop. Similar results were also obtained in (Nagy & Kovacs, 2005).

Fruit diameter showed that there were significant differences among the treatments. Maximum fruit diameter (16.67) was recorded under the treatment T<sub>10</sub> is at par with T<sub>5</sub> (15.67) followed by T<sub>9</sub> (13.67) respectively. Minimum fruit diameter was recorded under the treatment T<sub>0</sub>(control) (7.67) which is at par with T<sub>1</sub> (8.00) and T<sub>2</sub> (8.33) respectively. Fruit length showed that there were significant differences among the treatments. Maximum fruit length (23.00) was recorded under the treatment T<sub>10</sub> followed by T<sub>5</sub> (21.77), T<sub>9</sub> (20.33) and T<sub>8</sub> (19.17) which was at par with T<sub>7</sub> (18.33). Minimum fruit length (12.83) was recorded under the treatment T<sub>0</sub> (Control) which is at par with T<sub>1</sub> (13.37). Maximum fruit weight (4.23) was recorded under the treatment T<sub>10</sub> followed by T<sub>5</sub> (4.00), T<sub>9</sub> (3.57) which was at par in with T<sub>8</sub> (3.40). Minimum fruit weight (1.63) was recorded under the treatment T<sub>0</sub> (Control) followed by the treatments T<sub>1</sub> (1.90), T<sub>2</sub> (2.23) respectively. Present investigation showed that the maximum fruit weight on an average (4.23kg) was obtained in T<sub>10</sub> as compared to control (1.63kg). It might be due to use of growth regulators to improve the use of stored carbohydrates, nitrogen and other variables, Singh (2013) in *Citrus limon*, Kaur Sukhjiti (2017) in Florida guard peach and Siddiqua *et al.* (2018) in dragon fruit, also recorded these results.

## **2. Fruit yield per tree and fruit yield per hectare**

Maximum fruit yield per tree (3.10) was recorded under the treatment T<sub>10</sub> followed by T<sub>5</sub> (2.57), T<sub>9</sub> (1.73) and T<sub>8</sub> (1.50) respectively. Minimum fruit yield per tree (0.34) was recorded under the treatment T<sub>0</sub> (control) followed by T<sub>2</sub> (0.59). The maximum yield obtained was (12.40 q/ha) T<sub>10</sub>. Fruit yield per hectare showed that there were significant differences among the treatments. Maximum fruit yield per hectare (12.40) was recorded under the treatment T<sub>10</sub> followed by T<sub>5</sub> (10.27). Minimum fruit yield per hectare (1.37) was recorded under the treatment T<sub>0</sub> (control) followed by T<sub>2</sub> (Zinc @ 100 ppm) (2.35), T<sub>3</sub> (2.45) respectively. The beneficial effect of boron and zinc in increasing fruit yield might be due to the higher availability of photosynthesis, and/or their role in increasing the percent of perfect flowers and these chemicals are also associated with hormone metabolism which promotes synthesis of auxin, essential for fruit set and growth. The results are in accordance with Kazemi (2014). Similar results of increased yield due to the application of ZnSO<sub>4</sub> were reported by Pathak *et al.* (2004) in olive.

## **C) Quality attributes**

### **1. TSS, oil content**

Oil content showed that there were significant differences among the treatments.

Maximum Oil content (20.83) was recorded under the treatment T<sub>10</sub> followed by T<sub>5</sub> (19.70), T<sub>9</sub> (18.67) respectively. Minimum Oil content (10.20) was recorded under the treatment T<sub>0</sub> (Control) followed by T<sub>1</sub> (11.53). The results observed are also in agreement with that obtained by Shaheen (1995), Weisman *et al.* (2002) and Hassan (2000) who found great increase in fruit oil content of olive trees due to boron treatments. Moreover, Kamal (2002) found that boron and Zinc with a combination of GA<sub>3</sub> applications increased the oil percentage in olive fruits.

TSS of fruit showed that there were significant differences among the treatments. Maximum TSS of fruit (13.80) was recorded under the treatment T<sub>10</sub> followed by T<sub>5</sub> (13.37), T<sub>9</sub> (GA3 @ 150 ppm) (12.90) respectively. Minimum TSS of fruit (10.10) was recorded under the treatment T<sub>0</sub> (control) followed by T<sub>1</sub> (10.60), T<sub>2</sub> (10.90) respectively. The combination of micronutrients and plant growth regulators have shown positive affect on the TSS of olive fruit. The maximum TSS have shown was 13.80 in the treatment T<sub>10</sub>. The studies pertaining to bio-chemical status of fruits reflected that TSS (%) was affected significantly by the micronutrients or the plant growth regulators in the finding of Rajkumar *et al.* (2014).

### **Conclusion**

On the basis of the present investigation, it is concluded that among the various treatments applied to enhance the Flowering, Fruit set, fruit Yield, Fruit quality and Oil quality of Olive cultivars, treatment T<sub>10</sub> (Boron @ 150 ppm + Zinc @ 150 ppm + NAA@ 150 ppm + GA3 @ 150 ppm) was found to be superior among others, followed by treatment T<sub>5</sub> (Boron @ 100ppm + Zinc @ 100 ppm + NAA @ 100ppm + GA3 @ 100ppm) and treatment T<sub>0</sub> (control) was found to be inferior. It is also concluded that among all cultivars, Arbequina cultivar was found best. Thus, treatment T<sub>10</sub> (Boron @ 150 ppm + Zinc @ ppm + NAA@ 150 ppm + GA3 @ 150 ppm) with Arbequina cultivar is best recommended for overall flowering and fruiting characters of Olive like, Days to flowering, Number of flowers per panicle, Fruit drop percentage, Fruit diameter, Fruit length, Fruit weight, Fruit yield per tree, Fruit yield per hectare, Oil content, TSS of fruit.

**Table 1 Performance of different micronutrients application on various parameters of olive.**

<b>Treatment symbol</b>	<b>Treatment Details</b>	<b>Days to flowering</b>	<b>Flowers per panicle</b>	<b>Fruit drop (%)</b>	<b>Fruit Diameter (mm)</b>	<b>Fruit length (mm)</b>	<b>Fruit weight (g)</b>	<b>Fruit yield (Kg/tree)</b>	<b>Fruit yield (q/h)</b>	<b>Oil content (%)</b>	<b>TSS (°Brix)</b>
<b>T<sub>0</sub></b>	Control	28.67	9.33	37.00	7.67	12.83	1.63	0.34	1.37	10.20	10.10
<b>T<sub>1</sub></b>	Boron @ 100 ppm	26.67	13.33	34.67	8.00	13.37	1.90	0.49	1.95	11.53	10.60
<b>T<sub>2</sub></b>	Zinc @ 100 ppm	26.00	15.67	31.33	8.33	14.00	2.23	0.59	2.35	12.73	10.90
<b>T<sub>3</sub></b>	NAA @100 ppm	25.00	16.33	34.33	9.67	15.07	2.60	0.61	2.45	13.17	11.17
<b>T<sub>4</sub></b>	GA <sub>3</sub> @ 100 ppm	23.00	17.00	31.00	10.00	16.33	2.80	0.87	3.49	14.60	11.50
<b>T<sub>5</sub></b>	Boron@ 100 ppm+ Zinc @ 100 ppm + NAA @100 ppm +GA <sub>3</sub> @100 ppm	15.33	24.33	21.33	15.67	21.77	4.00	2.57	10.27	19.70	13.37
<b>T<sub>6</sub></b>	Boron @ 150 ppm	21.33	20.00	30.00	11.00	17.43	3.00	1.01	4.04	15.20	11.80
<b>T<sub>7</sub></b>	Zinc @ 150 ppm	20.67	20.67	28.33	12.33	18.33	3.10	1.20	4.80	16.77	12.13
<b>T<sub>8</sub></b>	NAA @150 ppm	19.33	21.33	27.00	13.67	19.17	3.40	1.50	6.00	17.13	12.60
<b>T<sub>9</sub></b>	GA <sub>3</sub> @ 150 ppm	18.33	22.00	25.67	14.00	20.03	3.57	1.73	6.93	18.67	12.90
<b>T<sub>10</sub></b>	Boron @ 150 ppm+ Zinc @ 150 ppm + NAA @150 ppm + GA <sub>3</sub> @ 150 ppm	13.33	26.00	19.67	16.67	23.00	4.23	3.10	12.40	20.83	13.80
<b>F-Test</b>		<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>
<b>S.E.(m) (±)</b>		<b>1.28</b>	<b>1.17</b>	<b>0.97</b>	<b>0.66</b>	<b>0.40</b>	<b>0.08</b>	<b>0.07</b>	<b>0.31</b>	<b>0.13</b>	<b>0.09</b>
<b>CD (5%)</b>		<b>2.67</b>	<b>2.44</b>	<b>2.03</b>	<b>1.38</b>	<b>0.84</b>	<b>0.18</b>	<b>0.16</b>	<b>0.66</b>	<b>0.28</b>	<b>0.20</b>
<b>CV</b>		<b>7.26</b>	<b>7.60</b>	<b>4.09</b>	<b>7.03</b>	<b>2.86</b>	<b>3.62</b>	<b>7.69</b>	<b>7.69</b>	<b>1.09</b>	<b>1.01</b>

## References

- A.O.A.C. (1975).** Association of official Agricultural Chemists. Official Methods of Analysis, 12th ed., P.O. Box 450, Benjamin Franklin Station, Washington DC., pp: 832. 6. Kates, M., 1972. Techniques of Lipidology, Isolation, Analysis and Identification of Lipids. *North Holland Publishing Company*, Amsterdam, London.
- A.O.A.C. (2000).** Official Methods of Analysis: Association of Official Analytical Chemists. 15th (Ed). Washington, DC.
- A.O.C.S. (1964).** American Oil Chemists Society Official and Tentative Methods. 2nd Ed. The Society, Chicago, Illinois.
- A.R. Bonyanpour, G.R. Moafpourian and B. Jamali. (2017)** Responses of ‘Belidi’ Olives to foliar and soil applied iron, manganese and zinc fertilizers, *Journal of Applied Horticulture* 19(2): 125-129.
- Abo- El-Ez, A. E. T.; Shereen A. Shaheen and Hadeer M. Abo-Eloun (2018)** Effect of Foliar Application some Plant Growth Regulators on Growth and Productivity of Olive Trees under South Egypt Conditions. *Journal of Plant Production*, Mansoura University, 9 (10): 775–783.
- Abdrabboh G.A. (2013)** Effect of Some growth regulators on yield and fruit quality of Manzanillo Olive trees. *Nature and Science* 11(10):143-151.
- Abd El-Naby, S.K.M, M.R. El-Sonbaty, E.S. Hegazi, M.M. Samira and T.F. El-Sharony (2012)** Effect of Gibberellic Acid Spraying on Alternate Bearing of Olive Trees. *Journal of Applied Sciences Research*, 8(10): 5114-5123.
- Anshuman Singh, S. Upadhyay and R G Upadhyay (2017).** Effect of plant growth regulators (NAA, 2, 4-D and GA3) on Fruit Retention and Quality of Mango cv. Dasehri. *International Journal of Tropical Agriculture*,35(4): 967-974.
- A.R. Bonyanpour, G.R. Moafpourian and B. Jamali (2017).** Responses of ‘Belidi’ Olives to foliar and soil applied iron, manganese and zinc fertilizers. *Journal of Applied Horticulture*, 19(2): 125-129.
- Askarieh, A., Suleiman, S. and Tawakalna, M. (2021).** Sweet Cherry (*Prunus avium* L.) Fruit Drop Reduction by Plant Growth Regulators (Naphthalene Acetic Acid NAA and Gibberellic Acid GA3). *American Journal of Plant Sciences*, 12: 1338-1346.
- Badal, D.S. and Tripathi, V.K. (2021).** Influence of foliar feeding of NAA and boron on growth, flowering, fruiting, and yield of winter season guava (*Psidium guajava* L.) cv. L-49. *Biological Forum – An International Journal*, 13(3): 387-391.

- Chandramukhi Sahu and Damendra Kumar Sahu (2019)** Effect of plant growth regulators on yield and yield attributing characteristics of mango (*Mangifera indica* L) cv. Dashehari, *International Journal of Horticulture and Food Science*, 2(1): 52-54.
- Desouky IM, Laila F. Haggag, MMM, Abd El-Migeed, Kishk YFM and El- Hady (2006)**. Effect of boron and calcium nutrients sprays on fruit set, oil content and oil quality of some Olive Cultivars. *World Journal of Agricultural Sciences*, 5 (2): 180-185.
- R. Draie, W. Alhaj-Rabie, A. AL-Mahmoud (2020)**. Influence of foliar spraying with macro and microelements on the growth and productivity of olive trees in Idleb Province. *International Research Journal of Innovations in Engineering and Technology - IRJIET*, 4(4):12-24.
- E. G. Mikhail and K. G. Goargios (2014)** Effect of calcium Nitrate and Gibberellic Acid Foliar Sprays on Fruiting and Fruit Quality of “Manzanillo” and “Dolce” Olive Cvs, *Egyptian Journal of Horticulture*, 41(2):169-182.
- El- Khawaga, A.S. (2003)**. Effect of girdling and foliar application of some nutrients on growth, flowering, yield and fruit quality of Manzanillo olive trees grown in sandy soil. *Journal of Agricultural of Science*, Mansoura University 28(3): 2124-2124.
- E.S. Hegazi, R.A. El-Motaium, T.A. Yehia and M.E. Hashim (2015)**. Effect of Boron Foliar Application on Olive (*Olea europea* L.) Trees 1- Vegetative growth, flowering, fruit set, yield and fruit quality. *Journal of Horticultural Science & Ornamental Plants*, 7(1): 48-55.
- Esmail A. E. Genaidy, Nazmy Abd-Alhamid Hassan S. A. Hassan Ahmed M. Hassan and Laila F. Hagagg (2020)** Effect of foliar application of boron trioxide and zinc oxide nanoparticles on leaves chemical composition, yield and fruit quality of *Olea europaea* L. cv. Picual. *Bulletin of the National Research Centre*, 44:106.
- Ettore Barone, Michele La Mantia, Annalisa Marchese, Francesco Paolo Marra (2014)** Improvement in yield and fruit size and quality of the main Italian table olive cultivar Nocellara del Belice. *Scientia Agricola*, 71(1): 52-57.
- Fageria, N. K., Baligar, V. C. and Zobel, R. W. (2007)** 'Yield, Nutrient Uptake, and Soil Chemical Properties as Influenced by Liming and Boron Application in Common Bean in a No-Tillage System', *Communications in Soil Science and Plant Analysis*, 38:11, 1637 - 1653.
- Ferreira I. Q, Rodrigues M. A, Arrobas M (2019)**. Soil and foliar applied boron in Olive: tree crop growth and yield, and boron remobilization within plant tissues. *Spanish Journal of Agricultural Research*,17(1): e0901.

- Faisal Khalil, Khalid Mahmood Qureshi, Ammaz Khan, Fakhar-ul-Hassan and Nabila Bibi (2012)** effect of girdling and plant growth regulators on productivity in olive (*Olea europaea*) *Pakistan Journal of Agricultural Research* 25(2).
- Gurteg Singh and Harsimrat Kaur Bons (2020)** Influence of naphthalene acetic acid on fruit setting, fruit quality and yield of *Manilkara achras* L. cv. Kalipatti under subtropical conditions, *Indian Journal of Experimental Biology*, 58: 661-666.
- Gowhar A. Dar. (2017)** Impact of Boron Nutrition in Fruit crops, *International journal of Current Microbiology and Applied Sciences*, 6(12): 4145 – 4155.
- H. A. Hifny, S.M. Khalifa, A. E. Hamdy and A.N. Abd El-Wahed (2017)** Effect of GA3 and NAA on Growth, Yield and Fruit Quality of Washington Navel Orange. *Egyptian Journal of Horticulture*,44(1): 33- 43.
- Haggag, L. F., Abd El-Migeed, M. M. M., Fawzi, M. I. F., Shahin, M. F. M. and Merwad M. A. (2015)** Influence of spraying Zinc sulphate and Gibberlic acid on yield and fruit properties of “Manzanillo” Olives. *International Journal of Agricultural Technology* 11(7):1599-1611.
- Harhash M.M, W.M. E. Nasr Alla and Walid F. A. Mosa (2022).** Response of “Kalamata” olive cultivar to the foliar application of moringa extract, boron, and zinc. *Egyptian Academic Journal of Biological Sciences*, 13(2):171-178.
- Hassan HSA, Sarrwy SMA, Mostafa EAM (2010).** Effect of foliar spraying with liquid organic fertilizer, some micronutrients and gibberellins on leaf minerals content, fruit set, yield, and fruit quality of “Hollywood” plum trees. *Agriculture and Biology Journal of North America*, 1: 638-643.
- Jagtap, V.M., Patel, H.C., Nehete, D.S. and Godage, S.S. (2013).** Effect of foliar application of plant growth regulators and micronutrients on yield and quality of acid lime cv. KAGZI (*Citrus aurantifolia* Swingle), *Asian Journal of Horticulture*, 8(1): 57-59.
- Laila, F. Hagagg, M.M.M. Abd El-Migeed, Attia, M.F., M.F.M. Shahin, M.A. Merwad and E. A.-E. Genaidy (2014).** Impact of foliar application of zinc sulphate and gibberlic acid on fruit quality and quantity of “Kalamata” olives. *Middle East Journal of Agriculture Research*, 3(4): 745-750.
- Lewis DH (1980)** Are there interrelations between metabolic role of boron, synthesis of phenolic phytoalexin and the germination of pollen? *New Phytol.*, 84: 261-270.
- M. Abou El-Wafa, Ayman A.M. Ali, Mohamed G. El Barbary (2021)** Productivity of Olive Trees as Influenced by Spraying with Glucose, Phosphorus and Gibberellic Acid, *American-Eurasian Journal of Agriculture and Environmental Science*, 21 (2): 117-125.

- M.M., Samira and T.F.; El-Sharony. (2012).** Effect of Gibberellic Acid Spraying on Alternate Bearing of Olive Trees *Journal of Applied Sciences Research*, 8(10): 5114-5123.
- McLaughlin, S.B., Wimmer, R. (1999).** Tansley Review No. 104 Calcium physiology and terrestrial ecosystem processes. *The New Phytologist*, 142(3), 373-417.
- M. Hasani, Z. Zamani, G. Savaghebi, R. Fatahi (2012)** Effects of zinc and manganese as foliar spray on pomegranate yield, fruit quality and leaf minerals, *Journal of Soil Science and Plant Nutrition*, 12(3): 471- 480.
- Mahaveer Suman, Pency D Sangma, Deva Ram Meghawal and Om Prakash Sahu (2017)** Effect of plant growth regulators on fruit crops. *Journal of Pharmacognosy and Phytochemistry*, 6(2): 331-337.
- Mervat S.M. Sourour, Eman E. K. Abd. Ella and Wafaa, A. El- Sisy (2011).** Growth and productivity of Olive tree as influenced by foliar spray of some micronutrients, *Journal of Agriculture and Environmental Science*, Alex University, Egypt, 10(2).
- Muengkaew R, Chaiprasart P and Wongsawad P, (2002)** Calcium, Boron addition Promotes Pollen Germination and Fruit set in Mango, *International Journal of Fruit Science*.
- Muhammad Sajid, Abdur-Rab, Nawab Ali, Muhammad Arif, Louise Ferguson and Masood Ahmed (2010).** Effect of foliar application of zinc and boron on fruit production and physiological disorders in sweet orange cv. blood orange. *Sarhad Journal of Agriculture*, 26(3): 355-360.
- Muhammad Ali, Mukhtar Ahmad, Naveed Anjam, Hafeez-ur-Rehman, M. Imran Kasana, Sudheer Tariq and Aasia Ramzan (2014)** Efficacy of boron & gibberellic acid on growth and fruit yield in olive (*Olea europaea* L.) cv. Gemlik. *International Journal of Biology and Biotechnology*, 11 (2-3): 295-298.
- Mukhtar Ahmad, Muhammad Ali Khan, Hafeez-Ur-Rahman, Niaz Ahmad, Sudheer Tariq and Aasia Ramzan (2011)** effect of boron and gibberellic acid on growth and fruit yield of Olive cv. "Uslu". *International journal of biology and biotechnology* 8 (1): 123-126.
- Osama, H. M. El Gammal (2022)** Effect of Zinc and Boron on yield and fruit quality of Manzanillo Olive trees under Siwa oasis conditions, *Alexandria Science Exchange Journal* 4.
- P.P.S. Gill and J.S. Bal (2009).** Effect of growth regulator and nutrients spray on control of fruit drop, fruit size and quality of Ber under sub-montane zone of Punjab, *Journal of Horticultural Sciences*, 4(2): 161-163.
- P. Sayyad-Amin, A.R. Shahsavar, E. Aslmoshtagh (2015).** Study on foliar application nitrogen, boron and zinc on olive tree, *Trakia Journal of Sciences*, 2:131-136.

- Poornima Devi, R.K.S. Gautam, Jyoti Singh, Sunil Kumar Maurya and Aneeta Chaudhary (2019).** Effect of foliar application of NAA, GA3 and zinc sulphate on fruit drop, growth and yield of ber (*Zizyphus mauritiana* Lamk.) cv. Banarasi Karaka. *International Journal of Current Microbiology and Applied Sciences*, 8(1): 1679-1683.
- Raghvendra Sharma And Rajesh Tiwari (2015).** Effect of growth regulator sprays on growth, yield and quality of guava under malwa plateau conditions, *Annals of Plant and Soil Research*, 17(3): 287-291.
- Rajkumar, J. P. Tiwari and Shant Lal (2014).** Effect of foliar application of zinc and boron on fruit yield and quality of winter season guava (*Psidium guajava*) cv. Pant Prabhat *Annals of Agri Bio Research*, 19(1):105-108.
- S.N. Ghosh, B. Bera, S. Roy, A. Kundu and S.K. Dutta Roy (2009).** Effect of nutrients and plant growth regulators on fruit retention, yield and physio-chemical characteristics in Aonla cv. NA-10, *Journal of Horticultural Sciences*, 4(2): 164-166.
- Saadati S, Moallemi N, Mortazavi SMH, Seyyednejad SM (2016).** Foliar Applications of Zinc and Boron on Fruit Set and Some Fruit Quality of Olive. *Vegetos*, 29:2.
- Sabour, A.M.T. (2001).** Influence of Calcium and Zinc on controlling the inflorescence malformation and improving fruiting of *Hindy bisinnara* mango trees. *Journal of Agricultural Science Mansoura University*, 26(4): 2263-2269.
- Sadrollah Ramezani and Akhtar Shekafandeh (2009)** Roles of gibberellic acid and zinc sulphate in increasing size and weight of Olive fruit, *African Journal of Biotechnology* 8 (24): 6791-6794.
- Sadrollah Ramezani, Akhtar Shekafandeh & Mohammad Reza Taslimpour (2010).** Effect of GA3 and Zinc Sulphate on Fruit Yield and Oil Percentage of 'Shengeh' Olive Trees, *International Journal of Fruit Science*, 10(3): 228-234.
- Sayyad-Amin, A.R. Shahsavar, E. Aslmoshtagh (2015).** Study on foliar application nitrogen, boron and zinc on Olive tree, *Trakia Journal of Sciences* 2:131-136
- Shabaq M. Nafea, Halala K. Abdulfatah (2014)** effect of Foliar application of GA3 and NAA for reducing alternate bearing of Olive trees (*Olea europaea* L. cv. Ashrasie) *Journal of Agriculture and Veterinary Science*,7(1): 08-12.
- Shaheen, S.A.A., (1995).** Effect of foliar sprays of some nutrients on flowering and fruiting of Olive trees. M.Sc. Thesis. Faculty of Agriculture., Cairo Univ. pp: 122.
- Sheikh K.H.A, Singh, B. Haokip, S.W. Shankar, K. Debbarma, R. Gaitri Devi, A. Nengparmoi, T.H. (2021).** Response of Yield and Fruit Quality to Foliar Application of Micronutrients in Lemon [*Citrus limon* (L.) Burm.] cv. Assam Lemon, *Journal of Horticultural Sciences*, 16(2).

- Shereen A. and Hadeer M. Abo-Eloun (2018).** Effect of Foliar Application some Plant Growth Regulators on Growth and Productivity of Olive Trees under South Egypt Conditions. *Journal of Plant Production*, Mansoura University, 9(10): 775 – 783.
- Singh, Anshuman and Singh, H. K. (2015)** Application of Plant Growth Regulators to improve Fruit Yield and Quality in Indian Gooseberry (*Emblica officinalis* Gaertn.). *Journal of AgriSearch*, 2(1): 20-23.
- Slavko Perica, Patrick H. Brown<sup>1</sup>, Joseph H. Connell, Agnes M.S. Nyomora, Christos Dordas, and Hening hu, (2001)** Foliar boron application Improves flower fertility and fruit set of Olive, *HortScience* 36(4):714–716.
- Swatantra Yadav, J.P. Singh, Shubham Gupta and Jitendra Singh Yadav (2021).** A Study on Foliar Feeding of GA<sub>3</sub> and NAA on Fruit Drop, Retention, Yield and Quality of Ber Fruit (*Ziziphus mauritiana* Lamk.) cv. “Banarasi Karaka” *Biological Forum – An International Journal* 13(3): 608-612.
- Talaie A, Badm Mahmoud MT, Malakout MJ (2001).** The effect of foliar application of N, B and Zn on quantitative and qualitative characteristics of olive fruit. *Iranian Journal of Agricultural Science* 32: 727-736. *Vegetos* 29:2.
- Wiesman, Z., A. Ronen A, Ankarion Y, Maranz S, Chpagain B and Abramovich Z, (2002).** Effect of Olive-Nutri-vant on yield and quality of Olives and oil. *Acta Horticulturae*, 594: 557-562.
- Yadav et al., (2022)** Effect of Foliar Application of Micronutrients on Yield and Quality of Pomegranate. *International Journal of Bio-resource and Stress Management*, 13(9): 914-920.
- Yogendra Singh, Nidhika Thakur and Nirmal Kumar Meena (2018)** Studies on the effect of foliar spray of Zn, Cu and B on growth, yield and fruit quality of sweet orange (*Citrus sinensis* L.) cv. Mosambi, *International Journal of Chemical Studies*, 6(5): 3260-3264.