

Original Research Article

Fast Food Consumption and Dental Caries among Teenagers

ABSTRACT

Background: Dental caries in teenagers is significant public health concern. Fast food consumption among teenagers in Bangladesh is steadily increasing. There is a close relationship between fast food consumption and dental caries. The objectives of the study were to determine the association between fast food consumption and dental caries among teenagers.

Methods: This cross-sectional study was conducted from January to December 2022. Sample size was 345 teenagers from two secondary schools in Dhaka city. The convenience sampling technique was used to collect data through face-to-face interviews with a pretested, semi-structured questionnaire. A checklist was also used to collect data through clinical examination of the oral cavity. Data were edited for quality control by checking and rechecking. Statistics software was used to analyze the data.

Results: Among 345 teenagers, the mean \pm SD age was 14.88 \pm 1.17 years. The majority of the teenagers (53.6%) were female and Muslim (95%). Around 59.4% of the teenagers' fathers were business owners, while 37.4% were government employees. The majority of respondents (73.3%) ate breakfast before going to school, 52.2% ate tiffin from the school canteen, and 54.8% ate snacks once a day. The majority (93.3%) ate at an outside restaurant, with 90.4% eating fast food. 59.7% of people ate fast food at least once a week. 53% of teenagers brushed their teeth on a regular basis. Dental caries affected 36.8% of the teenagers. 98.0% of teenagers who ate fast food had dental caries, which was statistically significant. ($p < 0.05$). A significant association was discovered between dental caries and fast food consumption such as Burger (75.2%), Pizza (60%), Sandwich (40.8%), and Chicken Grill (60.0%) per week ($p < 0.05$). There was also a significant correlation (96.0%) between dental caries and regular tooth brushing. ($p < 0.05$)

Conclusion: To prevent dental caries, teenagers should avoid eating fast food and snacks between meals and brush their teeth on a regular basis.

Keywords : *Dental caries, fast food, teenagers.*

1. INTRODUCTION

Globalization has resulted in a nutritional shift away from home cooking and toward fast food, which has contributed to an increase in dental caries among teenagers [1]. Fast food lacks micronutrients such as vitamins, minerals, amino acids, and fiber. Fast food refers to foods that are perceived to have little or no nutritional value. It contains a lot of refined sugar, white flour, trans fat, poly saturated fat salt, and a lot of food additives. It also lacks proteins, vitamins, minerals, and fiber [2]. Diet and nutrition impact oral cavity growth, health, and disease development, affecting tooth integrity, pH, saliva, and plaque makeup [3]. The frequency, with which cariogenic foods are consumed, particularly between meals, is strongly related to the risk of tooth decay [4]. Dental caries is a multifaceted disease. Dental caries remains a major public health issue around the world. Dental caries is increasing in both developed and developing countries. Caries prevalence has been reported to be up to 12% in affluent countries. In Bangladesh, dental caries is a prevalent oro-dental issue. Dental caries affects almost 40% of children under the age of five. It has been observed that around 88% of people aged 13 to 22 are missing one or more permanent teeth as a result of dental caries or a related condition [5]. The aim of the study was to determine the association between fast food consumption and dental caries among teenagers.

2. METHODS

This was a cross-sectional study that was carried out from January 2022 to December 2022. The study place was P.M Pilot boys' school, Zinzira, Keranigong, Dhaka. And P.M Pilot girls' school, Zinzira, Keranigong, Dhaka. The study population comprised of teenagers aged 13 to 19 years from above mentioned institutions in Dhaka district. Convenience sampling technique was used for sample selection. Face-to-face interviews were conducted using a semi-structured questionnaire and a clinical examination checklist. Before data collection pretesting was done among 35 teenagers at Shahid Cadet Academy School, Keranigonj, Dhaka to check the accuracy and degree of reliability of the questionnaire. To ensure consistency, the data were manually checked and edited after collection, and then coded, entered and analyzed with SPSS 22. The descriptive analysis made use of frequency, mean, standard deviation, and percentage. Chi-square test used to determine the association between fast food consumption and dental caries among teenagers.

3. RESULTS

In our study, among 345 students, 125 (36.3%) respondents belonged to 13-14 years and 220 (63.7%) belonged to 15- 17 years (Table 1). Whereas the majority of respondents (322 (93.3%) ate at a restaurant, 23 (6.7%) did not eat at a restaurant (Table 2).

Table 1: Distribution of the respondents according to their age (n-345)

Age of respondents	Frequency	Percentage	Mean \pm SD*
13 - 14 Years	125	36.3	14.88(\pm 1.17)
15- 17 Years	220	63.7	
Total	345	100.0	

*SD= Standard deviation

Table 2: Respondent distribution at an outside restaurant (n-345)

Eating in outside Restaurant	Frequency	Percentage
Yes	322	93.3
No	23	6.7
Total	345	100.0

Most of the respondents 312 (96.9%) liked to eat fast foods in restaurants, 21 (6.5%) of the respondents ate Chinese food in restaurants, 8 (2.5%) of the respondents ate Indian food in restaurants, 19 (6%) of the respondents ate Bangla food in outside restaurants (Table 3). Table 4 also revealed that majority of the respondents 206 (59.7%) ate fast food weekly, 16 (4.6%) respondents ate fast food daily, 38 (11.0%) respondents ate fast food monthly, 52 (15.1%) respondents ate fast food occasionally.

Table 3: The distribution of respondents based on the types of foods consumed in outside restaurants(n-322)

Types of outside Restaurants	Frequency	Percentage
Chinese food	7	2.2
Indian food	2	0.6
Fast food	312	96.9
Bangla food	1	0.3
Total	322	100.0

Table 4: Distribution of respondents based on frequency of fast food consumption (n-312)

Frequency of fast food consumption	Frequency	Percentage
Daily	16	4.6
Weekly	206	59.7
Monthly	38	11.0
Occasionally	52	15.1
Total	312	100.0

According to Table 5, the majority of the respondents, 218 (63.2%), had no caries, while 127 (36.8%) had caries. Table 6 shows the relationship between dental caries and respondents' preference for eating out. The findings revealed a significant relationship between dental caries and respondents' preference for eating out. ($p < 0.05$). Table 7 shows the relationship between dental caries and frequency of fast food consumption. The results showed that there was a significant relationship between dental caries and frequency of fast food consumption ($p < 0.05$).

Table 5: The distribution of respondents according to the presence of dental caries. (n-345)

Dental caries	Frequency	Percentage
Present	127	36.8
Absent	218	63.2

Total	345	100.0
--------------	------------	--------------

Table 6: Association between dental caries and the respondents' preference to eat at an outside restaurant(n=345)

Preference to eat at an outside restaurant	Dental caries		Total f(%)	Statistics
	Yes f (%)	No f(%)		
Yes	125(36.23%)	195(56.52%)	320(92.8%)	Chi-square test $\chi^2 = 9.619$ p=0.001 (p<0.05)
No	2(0.57%)	23(6.66%)	25(7.2%)	
Total	127	218	345	

*Statistically significant

Table 7: Association between dental caries and frequency of eating fast foods of the respondents (n=312)

Frequency Eating fast foods	Dental caries		Total f(%)	Statistics
	Yes f(%)	No f(%)		
Daily	154(4.8%)	1(0.3%)	16(5.1%)	Chi-square test $\chi^2 = 24.256$ p=0.000 (p<0.05)
Weekly	80(25.6%)	126(40.4%)	206(66.0%)	
Monthly	10(3.2%)	28(9.0%)	38(12.2%)	
Occasionally	16(5.1%)	36(11.5%)	52(16.7%)	
Total	121	191	312	

*Statistically significant

4. DISCUSSION

According to the findings of this study, 125 (36.3%) of 345 respondents were between the ages of 13 and 14, while 220 (63.7%) were between the ages of 15 and 17. In a similar study conducted in India in 2017, the majority (51.8%) of the participants were under the age of 12 [6]. Another study conducted in Hanoi, Vietnam in 2021 found that 1153 (58.1%) of the respondents were between the ages of 11 and 12 years old, and 832 (41.9%) were between the ages of 13 and 14 years old. [7]. The majority of the respondents (312 (90.4%)) ate fast food, while 33 (9.6%) did not like fast food. Similar research was done in Dhanmondi, Bangladesh, where 197 (98.5%) students responded about their choice for fast food, with about 22% reported they ate it four days per week and 21.3% reporting they ate it every day of the week.

[3].The majority of respondents 206 (59.7%) ate fast food weekly, 16 (4.6%) ate fast food daily, 38 (11.0%) ate fast food monthly, 52 (15.1%) ate fast food occasionally, and 33 (9.6%) did not eat fast food. A similar study conducted in Benghazi City, Libya, discovered that more than two-thirds of the study population (68.5%) consumed fast foods three times or more per week, while 17% consumed fast foods at least once a week, and (14.5%) twice weekly. Furthermore, 63.5 percent of participants consume fast food on a daily and weekly basis, compared to 28 percent during holidays, 8.3 percent during celebrations, and only 0.2 percent during vacation [8]. Caries were absent in 218 (63.2%) of the respondents, while they were present in 127 (36.8%) of the respondents. In a similar study conducted in Tamil Nadu, India, 80% of 59 students in rural schools had caries and 20% did not. In an urban school, 84% of 115 students had caries, while 16% had none. Caries affects more urban students than rural students. [3]. Another study was conducted in Chennai, India, among 916 teenagers, and the prevalence of caries was found to be 36%, which was very similar to the current study [9]. This study also found a link between dental caries and respondents' preference for eating out. The findings revealed a significant relationship between dental caries and respondents' preference to eat in restaurants ($P<0.05$). This study found a significant association between dental caries and frequency of eating fast foods ($P<0.05$). In terms of dietary habits, both eating breakfast and eating out frequently were strongly linked to dental caries ($P<0.001$) [10].

8. ASSENT

All authors declare that "written informed assent from the participant was obtained for publication of this research report and accompanying images." This journal's Editorial Office/Chief Editor/Editorial Board members can review a copy of the written consent

9. ETHICAL APPROVAL

Ethical approval was taken from the IRB of National Institute of Preventive and Social Medicine (NIPSOM), Bangladesh.

10. REFERENCE

1. Athavale P, Khadka N, Roy S, Mukherjee P, Chandra Mohan D, Turton B, Sokal-Gutierrez K. Early childhood junk food consumption, severe dental caries, and undernutrition: a mixed-methods study from Mumbai, India. *International journal of environmental research and public health*. 2020 Nov;17(22):8629.
2. Das JC. Fast food consumption in children: A review. *Medical & Clinical Reviews*. 2015;1(1):1-4.
3. Gunaseelan BV, Gunasekaran V. Prevalence of Dental Caries among Junk Food-and Non-junk Food-eating Schoolchildren in Urban and Rural Areas of Mayiladuthurai, Tamil Nadu, India. *Journal of Scientific Dentistry*. 2020 Jan;10(1):7-9.
4. Punitha VC, Amudhan A, Sivaprakasam P, Rathanaprabu V. Role of dietary habits and diet in caries occurrence and severity among urban adolescent school children. *Journal of pharmacy & bioallied sciences*. 2015 Apr;7(Suppl 1):S296.
5. Khan MI, Polan M, Nahar A, Raihan M. Factors related to dental caries among the patients attending at the outpatient department (OPD) of Dhaka Dental College and Hospital. *Bangladesh J. Dent. Res. Educ*. 2013 Oct 14;3:16-20.
6. Asawa K, Sen N, Bhat N, Tak M, Sultane P, Patil V. Association of sugary foods and drinks consumption with behavioral risk and oral health status of 12-and 15-year-old Indian school children. *Journal of education and health promotion*. 2018;7.
7. Van Chuyen N, Van Du V, Van Ba N, Long DD, Son HA. The prevalence of dental caries and associated factors among secondary school children in rural highland Vietnam. *BMC Oral Health*. 2021 Dec;21:1-7.
8. TAGURI HS, BUZGEIA MH, HAMZA MS. Fast Foods Consumption in Benghazi City, Libya (Doctoral dissertation, University of Benghazi).

9. Punitha VC, Amudhan A, Sivaprakasam P, Rathana prabu V. Role of dietary habits and diet in caries occurrence and severity among urban adolescent school children. Journal of pharmacy & bioallied sciences. 2015 Apr;7(Suppl 1):S296.

10. Bae JH, Obounou BW. Presence of dental caries is associated with food insecurity and frequency of breakfast consumption in Korean children and adolescents. Preventive nutrition and food science. 2018 Jun;23(2):94.

UNDER PEER REVIEW