

Original Research Article

EFFECT OF INTEGRATED NITROGEN MANAGEMENT ON GROWTH AND YIELD OF OKRA AT SUNDARBAZAR, LAMJUNG, NEPAL

ABSTRACT

Integrated Nutrient Management (INM) is a crucial approach to improving the sustainable yield of crops in an environmentally friendly manner. In this study, a field experiment was conducted at Lamjung Campus, Sundarbazar, Lamjung, from March 2022 to June 2022 to investigate the effect of integrated nitrogen management on the growth and yield of okra (cv. Arka Anamika). A randomized full block design was utilized in the investigation, with seven treatments and three replications. The treatments consisted of a control (T₁) and six other combinations of nitrogen sources, with each treatment receiving 50% of its nitrogen from chemical fertilizers and 50% from organic sources. The amount of organic manure was calculated based on the nitrogen content of the manures. Farmyard manure (FYM), poultry manure, vermicompost, mustard cake, and goat manure were tested as organic sources. The effect of treatment combinations was evaluated in terms of plant height, stem diameter, leaf numbers, days to first flower opening, days to first fruit maturity, number of pods per plant, length, and diameter of pods, and yield ha⁻¹. The findings revealed that integrated nitrogen management strategies had a substantial impact on okra growth and output. Among all the treatments, 50% recommended N through chemical fertilizer + 50% N through poultry manure (T₄), which gave the highest number of pods per plant (19.87), yield per ha (13.59 t/ha), and shortest days taken to first flowering (46.54 days) and plant height (54.80 cm), followed by T₂, while the control treatment (T₁) yielded the lowest. These findings suggest that the application of 50% recommended N through chemical fertilizer + 50% N through poultry manure is an effective strategy for obtaining a high yield of okra in the study area.

Keywords: Growth, Nitrogen, Okra, Poultry manure, Yield

1. INTRODUCTION

Okra is one of the well-known vegetable crops that belong to the genus *Abelmoschus*, family Malvaceae (Siemonsma, 1982) having somatic chromosome number of Okra $2n=130$ (Baviskar et al., 2011). It was initially planted in Egypt in the 12th century after being discovered in tropical America (Maurya et al., 2013). It is called Lady's Finger in England, Gumbo in the United States of America, and Bhindi in Nepal. It is a short-duration hardy vegetable crop; the immature green fruits are used as a vegetable and are generally marketed in fresh but sometimes in canned or dehydrated form. It contains plenty of vitamins, calcium, potassium, and other minerals. The present consumption of vegetables is 195g/day/capita which is very low compared to the recommended dose by dietitians 375g/day/capita (FAO/WHO, 2003). Therefore, there is a big gap between the requirement and the supply of vegetables in Nepal. Successful okra production may contribute partially to solving vegetable scarcity in summer. Over a total area of 9,584 hectares, the nation produced 110,565 metric tons of okra, with productivity of 11.54 tons per hectare (MOAD, 2021).

Despite its yield potential, it has not yet achieved optimum yields due to a continuing deterioration in soil fertility (Owa et al., 2015) which could be attributed to the imbalanced use of fertilizers in the mid-hill of Nepal. Crop nutrient requirements are determined by soil texture, preceding vegetation cover type, cropping intensity, and soil moisture (Denton & Swarup, 1983). Nitrogen (N) is the most crucial, among various nutrients required for adequate nutrition and a high yield of okra (Owa et al., 2015). Inappropriate nitrogen sources, uneven fertilizer application, and a high rate of nitrogen leaching are all factors in low crop yields (Akhtar et al., 2010). Nitrogen's primary role is to contribute to plant growth characteristics, thereby influencing yield and quality parameters, which are generally directly correlated (Sati et al., 2018) and it is major essential for the synthesis of chlorophyll, protein, nucleic acid, hormones, and vitamins, as well as cell division and elongation (Firoz, 1970). Between 1997 and 2003, the use of nitrogen fertilizers for vegetable production increased by 21% (Mubashir et al., 2010).

Unfortunately, N deficiency is widespread in Nepal on account of low available soil N and organic matter content as a result of nitrification and leaching losses. Imbalanced application of chemical fertilizers for obtaining higher yields is linked to soil acidity, poor physical structure, and nutrient retention characteristics, which negatively affect crop growth and yield. Moreover, it increases the cost of production and causes economic loss (Saurabh et al., 2021). The use of organic manure should be promoted to sustain soil fertility over time. This is because the nutrients in organic manures

are released more slowly and are kept in the soil for a longer period, ensuring a lasting residual effect (Sharma & Mitra, 1991). According to F et al., 2016, it has also been reported that organic manure has a direct impact on plant growth by providing all necessary macro and micronutrients in accessible forms during mineralization, hence increasing the physical and physiological qualities of soils. There are different types of nitrogen sources including farmyard manure, poultry manure, goat manure, vermicompost, mustard cake, etc. Combining the use of organic and chemical fertilizers increases yields, improves soil health, and reduces pollution problem issues brought on by improper use of chemical fertilizers alone (Basnet et al., 2021). Keeping all the above facts in view this experiment was conducted under mid-hill condition of Nepal to evaluate the effect of integrated nitrogen management on the growth and yield parameters of okra and to find out the suitable nitrogen management system in okra.

2. MATERIALS AND METHODS

2.1. Experimental site and soil analysis

The experiment was laid out in the field of horticulture research farm of Lamjung Campus, Sundarbazar, Lamjung. Geographically, the experimental field is situated at 28° 07' 01.49" N latitude, 82° 17' 48.40" E longitude, and an elevation of 857 masl. The duration of the experiment was from March 2022 to June 2022. The location falls in the mid-hill area with a humid subtropical climate.

The soil of the studied area was tested before commencing the experiment, and soil falls on the texture loam having a pH value of about 5.06, organic matter 1.826, total nitrogen (0.09%), available phosphorus (6.36 kg/ha), and available potassium (125.6 kg/ha). Moisture percentage was recorded at 25.85%.

2.2. Experimental materials

The Arka Anamika variety of okra, which was registered in Nepal in 2010, was used in the experiment (Krishi Diary, 2078). The seed of Arka Anamika was taken from Krishi Sahayog Kendra Tatha Biyu Bhijan Agro-Vet Sundarbazar, Lamjung.

2.3. Experimentation

The experiment was laid out in Randomized Complete Block Design (RCBD) with seven treatments and three replications. The net area of the experimental field was 84.6 m² (14.1 m × 6 m) in which the individual plot measured 2.7m² (1.8m × 1.5 m). There were 21 plots in

total, with a spacing of 0.25m between each plot and a spacing of 0.75m between each replication. The recommended dose of fertilizer for okra is 160 N: 120 P₂O₅: 60 K₂O as per Kanhaiya Prasad Singh, 2019 which was used in the study.

Table 1: Treatment details of the experimental field

SN	Treatment	Details
1	T ₁	Control
2	T ₂	100% recommended N through chemical fertilizer
3	T ₃	50% recommended N through chemical fertilizer + 50% N through farm yard manure (FYM)
4	T ₄	50% recommended N through chemical fertilizer + 50% N through poultry manure
5	T ₅	50% recommended N through chemical fertilizer + 50% N through vermicompost
6	T ₆	50% recommended N through chemical fertilizer + 50% N through mustard cake
7	T ₇	50% recommended N through chemical fertilizer + 50% N through goat manure

One-month-old, decomposed poultry manure, FYM, goat manure, vermicompost, and freshly prepared mustard cake were collected and sun-dried to reduce moisture. Nutrient content in manures was tested in the regional Soil and Fertilizer Testing Laboratory, Pokhara, Kaski. The quantities of organic manures for the experiment were calculated based on their respective nitrogen content.

Table 2. Nutrient content of organic manures

Amending substances	Nutrients containing %		
	N	P	K
FYM	0.5%	0.5%	0.5%

Poultry manure(layers)	2.85%	4.1%	2%
Goat manure	2.21	0.77	3.89
Vermicompost	2.52	1.69	2.08
Mustard Cake	4.61	1.5	1.3

2.4. Cultural activities

Cultural practices, such as seed and sowing, irrigation, gap filling, weeding, plant protection measures, and picking of fruits, were followed in the experiment. Water-primed seeds were sown on March 4th with two seeds placed in each spot to reduce missing plants, and the soil was compacted over the seeds to facilitate germination. Light irrigation was applied immediately after sowing, and medium irrigation was done every second day until the first month, after which heavy irrigation was done every week. Gap filling was done 16 days after sowing (DAS) using plants from the same treatment as filler plants. Weeding was done three times, at 30 DAS, 40 DAS, and 50 DAS, while plant protection measures, such as spraying of Imidacloprid 17.5 SL (0.25%) to protect against insect pests like jassids and whitefly and spraying of Dimethoate 30 EC (0.03%) to control fruit borers in okra crops, were applied as needed. The fruits were picked manually when they were green tender and at a marketable size and were immediately weighed and subjected to other observations.

2.5. Harvesting and data collection

Five randomly selected plants from each plot were used to collect various types of growth and yield contributing data for okra and their average value was considered as one plot for each parameter. Plant height, stem girth, number of leaves per plant, Days to first flowering, and days to first fruit maturity were measured at different intervals after sowing. Fruits were collected every third day after the first harvest up to 20 times. The number of fruits per plant, fruit length, fruit girth, and individual fruit weight were determined at each harvest. Yield per plot was converted to ton per hectare

2.6. Statistical analysis

Data regarding growth and yield traits were analyzed statistically by one-way analysis of variance (ANOVA) by using R- Studio 4.2.2. Treatment means were separated by least

significant differences (LSD) at 5%, 1%, and 0.1% probability outlined by Gomez & Gomez, 1984.

3. RESULTS

3.1. Growth attributes

3.1.1. Germination%

Table 3. Effect of integrated nitrogen management on germination Percentage of okra

Treatments	Germination %
T ₁	62.43 ^b
T ₂	65.53 ^{ab}
T ₃	65.20 ^{ab}
T ₄	68.60 ^a
T ₅	65.06 ^{ab}
T ₆	65.46 ^{ab}
T ₇	65.41 ^b
SEM (±)	1.20
LSD (α=0.05)	3.72
CV	3.20%
Grand mean	65.24
p-value	NS

*Means in column followed by similar letter/s are not significantly different, CV=coefficient of variation, NS= non-significant; *= p<0.05; **p<0.01; ***= p<0.001 in LSD*

There was a non-significant difference in germination percentage among a different combination of inorganic and organic manures but T₄ (50% recommended N through chemical fertilizer + 50% N through poultry manure) recorded the maximum germination percentage which was 68.60 % followed by 65.53% at T₂ (50% recommended N through chemical fertilizer + 50% N through FYM). The minimum plant germination percentage (62.43 %) was recorded at T₁ (control).

3.1.2. Plant height, number of leaves per plant, and stem diameter

Table 4. Effect of integrated nitrogen management on the growth attributes of okra

Treatments	Plant height(cm)	Number of leaves per	Stem diameter(mm)
------------	------------------	----------------------	-------------------

	plant								
	30	45	60	30	45	60	30	45	60
	DAS	DAS	DAS	DAS	DAS	DAS	DAS	DAS	DAS
T ₁	10.15 ^c	23.02 ^c	40.13 ^c	3.36 ^e	6.345 ^c	10.85 ^c	3.21 ^c	6.91 ^c	11.11 ^c
T ₂	11.97 ^{bc}	27.04 ^{abc}	45.30 ^b	3.95 ^b	6.980 ^{bc}	12.54 ^{bc}	3.74 ^{bc}	8.56 ^{ab}	12.83 ^b
T ₃	13.53 ^{ab}	28.48 ^{ab}	45.46 ^b	3.85 ^{bc}	7.79 ^b	13.06 ^b	4.30 ^{ab}	8.27 ^{ab}	14.26 ^{ab}
T ₄	15.39 ^a	31.32 ^a	54.80 ^a	4.32 ^a	9.39 ^a	14.70 ^a	5.10 ^a	9.29 ^a	15.50 ^a
T ₅	12.71 ^{bc}	28.36 ^{ab}	44.80 ^{bc}	4.04 ^b	7.40 ^b	11.40 ^{bc}	4.52 ^{ab}	8.80 ^{ab}	13.38 ^b
T ₆	11.67 ^{bc}	25.25 ^{bc}	44.26 ^{bc}	3.52 ^{de}	7.03 ^{bc}	11.66 ^{bc}	3.61 ^{bc}	7.73 ^{bc}	14.25 ^{ab}
T ₇	11.96 ^{bc}	25.14 ^{bc}	42.47 ^{bc}	3.66 ^{cd}	7.26 ^{bc}	11.70 ^{bc}	4.07 ^{bc}	8.19 ^{ab}	12.83 ^b
SEM (±)	0.8043	1.528	1.453	0.0838	0.314	0.518	0.277	0.341	0.528
LSD	2.478	4.70	4.479	0.258	0.968	1.597	0.855	1.05	1.63
($\alpha=0.05$)									
CV	11.15%	9.82%	5.55%	3.80%	7.29%	7.32%	11.76%	7.16%	6.79%
Grand mean	12.48	26.94	45.31	3.81	7.45	12.27	4.08	8.25	13.46
p-value	*	*	***	***	***	**	**	**	**

Means in column followed by similar letter/s are not significantly different, CV=coefficient of variation, NS= non-significant; *= $p<0.05$; **= $p<0.01$; ***= $p<0.001$ in LSD

3.1.3. Plant height

The vegetative growth parameters of plant height were influenced significantly at 30, 45, and 60 DAS due to different treatments of integrated nitrogen management as shown in Table 4. The highest plant height was recorded in T₄ at 60 DAS (54.80cm) while the lowest plant height was recorded in Control treatment T₁ (40.13 cm).

3.1.4. Number of leaves per plant

The number of leaves per plant also shows significant differences when the okra plant was treated with various treatments of integrated nitrogen management. The maximum number of leaves per plant was observed in T₄ at 60 DAS (14.70) while the minimum number of leaves per plant was seen in the control treatment i.e., T₁ (10.85).

3.1.5. Stem diameter (mm)

Integrated nitrogen management exerted a significant difference in the stem diameter. The maximum stem diameter was found in the treatment T₄ at 60 DAS (15.50 mm) which was seen to be statistically par with T₃ (14.26 mm) and T₆ (14.25 mm). The maximum stem diameter was then followed by T₂, T₇, and T₁ respectively. The minimum stem diameter was found in the control treatment (11.11 mm).

3.2. Reproductive attributes

Table 5. Effect of integrated nitrogen management on the reproductive attributes of okra

Treatments	Days to first flower opening	Days to first fruit maturity
T ₁	53.37 ^a	59.37 ^a
T ₂	48.53 ^d	54.70 ^{cd}
T ₃	48.88 ^{cd}	54.83 ^{cd}
T ₄	46.54 ^e	52.54 ^e
T ₅	48.26 ^{de}	54.07 ^{de}
T ₆	52.67 ^b	57.04 ^b
T ₇	50.46 ^{bc}	56.11 ^{bc}
SEM (±)	0.5637	0.5522
LSD (α=0.05)	1.7372	1.7016
CV	1.9694%	1.7226%
Grand mean	49.583	55.526
p-value	***	***

*Means in column followed by similar letter/s are not significantly different, CV=coefficient of variation, NS= non-significant; *= p<0.05; **p<0.01; ***= p<0.001 in LSD*

3.2.1. Days to first flower opening

The results revealed that the different combinations among treatments show significant differences in the days of the first flower opening. The least days to took to first flowering was observed when plants were treated with (T₄) 50% recommended N through chemical fertilizer + 50% N through poultry manure (46.54 days) which was statistically par with treatment (T₅) 50% recommended N through chemical fertilizer + 50% N through

vermicompost (48.26 days). The maximum days it took to first flowering was observed in the control treatment T₁ (53.37 days).

3.2.2. Days to first fruit maturity

There was a significant difference in days to first fruit maturity among various treatments of combined use of inorganic and organic manures. (T₄) 50% recommended N through chemical fertilizer + 50% N through poultry manure shows the least days to take to first fruit maturity (52.54 days) which was statistically similar to the treatment T₅ (54.07 days). While control Treatment T₁ (59.37 days) took the maximum of days to first fruit maturity.

3.3. Yield attributes

Table no.6: Effect of integrated nitrogen management on the yield attributes of okra

Treatments	Number of pods per plant	Length of pods(cm)	Diameter of pods(mm)	Yield ton per hectare
T ₁	12.16 ^d	9.523 ^c	10.70 ^c	8.20 ^c
T ₂	17.746 ^{ab}	12.436 ^{ab}	12.31 ^{ab}	11.18 ^b
T ₃	15.67 ^{bc}	11.63 ^b	11.917 ^{ab}	10.24 ^{bc}
T ₄	19.87 ^a	13.74 ^a	12.77 ^a	13.59 ^a
T ₅	17.44 ^{ab}	12.01 ^{ab}	12.14 ^{ab}	10.82 ^b
T ₆	13.62 ^{cd}	11.13 ^b	11.38 ^{bc}	9.72 ^{bc}
T ₇	15.19 ^{bc}	11.37 ^b	11.87 ^{ab}	9.99 ^{bc}
SEM (±)	0.8355	0.55250	0.34280	0.6855
LSD (α=0.05)	2.574	1.7024	1.056279	2.112
CV	9.06%	8.184%	5.001%	11.26%
Grand mean	15.96	11.692	11.872	10.53
p-value	***	**	*	**

*Means in column followed by similar letter/s are not significantly different, CV=coefficient of variation, NS= non-significant; *= p<0.05; **p<0.01; ***= p<0.001 in LSD*

3.3.1. Number of pods per plant

Integrated application of organic and inorganic fertilizers showed significant differences in the result on the number of pods per plant. The number of pods *per plant* was highest (19.87) in the treatment T₄ (50% recommended N through chemical fertilizer + 50% N through poultry manure) which was at par with the plots that received 100% recommended N through

chemical fertilizer (T₂) and 50% recommended N through chemical fertilizer + 50% N through vermicompost (T₅). The lowest number of pods per plant was found in control (T₁).

3.3.2. Length of pods (cm)

Integrated nitrogen management on different treatments shows significant differences in the pod length of the okra pod. Maximum pod length was recorded in treatment T₄ (13.74 cm) which was followed by T₂, T₅, T₃, T₇, and T₆ respectively. Treatment T₄ was found statistically par with treatment T₂ (12.43 cm) and treatment T₅ (12.01 cm). The minimum pod length was recorded in the control treatment (T₁) of pod length (9.52 cm).

3.3.3. Diameter of pods (mm)

There was a significant difference in the diameter of pods among different treatments of integrated nitrogen management. The maximum diameter of the pod was observed in treatment T₄ (12.77 mm) which was also statistically similar to treatments T₂ (12.31mm), T₅ (12.14 mm), T₃ (11.91 mm), and T₇ (11.87 mm). The minimum diameter of pods was observed in the control T₁ (10.70 mm).

3.3.4. Yield ton per hectare

The study revealed that nitrogen-based integrated nitrogen management on different treatments exerted a significant difference in the yield of okra. Treatment (T₁) control gave the lowest 8.20-ton yield per hectare while the maximum yield per hectare was observed in treatment T₄ (13.59 tons/ha).

4. DISCUSSION

In comparison to the control plot, the number of pods, fruit weight per plant, pod length, pod diameter, and yield per hectare increased significantly, and high values were observed in the poultry manure treated plot, indicating that poultry manure was available in the best form for easy absorption by the plant roots, increasing the plant's morphological growth. The results of morphological parameters such as plant height, number of leaves, and stem diameter improved significantly. When compared to other sources of organic manure, okra grows better on poultry manure in terms of plant height. This demonstrates that poultry manure was easily accessible and in the best form for root absorption, leading to an increase in the morphological growth of plants which is corroborated by the findings of (Ajari et al., 2004).

The increased number of leaves per plant with poultry manure application in treatments was related to the importance of improved crop plant vegetative growth (Tindall, 1983). The applications of poultry manure may have increased the amount of nitrogen available to plants through mineralization, and nitrogen is known to stimulate plant growth. (Anyaegbu et al., 2010). The effects of integrated nitrogen management on the number of leaves were found to be greatest in poultry manure, correlating with previous research by Fagwalawa & Yahaya, 2016. An increase in stem diameter might be due to an increase in the vegetative growth of the plant. It has been discovered that nutrient concentration influences how many days it takes for flowers to open (Dauda et al., 2008) hence the earlier reproduction of okra under poultry manure could be attributed to the rapid release of nitrogen into the soil (Havlin et al., 2014).

The study of Adekiya & Agbede, 2017 shows the nutrient release may have coincided with the specific phenological demands of okra. The increased number and weight of okra pods as a result of poultry manure application could be linked to the easier solubilization of liberated plant nutrients, resulting in improved nutrient status and soil water-holding capacity. The findings were consistent with those of (Sanwal et al., 2007) in turmeric (*Curcuma longa*) Premsekhar and Rajashree, (2009) in okra (*A. esculentus*) in which they found that superior physical and biological features of the soil, resulting in greater nutrient availability to the plants, might be related to increased crop production response to application of organic manure.

In comparison to other organic manures, poultry manure considerably increased plant growth, yield, mineral content, and proximate composition of okra due to its high soil chemical characteristics, which may be connected to its lowest C: N ratio, lignin, and lignin: N ratio (Adekiya et al., 2020). Similarly, both liquid and solid excreta are excreted without loss of urine due to which poultry manure is richer in organic matter and essential plant nutrients than the manure of other animals (Ewulo, 2005).

Although no significant effect was recorded on the germination percentage of plants, T4 (50% recommended N through chemical fertilizer + 50% N through poultry manure) recorded the maximum germination percentage. Study of Boateng et al., 2009 showed that poultry manure decomposes rapidly and releases all essential nutrients for the crops, which facilitates quick and efficient nutrition uptake by the plants, potentially leading to maximum germination percentage in the treated crops.

CONCLUSION

The combination of 50% recommended N through chemical fertilizer and 50% N through poultry manure management practice resulted in significant improvement in most growth attributes of okra, including yield and yield attributes. Based on the findings, it is suggested to use the combination of 50% recommended N through chemical fertilizer + 50% N through poultry manure to obtain a higher growth and yield of okra in a mid-hill place like Sundarbazar, Lamjung.

REFERENCES

- Adekiya, A. O., & Agbede, T. M. (2017). Effect of methods and time of poultry manure application on soil and leaf nutrient concentrations, growth and fruit yield of tomato (*Lycopersicon esculentum* Mill). *Journal of the Saudi Society of Agricultural Sciences*, 16(4), 383–388. <https://doi.org/10.1016/J.JSSAS.2016.01.006>
- Adekiya, A. O., Ejue, W. S., Olayanju, A., Dunsin, O., Aboyeji, C. M., Aremu, C., Adegbite, K., & Akinpelu, O. (2020). Different organic manure sources and NPK fertilizer on soil chemical properties, growth, yield and quality of okra. *Scientific Reports 2020 10:1*, 10(1), 1–9. <https://doi.org/10.1038/s41598-020-73291-x>
- AjarI, O. I., Tsado, E. K., Oladiran, J. A., & Salako, E. A. (2004). Plant height and fruit yield of okra as affected by field application of fertilizer and benlate in bida, nigeria. *Nigeria Agricultural Journal*, 34(1). <https://doi.org/10.4314/NAJ.V34I1.3173>
- Akhtar, M., Khan, M. Z., Rashid, M., Ahsan, Z., & Ahmad, S. (2010). Effect of potash application on yield and quality of tomato (*Lycopersicon esculentum* Mill.). *Pakistan Journal of Botany*.
- Anyaegbu P O, And, I. U. P., & Ekwugha E U. (2010). 82 Nutrient uptake and root yield of cassava as influenced by liming and poultry manure under different cropping systems. 12 3.
- Basnet, B., Aryal, A., Neupane, A., K.C., B., Rai, N. H., Adhikari, S., Khanal, P., & Basnet, M. (2021). Effect of integrated nutrient management on growth and yield of radish. *Journal of Agriculture and Natural Resources*, 4(2), 167–174. <https://doi.org/10.3126/janr.v4i2.33712>
- Baviskar, M. N., Bharad, S. G., Dod, V. N., & Barne, V. G. (2011). Effect of integrated nutrient management on yield and quality of sapota. *Plant Archives*, 11(2), 661–663.
- Boateng, S., Zickermann, J., & Kornahrens, M. (2009). Poultry manure effect on growth and yield of maize. *West African Journal of Applied Ecology*, 9(1). <https://doi.org/10.4314/WAJAE.V9I1.45682>
- Dauda, S., Ajayi, F., & Ndor, E. (2008). Growth and yield of water melon (*Citrullus lanatus*) as affected by poultry manure application. *Electronic Journal of Environmental, Agricultural and Food Chemistry*.

- Denton, S., & Swarup, V. (1983). Tomato cultivation and its potential in Nigeria. *Acta Horticulturae*, 123, 257–272. <https://doi.org/10.17660/ACTAHORTIC.1983.123.23>
- Ewulo, B. S. (2005). Effect of Poultry Dung and Cattle Manure on Chemical Properties of Clay and Sandy Clay Loam Soil. *Journal of Animal and Veterinary Advances*, 4(10), 839–841. <https://www.researchgate.net/publication/26591139>
- F, F. Z., Shedeed, S. I., & K Hassan, N. M. (2016). A Review of Organic Agricultural of Some Vegetables Crops. *American Journal of Food Science and Health*, 2(3), 25–31. <http://www.aiscience.org/journal/ajfshhttp://creativecommons.org/licenses/by/4.0/>
- Fagwalawa, L., & Yahaya, S. M. (2016). Effect Organic Manure On The Growth And Yield Of Okra. *Imperial Journal of Interdisciplinary Research*.
- Firoz, Z. (1970). Impact of nitrogen and phosphorus on the growth and yield of okra [*Abelmoschus esculentus* (L.) Moench] in hill slope condition. *Bangladesh Journal of Agricultural Research*, 34(4), 713–722. <https://doi.org/10.3329/bjar.v34i4.5846>
- Gomez, A. A., & Gomez, K. A. (1984). Statistical procedures for agricultural research: second Edition. A Wiley-Interscience Publication, 6, 1–690. <https://www.wiley.com/en-us/Statistical+Procedures+for+Agricultural+Research%2C+2nd+Edition-p-9780471870920>
- Havlin, J. L., Tisdale, S. L., Nelson, W. L., & Beaton, J. D. (2014). Soil Fertility and Fertilizers, 8th Edition. *Pearson*.
- Kanhaiya Prasad Singh, R. R. B. (2019). Vegetable Crops Production Technology. [https://www.samikshapublication.com.np/Vegetable Crops Production Technology](https://www.samikshapublication.com.np/Vegetable%20Crops%20Production%20Technology)
- Krishi Diary. (2078). Krishi Diary. *Ministry of Agriculture and Livestock Development*, 356.
- M. Premsekhar V. Rajashree. (2009). *Influence of organic manures on growth, yield and quality of okra*. American-Eurasian Journal of Sustainable Agriculture. https://www.researchgate.net/publication/290005224_Influence_of_organic_manures_on_growth_yield_and_quality_of_okra
- Maurya, R., Bailey, J., & Of, J. C. (2013). Impact of plant spacing and picking interval on the growth, fruit quality and yield of okra (*Abelmoschus esculentus* (L.) Moench). *American Journal of Agriculture and Forestry*, 1(4), 48–54. <https://doi.org/10.11648/j.ajaf.20130104.11>
- Ministry of Agricultural Development. (2021). *Statistical Information on Nepalese Agriculture*. 274.
- Mubashir, M., Malik, S., Khan, A., Ansari, T., Wright, S., Brown, M., & Islam, K. R. (2010). Growth, yield and nitrate accumulation of irrigated carrot and okra in response to nitrogen fertilization. *Pakistan Journal of Botany*.
- Owa, S. K., Etukudo, F. D., & Ibitoye-Ayeni, O. O. (2015). Influence of Different Nitrogen Sources on the Growth and Yield of Three Varieties of Okra (*Abelmoschus esculentus*) in Kabba, Kogi State, Nigeria. *Agricultural Sciences*, 6, 1141–1147. <https://doi.org/10.4236/as.2015.610109>

- Sanwal, S., Laxminarayana, K., Yadav, R., Rai, N., Yadav, D. S., & Bhuyan, M. (2007). Effect of organic manures on soil fertility, growth, physiology, yield and quality of turmeric. *Indian Journal of Horticulture*.
- Sati, U. C., Raghav, M., Singh, D. K., Singh, S., Singh, D. K., & Singh, D. K. (2018). Integrated nitrogen management in okra under Tarai conditions of Uttarakhand. *International Journal of Chemical Studies*, 6(1), 1118–1122.
- Saurabh, Jaiswal, R., Ali, S., & Khandwe, R. (2021). Impact of integrated nutrient management on growth and yield of okra [*Abelmoschus esculentus* (L.) Moench]. *International Journal of Chemical Studies*, 9(2), 302–305. <https://doi.org/10.22271/chemi.2021.v9.i2e.11824>
- Sharma, A. R., & Mitra, B. N. (1991). Effect of different rates of application of organic and nitrogen fertilizers in a rice-based cropping system. *The Journal of Agricultural Science*, 117(3), 313–318. <https://doi.org/10.1017/S0021859600067046>
- Siemonsma, J. S. (1982). The cultivation of okra (*Abelmoschus* spp.), a tropical fruit vegetable, with special reference to the Ivory Coast. *The Cultivation of Okra (Abelmoschus Spp.), a Tropical Fruit Vegetable, with Special Reference to the Ivory Coast*.
- Tindall, H. D. (1983). *Vegetables in the Tropics*. <https://doi.org/10.1007/978-1-349-17223-8>