

Original Research Article

Assessment of Available Major and Micronutrient Status of soils under the barren land in central plain of Uttar Pradesh

Abstract

Soil fertility evaluation of barren land is the most basic decision-making tool for an effective sustainable plan of a particular area. Thus, the present study was carried out to evaluate the soil fertility status in session variation of the two blocks of Kanpur Dehat (Akabrpur and Maitha). The soil samples were randomly collected based on the variability of land at a depth of 0-15 cm, 15-30cm and 30-60 cm in 5-5 sites in both blocks. A GPS device was used to identify the location of the soil sampling points. Soil samples were analyzed for texture, pH, OC, EC, N, P, K, S, Fe, Zn, Cu, Mn, and exchangeable cations status following standard analytic methods in the laboratory of Soil Science and Agricultural chemistry, C.S.A. University of Agriculture and Technology, Kanpur, UP. The soil organic carbon ranged from 0.18 to 0.34% of both blocks. Available nitrogen ranged from 102.78 to 138.39 kg ha⁻¹, available phosphorous ranged from 9.89 to 16.47 kg ha⁻¹ and available potassium ranged from 230.65 to 276.38 kg ha⁻¹ in the surface soil of Maitha, all of which showed decrease in value with increase in depth. Exchangeable calcium ranged from 4.58 to 6.34 (cmol (p+) kg⁻¹), exchangeable magnesium ranged from 2.20 to 4.40 (cmol (p+) kg⁻¹) and the pH of soil in both blocks highly alkaline in nature, all of which varied significantly with site and depth. The results indicated that soils are not good for cultivation of various crops. Farmers are required to maintain Soil Health Card which helps them to adopt suitable management practices and provide proper nutrition to soil.

Key word - Barren land, ~~Maero-nutrients~~ Macronutrients, Micronutrients, GPS, Organic carbon and Suitability.

Introduction

The barren land of the central plain in Uttar Pradesh, India, presents a significant challenge for agricultural productivity and land restoration efforts (Sharma and Singh, 2015). These barren lands are characterized by depleted organic carbon content and poor soil aggregate stability, which severely limit the establishment of vegetation and the overall health of the ecosystem (Mensah, 2015). Understanding the relationship between organic carbon and soil aggregate stability is crucial for devising effective strategies to restore and improve the fertility of these lands (Babur, et al. 2021). Organic carbon plays a fundamental role in soil health and functioning. It is a vital component of soil organic matter, which influences numerous physical, chemical, and biological properties of the soil (Fageria, 2012). Organic carbon content directly affects soil fertility, nutrient availability, water holding capacity, and microbial activity, all of which are critical for plant growth and ecosystem sustainability (Carter, 2002). Moreover, organic carbon is intimately linked to soil aggregate stability, a key indicator of soil structure and resistance to erosion (Allen et al., 2011).

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Soil testing is often used to determine the available nutrient status and nutrient supplying power of soil which helps in developing cost effective nutrient management practices that serve as a basis for amendments and sound fertilizer recommendations, which in turn leads to sustainable long term agriculture production by way of adoption of good agronomic management practices by farmers in the study area (Subbaiah et al 2022). Macronutrients (N, P, K, S) and micronutrients (Zn, Fe, Cu, Mn and B) are very important soil elements that determine crop yields in general and physicochemical properties like pH, EC, Organic Carbon in particular under normal environmental conditions (Dhaliwal, et al., 2019). Soil fertility is one of the important factors controlling crop yields and soil characterization in relation to evaluation of soil fertility of an area or a region is an important aspect in the text of sustainable production (Prasad et al 2020b). In present context, maintaining soil fertility is a key problem in Indian agriculture, especially under the country's rapidly growing population in recent decades. Erratic rainfall, minimal recycling of farm residues including livestock waste, no adoption of soil and water conservation measures, continuous cultivation of exhaustive crops and imbalanced fertilizer application are some of the major causes of soil degradation in India.

The central plain of Uttar Pradesh, known for its fertile soils and agricultural productivity in some areas, also harbors extensive stretches of barren land. These areas often suffer from a combination of factors, including overgrazing, deforestation, unsustainable land management practices, and improper irrigation, leading to severe land degradation (Bhattacharyya, et al, 2015). Restoring these barren lands requires a comprehensive understanding of the role of organic carbon in soil aggregate stability and the underlying processes involved (Harris, 2003). This study aims to investigate the relationship between organic carbon and soil aggregate stability in the barren land of the central plain of Uttar Pradesh. By examining the spatial distribution of organic carbon content and soil aggregate stability, we can identify the ~~area's~~ areas most susceptible to degradation and prioritize restoration efforts. Additionally, we will explore the factors influencing organic carbon accumulation and its impact on soil aggregate stability, such as land use history, soil management practices, and vegetation cover (Udom, and Ogunwole, 2015). The findings from this study will contribute to the development of targeted and sustainable strategies for rehabilitating the barren lands of the central plain of Uttar Pradesh. By enhancing organic carbon content and improving soil aggregate stability, we can promote the establishment of vegetation, increase agricultural productivity, and mitigate the environmental challenges associated with land degradation (Ruiz-Colmenero, et al, 2013). Ultimately, this research aims to support the sustainable development and resilience of the region's agricultural systems and ecosystems.

Formatted: English (United States)

Materials and method

Study sites

The Kanpur Dehat district lies in the center of Uttar Pradesh. It lies between 26° 31' to 350 75.94' N latitude and 790 49' to 840 46.9'' E longitude with total geographical area 3021 sq. km having elevation of about 126 m above the mean sea level. The entire study area was carried out in 10 villages namely Rura, Muridpur, Mandauli, Gutaiya, Barra, Kashipur,

Tikari, Shobhan, Rampur Shiwali and Hathika, falling under two blocks namely Akbarpur and Maithawith five –five different sites taken from villages. Selection of the village was based on its Barren land soil type with the help of GPS, there was not found ~~the~~ vegetation, farming situation and crops grown. The climate in Kanpur is warm and temperate. When compared with winter, the summers have much more rainfall. The climate is classified as Cwa according to Köppen and Geiger (1936). The average annual temperature is 25.3 °C in Kanpur. The annual rainfall is 939 mm/37.0 ~~inches~~. The given location is in the northern hemisphere. The particular spot is situated in the upper half of the planet. Summer starts here at the end of June and ends in September. There are the months of summer: June, July, August and September.

Formatted: English (United States)

Sample collection and Laboratory analysis

The soil samples were collected at different depths of 0- 15 cm, 15-30 cm and 30-45 cm respectively at the site. The collected soil samples were air-dried, ground with wooden pestle and mortar and sieved through 2-mm sieve (0.2 mm sieve for organic carbon), labelled and stored at the Soil Laboratory of the Department of Soil Science and Agriculture Chemistry, C.S. Azad University of Agriculture and technology, Kanpur-208002, UP, India following standard laboratory procedures.

The soil samples were analyzed for soil pH and electrical conductivity (EC) were recorded in 1:2.5 soil to water suspension (Jackson, 1973), the organic carbon (OC) determined by Walkley and Black 1934 method, Exchangeable bases were collected using neutral normal ammonium acetate and the exchanged ion measured following the procedure outlined in Hesse (1970). The Ca⁺⁺ and Mg⁺⁺ was determined by complex metric titration, involving ethylene diamine tetra acetic acid (EDTA) and Na⁺ was determined. The macronutrients like available nitrogen and phosphorus by alkaline potassium permanganate method (Subbiah and Asija, 1956; Sharawat and Burford, 1982) and Spectrophotometer fallow by the method (Olsen et al.1954), respectively. Available potassium was determined by flame photometer with neutral normal ammonium acetate as an extractant (Hanway and Heidel, 1952). Available Sulphur content of the soil samples were determined by turbidimetric method (Black, 1965). Micronutrients viz., available zinc (DTPA extractable) was determined by using Atomic Absorption Spectrophotometer (Lindsay and Norvell, 1978).

Data analysis

Data were subjected to descriptive statistics such as minimum, maximum, mean, standard deviation and coefficient of variation using the SPSS, Ver. 2023.

Result and Discussion

1. Physico-chemical properties of soil in Akbarpur and Maitha Block.

The table no. 1. provides the physico-chemical properties of soil (from 0 to 60 cm) in barren land of Akbarpur and Maitha block. The measurements are categorized into different depth ranges (0-15 cm, 15-30 cm, and 30-60 cm). The properties measured include pH, electrical conductivity (EC), organic carbon (OC), calcium (Ca), magnesium (Mg), and

sodium (Na) content. The table also includes statistics such as the minimum, maximum, mean, standard deviation (SD), and coefficient of variation (CV) for each property and depth range.

1.1. Akbarpur block

Soil pH is a measure of the acidity or alkalinity of the soil. It plays an important role in the availability of nutrients essential for plant growth. The pH value of the barren land is highly alkaline of Akbarpur block. The values of pH ranged from 8.52 to 8.90 with a mean value of 8.67 with an SD value of 0.16 and a CV value of 0.02 % at the surface of soil and all sites of soil sample decreasing to increasing level surface to ~~sub-surface~~subsurface of sites. Soil electrical conductivity is a measure of salinity of soil. It could be an indicator of nutrient availability in the soil (Sahu *et al.*, 2016). The low status of EC may be due to leaching of salts to lower horizons as suggested by Singh and Mishra (2012). The sample had very low soil organic carbon content on both blocks of sites. The values were 0.18 to 0.26 % in the surface soils and decreased to 0.10 to 0.21% at 30-60 cm depth. The low OC content of these soils may be attributed to ~~the~~occasional addition of organic materials, lack of natural vegetation, poor decomposition due to low rainfall, oxidation due to high summer temperature and wind erosion. ~~The~~A similar result was also reported by Prasad (2010). Exchangeable calcium and magnesium were extracted using neutral normal ammonium acetate and measured by EDTA titration. The exchangeable Ca content in Akbarpur block varied between 4.58 to 6.34 Cmol (+) kg⁻¹, whereas, the Mg content varied between 2.20 to 4.40 C mol (+) kg⁻¹ in surface of soil (0-15 cm). There was enough calcium and magnesium content in the soil to support plant growth. But Exchangeable sodium was a high quantity so harmful to plant growth.

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

1.2. Maitha

Physio-chemical properties of soil ~~are~~ given the data in ~~table~~Table no. 1. The pH value of the Maitha block is highly alkaline. The pH ranges between 8.74 to 9.62 and the mean value of pH is 9.46. The pH of 100 % soil samples ~~was~~ found more than 8.5 pH. The high degree of sodium saturation which on hydrolysis gives OH⁻ ion and high carbonate and bicarbonate. Similar results were also reported by Singh *et al.* (2010). Whereas the EC value of the soils was in neutral range i.e. 2.54 to 4.48 dS/m and the average value of EC was 3.65 dS/m and CV value is 0.20%. The EC of 100% soil sample under the rating of electrical conductivity. These values are fairly comparable to the result reported by Singh *et al.* (2010). The organic carbon content of the soils was very low due to ~~a~~deficit of vegetation. It ranged between 0.25 to 0.34%. The average, SD and CV, value of OC was 0.29, 0.04 and 0.13% respectively. The OC of 100 per cent sample were found lower than 0.50 per cent of OC. The low OC content of these soils may be attributed to occasional addition of organic materials, lack of natural vegetation, poor decomposition due to low rainfall, oxidation due to high summer temperature and wind erosion (Swarup, *at al.*, 2019). The similar result was also reported by Prasad (2010). Available Ca (cmol(p+) kg⁻¹) and Mg (cmol(p+) kg⁻¹) were medium to high in all sites of different Barren land, ranged from 3.86 to 5.64 and 3.68 to 6.12 in surface soils and from 4.88 to 7.46 and 5.78 to 7.35 in subsurface soils(30-60cm), respectively. Mean available Ca and Mg (cmol (p+) kg⁻¹) in surface soils were 4.90 and 4.99 in barren land, 4.1

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

and 1.7, respectively. With depth, there was increase in content under different sites of Maitha block.

1. Nutrient status in soil of Akbarpur block and Maitha Block.

1.1. Akbarpur

Akbarpur block of soil nutrient data show the table no. 2. Nitrogen content in soil samples ranged from 105.25 to 140.86 kg ha⁻¹ with a mean and CV value of 120.45 kg ha⁻¹ and 0.12% recorded in the Akbarpur block. The lowest nitrogen content was observed in the ~~sub surface~~ subsurface (30-60 cm) of soil while the highest nitrogen content was in the surface of soil but this nitrogen content compared to ~~the~~ soil fertility index was overall very less amount content. The nitrogen content of the study region is low probably due to ~~the~~ low organic carbon content present in the soil (Mandal, et al., 2022). ~~Available~~ ~~The available~~ phosphorus content of soil samples ranged from 12.36 to 18.94 kg ha⁻¹ with an average, SD and CV value of 16.00 kg ha⁻¹, 2.45 and 0.15%, respectively. The lowest value of phosphorus content was observed in the subsurface soil, whereas the highest value was in ~~the~~ surface of soil. About 50% of phosphorus is found in organic form and decomposition of the organic matter produces humus which forms complexes with Al and Fe and ~~protect~~ ~~protects~~ the P fixation (Kirkby, et al., 2011). Available potassium content of soil ranged from 233.12 to 278.85 kg ha⁻¹ with a mean value of 256 kg ha⁻¹ in the surface of soil. In the study region potassium content is high, may be due to elite, rich potassium minerals found in the soil. The data on ~~micronutrient~~ ~~micronutrients~~ were recorded in which the nutrient index value indicates the Zn, Mn, Cu and B were deficient while the Fe content was sufficient. Overall, the Zn content of the studied block ranges from 0.44 to 0.68 mg kg⁻¹ with mean value 0.57 mg kg⁻¹ while the Mn ranges from 4.64 to 6.25 mg kg⁻¹ with mean value of 5.45 mg kg⁻¹. The Fe content ranges from 11.89 to 14.25 mg kg⁻¹ with mean value 13.03 mg kg⁻¹, the Cu content of the soil samples on the whole ranges from 1.4 to 1.89 mg kg⁻¹ with mean value 1.29 mg kg⁻¹. The similar work reported by Gautam, et al. (2023)

2.2. Maitha

The results of the soil nutrient status (Table 3) revealed that low to medium content of available N, P and K were found in all sites, and it varied from 102.78 to 138.39, 9.89 to 16.47 kg ha⁻¹ and 230.65 to 276.65 kg ha⁻¹ in surface soils; 22.89 to 31.20, 3.16 to 4.68 and 81.89 to 107.98 kg ha⁻¹ in subsurface soils under all sites of Maitha block. The content of available N, P and K was higher in the surface horizons as compared to subsurface horizons. A regular decreasing pattern of available N, P and K was observed with depth. The highest amount of available N in the surface horizon may be attributed to the high amount of carbon. The lower P in sub-surface horizons as compared to surface horizon was due to the fixation of released P by clay minerals, oxides of iron and aluminum. Available S (kg ha⁻¹) content was also low to medium in most soils of different sites of barren land and varied from 10.78 to 14.07 kg ha⁻¹ in surface soils and from 7.56 to 9.31 kg ha⁻¹ in subsurface soils, respectively. Mean available S (kg ha⁻¹) in surface soils was 12.33 in all sites of Maitha block. However, the highest available S was observed in the surface horizons and decreased with depth. The DTPA-Cu, Fe and Zn were higher in surface horizon than sub-surface under all sites of barren land and it ranged from 1.00 to 1.85, 9.42 to 11.78 and 0.40 to 0.64 (mg kg⁻¹

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

¹) in surface soils, from 0.32 to 0.61, 2.09 to 3.67 and 0.09 to 0.20 (mg kg⁻¹) in subsurface soils, respectively. Mean available Cu, Fe and Zn (mg kg⁻¹) in surface soils were 1.25, 10.56 and 0.053, respectively in the all sites of barren land. The DTPA- Mn varied from 6.08 to 7.69 with ~~the a~~mean value was 6.89 (mg kg⁻¹) in surface soils; from 3.88 to 5.69 with mean and SD ~~value-values~~were 4.84(mg kg⁻¹) and 0.75 in subsurface soils, respectively under all sites. Irrespective of sample sites, the Mn showed an increased to decreasing trend with depth.

Formatted: English (United States)

Formatted: English (United States)

The candidate manuscript does not have a robust scientific discussion, I suggest the authors incorporate the suggested paragraphs, in this way, it would improve the scientific quality of the manuscript:

Formatted: Highlight

Formatted: Highlight

Assessing the major and micronutrient status of soils in the barren land of the central plain of Uttar Pradesh is crucial to understanding the soil's fertility potential. It helps identify nutrient deficiencies or imbalances that may limit plant growth and productivity (Calero et al. 2022). By analyzing nutrient levels (Olivares et al. 2022a), researchers and farmers can determine the appropriate fertilization practices needed to optimize crop yields and sustainably manage the land (Olivares et al. 2022b)

Understanding the nutrient status of soils is essential for effective nutrient management (Olivares et al. 2022c). By comparing the results of the assessment with quality studies of tropical soils in Latin America, researchers can gain insights into common nutrient patterns or deficiencies (Araya-Alman et al. 2020). This information can guide the formulation of region-specific nutrient management strategies (Lobo et al. 2023), including the selection and application of fertilizers or organic amendments tailored to the specific nutrient requirements of crops.

Crop Productivity Improvement: Nutrient deficiencies can significantly impact crop productivity (Hernandez et al. 2018b). By identifying the major and micronutrient deficiencies in the soils of the barren land in the central plain of Uttar Pradesh and comparing them with tropical soils in Latin America (Olivares et al. 2011), researchers can draw parallels and learn from successful soil management practices employed in similar contexts (Olivares, 2016; Hernandez et al. 2017). This knowledge can help develop effective interventions to improve crop productivity by addressing nutrient deficiencies through targeted fertilization or soil amendment approaches (Hernandez et al. 2018a).

Sustainable Agriculture: Sustainable agriculture aims to optimize productivity while minimizing the environmental impact. Assessing nutrient status in soils and comparing it with studies from Latin America can provide valuable insights into sustainable soil management practices (Cortez et al. 2018; Hernandez and Olivares, 2020). By understanding the nutrient dynamics and interactions in these regions (Hernandez et al. 2018a; Zingaretti et al. 2016; Zingaretti and Olivares, 2018, researchers can recommend appropriate soil conservation and nutrient management practices that promote long-term soil health (Lopez-

Beltran et al. 2019; Lopez and Olivares, 2019), reduce nutrient runoff, (Olivares, 2022) and minimize environmental degradation (Parra et al. 2012; Cortez et al. 2018; Hernandez et al. 2020).

Knowledge Exchange and Collaboration: Comparing the results of soil assessments between different regions fosters knowledge exchange and collaboration among scientists, researchers, and agricultural practitioners. It allows for the identification of best practices and lessons learned from different contexts, leading to innovations in soil management and agricultural productivity improvement (Paredes et al. 2021; Pitti et al. 2021).

In conclusion, the assessment of available major and micronutrient status of soils in the barren land of the central plain of Uttar Pradesh and its comparison with quality studies of tropical soils in Latin America can provide valuable insights for understanding soil fertility, nutrient management, and strategies to enhance agricultural productivity sustainably. These findings can guide soil management practices, nutrient supplementation, and knowledge sharing, ultimately contributing to improved agricultural outcomes and sustainable land use.

conclusion

The present study reveals that the soils are highly deficient in nutrients and suffer from poor soil fertility status. One of the main reasons for declining productivity is the lack of knowledge, awareness among the farmers and also non-adoption of sustainable crop management strategies including comprehensive soil health management. Nowadays there is a growing need for balanced fertilization and ~~site-specific~~ site-specific fertilizer recommendations according to the crop type, yield level and soil conditions. With the obligatory need for intensification of crop production, the demand of crops for readily available soil nutrients increases. Strengthening and finding out the right sources of amendments for improving fertility of problem soils, which will also supply more than one nutrient to economize the crop production, may be adopted as a means to improve problem soil and make wealth out of poor fertile lands.

References

Allen, D.E., Singh, B.P. and Dalal, R.C., 2011. Soil health indicators under climate change: a review of current knowledge. *Soil health and climate change*, pp.25-45.

Araya-Alman, M., Olivares, B., Acevedo-Opazo, C. et al. 2020. Relationship Between Soil Properties and Banana Productivity in the Two Main Cultivation Areas in Venezuela. *J Soil Sci Plant Nutr.* 20 (3): 2512-2524. <https://doi.org/10.1007/s42729-020-00317-8>

Babur, E., Kara, O., Fathi, R.A., Susam, Y.E., Riaz, M., Arif, M. and Akhtar, K., 2021. Wattle fencing improved soil aggregate stability, organic carbon stocks and biochemical

Formatted: English (United States)

Formatted: English (United States)

Formatted: Font: Italic

Formatted: English (United States)

quality by restoring highly eroded mountain region soil. *Journal of Environmental Management*, 288, p.112489.

Bhattacharyya, R., Ghosh, B.N., Mishra, P.K., Mandal, B., Rao, C.S., Sarkar, D., Das, K., Anil, K.S., Lalitha, M., Hati, K.M. and Franzluebbbers, A.J., 2015. Soil degradation in India: Challenges and potential solutions. *Sustainability*, 7(4), pp.3528-3570.

Carter, M.R., 2002. Soil quality for sustainable land management: organic matter and aggregation interactions that maintain soil functions. *Agronomy journal*, 94(1), pp.38-47.

[Calero, J., Olivares, B.O., Rey, J.C., Lobo, D., Landa, B.B., Gómez, J. A. 2022. Correlation of banana productivity levels and soil morphological properties using regularized optimal scaling regression. *Catena*, 208: 105718. https://doi.org/10.1016/j.catena.2021.105718](https://doi.org/10.1016/j.catena.2021.105718)

Formatted: Font: Italic

[Cortez, A., Olivares, B., Parra, R., Lobo, D., Rodríguez, M.F., Rey, J.C. 2018. Descripción de los eventos de sequía meteorológica en localidades de la cordillera central, Venezuela. *Ciencia, Ingenierías y Aplicaciones*. I \(1\):22-44. DOI: http://dx.doi.org/10.22206/cyap.2018.v1i1.pp23-45](http://dx.doi.org/10.22206/cyap.2018.v1i1.pp23-45)

Formatted: Font: Italic

Formatted: English (United States)

Dhaliwal, S.S., Naresh, R.K., Mandal, A., Singh, R. and Dhaliwal, M.K., 2019. Dynamics and transformations of micronutrients in agricultural soils as influenced by organic matter build-up: A review. *Environmental and Sustainability Indicators*, 1, p.100007.

Fageria, N.K., 2012. Role of soil organic matter in maintaining sustainability of cropping systems. *Communications in soil science and plant analysis*, 43(16), pp.2063-2113.

Gautam, M.K., Ghosh, A.K., Kumar, S., Kamal, M.K., Lal, K. and Gautam, A.K., 2023. Characterization and Study of Dominated Alluvial Soil Profile of Mirzapur in Indo-Gangetic Plain (IGPs) of India. *International Journal of Plant & Soil Science*, 35(9), pp.160-177.

Harris, J.A., 2003. Measurements of the soil microbial community for estimating the success of restoration. *European Journal of Soil Science*, 54(4), pp.801-808.

[Hernández, R. Olivares, B. 2020. Application of multivariate techniques in the agricultural land's aptitude in Carabobo, Venezuela. *Tropical and Subtropical Agroecosystems*, 23\(2\):1-12. https://n9.cl/zeedh](https://n9.cl/zeedh)

[Hernández, R; Pereira, Y; Molina, JC; Coelho, R; Olivares, B., Rodríguez, K. 2017. Calendario de siembra para las zonas agrícolas del estado Carabobo en la República Bolivariana de Venezuela. Sevilla, Spain, Editorial Universidad Internacional de Andalucía. 247 p.](#)

[Hernandez, R.; Olivares, B.; Arias, A; Molina, JC., Pereira, Y. 2020. Eco-territorial adaptability of tomato crops for sustainable agricultural production in Carabobo, Venezuela. *Idesia*, 38\(2\):95-102. http://dx.doi.org/10.4067/S0718-34292020000200095](http://dx.doi.org/10.4067/S0718-34292020000200095)

[Hernández, R; Olivares, B. Arias, A; Molina, JC., Pereira, Y. 2018a. Zonificación agroclimática del cultivo de maíz para la sostenibilidad de la producción agrícola en Carabobo, Venezuela. *Revista Universitaria de Geografía*. 27 \(2\): 139-159. https://n9.cl/12m83](https://n9.cl/12m83)

Hernández, R; Olivares, B., Arias, A; Molina, J.C., Pereira, Y. 2018b. Identificación de zonas agroclimáticas potenciales para producción de cebolla (*Allium cepa* L.) en Carabobo, Venezuela. Journal of the Selva Andina Biosphere. 6 (2): 70-82. Disponible en http://www.scielo.org.bo/pdf/jsab/v6n2/v6n2_a03.pdf

Formatted: Font: Italic

Hernández, R; Olivares, B., Coelho, R., Molina, J.C., Pereira, Y. 2018c. Análisis espacial del índice hídrico: un avance en la adopción de decisiones sostenibles en los territorios agrícolas de Carabobo, Venezuela. Revista Geográfica de América Central. 60 (1): 277-299. DOI: <https://doi.org/10.15359/rgac.60-1.10>

Formatted: English (United States)

Kirkby, C.A., Kirkegaard, J.A., Richardson, A.E., Wade, L.J., Blanchard, C. and Batten, G., 2011. Stable soil organic matter: a comparison of C: N: P: S ratios in Australian and other world soils. *Geoderma*, 163(3-4), pp.197-208.

Köppen W & Geiger R (1936) Das Geographische System der Klimate. In: Handbuch der Klimatologie (eds W Köppen & R Geiger), Bd 1, Teil C. Verlag Gebrüder Bornträger, Berlin (DE) (in German).

Lobo, D; Olivares, B; Rey, J.C; Vega, A; Rueda-Calderón, A. (2023). Relationships between the Visual Evaluation of Soil Structure (VESS) and soil properties in agriculture: A meta-analysis. Scientia agropecuaria, 14 - 1, 67 - 78. <https://doi.org/10.17268/sci.agropecu.2023.007>

López-Beltrán, M., Olivares, B., Lobo-Luján, D. (2019). Changes in land use and vegetation in the agrarian community Kashaama, Anzoátegui, Venezuela: 2001-2013. Revista Geográfica De América Central. 2(63):269-291. DOI: <https://doi.org/10.15359/rgac.63-2.10>

López, M. Olivares, B. (2019). Normalized Difference Vegetation Index (NDVI) applied to the agricultural indigenous territory of Kashaama, Venezuela. UNED Research Journal. 11(2): 112-121. <https://doi.org/10.22458/urj.v11i2.2299>

Formatted: English (United States)

Mandal, A., Toor, A.S., Dhaliwal, S.S., Singh, P., Singh, V.K., Sharma, V., Gupta, R.K., Naresh, R.K., Kumar, Y., Pramanick, B. and Nanda, G., 2022. Long-Term Field and Horticultural Crops Intensification in Semiarid Regions Influence the Soil Physiobiochemical Properties and Nutrients Status. *Agronomy*, 12(5), p.1010.

Meena, M.D., Yadav, R.K., Narjary, B., Yadav, G., Jat, H.S., Sheoran, P., Meena, M.K., Antil, R.S., Meena, B.L., Singh, H.V. and Meena, V.S., 2019. Municipal solid waste (MSW): Strategies to improve salt affected soil sustainability: A review. *Waste management*, 84, pp.38-53.

Mensah, A.K., 2015. Role of revegetation in restoring fertility of degraded mined soils in Ghana: A review. *International journal of biodiversity and conservation*, 7(2), pp.57-80.

Olivares, B.O.; Rey, J.C.; Perichi, G.; Lobo, D. 2022a. Relationship of Microbial Activity with Soil Properties in Banana Plantations in Venezuela. Sustainability 14, 13531. <https://doi.org/10.3390/su142013531>

Olivares B, Vega A, Calderón MAR, Rey JC, Lobo D, Gómez JA, Landa BB. 2022b. Identification of Soil Properties Associated with the Incidence of Banana Wilt Using Supervised Methods. *Plants*, 11(15):2070. <https://doi.org/10.3390/plants11152070>

Olivares B, Rey JC, Lobo D, Navas-Cortés JA, Gómez JA, Landa BB. 2022c. Machine Learning and the New Sustainable Agriculture: Applications in Banana Production Systems of Venezuela. *Agricultural Research Updates*. 42, pp. 133 - 157. Nova Science Publishers, Inc.

Olivares, B. O. 2022. Determination of the potential influence of soil in the differentiation of productivity and in the classification of susceptible areas to banana wilt in Venezuela. Doctoral dissertation, Universidad de Córdoba, UCOPress, Spain.

Olivares, B. 2016. Descripción del manejo de suelos en sistemas de producción agrícola del sector Hamaca de Anzoátegui, Venezuela. *La Granja: Revista de Ciencias de la Vida*. 23(1): 14–24. <https://n9.cl/ycp08>

Olivares, B., Verbist, K., Lobo, D., Vargas, R. y Silva, O. 2011. Evaluation of the USLE model to estimate water erosion in an Alfisol. *Journal of Soil Science and Plant Nutrition of Chile*. 11 (2):71-84. <http://dx.doi.org/10.4067/S0718-95162011000200007>

Paredes, F., Olivares, B., Rey, J., Lobo, D., Galvis-Causil, S. 2021. The relationship between the normalized difference vegetation index, rainfall, and potential evapotranspiration in a banana plantation of Venezuela. *SAINS TANAH - Journal of Soil Science and Agroclimatology*, 18(1), 58-64. <http://dx.doi.org/10.20961/stjssa.v18i1.50379>

Parra, R., Olivares, B., Cortez, A. y Rodríguez, M.F. 2012. Patrones de homogeneidad pluviométrica en estaciones climáticas del estado Anzoátegui, Venezuela. *Revista Multiciencias*. 12 (Extraordinario): 11-17. <https://n9.cl/xbslq>

Pitti, J; Olivares, B; Montenegro, E. 2021. The role of agriculture in the Changuinola District: a case of applied economics in Panama. *Tropical and Subtropical Agroecosystems*. 25 - 1, 1 - 11. <http://dx.doi.org/10.56369/tsaes.3815>

Ruiz-Colmenero, M., Bienes, R., Eldridge, D.J. and Marques, M.J., 2013. Vegetation cover reduces erosion and enhances soil organic carbon in a vineyard in the central Spain. *Catena*, 104, pp.153-160.

Sharma, D.K. and Singh, A., 2015. Salinity research in India-achievements, challenges and future prospects.

Swarup, A., Manna, M.C. and Singh, G.B., 2019. Impact of land use and management practices on organic carbon dynamics in soils of India. *Global climate change and tropical ecosystems*, pp.261-281.

Udom, B.E. and Ogunwole, J.O., 2015. Soil organic carbon, nitrogen, and phosphorus distribution in stable aggregates of an Ultisol under contrasting land use and management history. *Journal of Plant Nutrition and Soil Science*, 178(3), pp.460-467.

Zingaretti, M.L., Olivares, B. 2018. Analysis of the meteorological drought in four agricultural locations of Venezuela by the combination of multivariate methods. UNED Research Journal. 10 (1):181-192. <http://dx.doi.org/10.22458/urj.v10i1.2026>

Zingaretti, M.L., Olivares, B., Demey Zambrano, J.A., Demey, J.R. 2016. Tipificación de los sistemas de producción agrícola y la percepción de la variabilidad climática en Anzoátegui, Venezuela. Revista FAVE - Ciencias Agrarias 15 (2): 39-50. <https://doi.org/10.14409/fa.v15i2.6587>

Formatted: English (United States)

UNDER PEER REVIEW

Table.2. Nutrient content of soil (from 0 to 60 cm) in barren land of Akbarpur.

	Depth	Macro Nutrients (kgha ⁻¹)				Micro Nutrients (Mgha ⁻¹)			
		N	P	K	S	Cu	Fe	Mn	Zn
Min.	0-15	105.25	12.36	233.12	13.25	1.04	11.89	4.64	0.44
	15-30	45.86	10.25	210.45	12.77	1.02	9.19	5.63	0.23
	30-60	25.36	5.63	84.36	10.12	0.36	4.56	2.44	0.13
Max.	0-15	140.86	18.94	278.85	16.54	1.89	14.25	6.25	0.68
	15-30	67.15	13.45	288.45	15.60	2.11	11.80	6.63	0.46
	30-60	33.67	7.15	110.45	11.78	0.65	6.14	4.25	0.24
Mean	0-15	120.45	16.00	259.86	14.80	1.29	13.03	5.45	0.57
	15-30	58.57	11.26	239.38	14.32	1.45	10.44	5.89	0.34
	30-60	30.44	6.35	94.18	10.95	0.50	5.54	3.40	0.20
SD	0-15	13.88	2.45	17.67	1.26	0.35	0.88	0.75	0.09
	15-30	8.17	1.34	29.16	1.15	0.41	1.01	0.43	0.09
	30-60	3.76	0.76	10.44	0.73	0.13	0.62	0.66	0.04
CV	0-15	0.12	0.15	0.07	0.09	0.27	0.07	0.14	0.16
	15-30	0.14	0.12	0.12	0.08	0.28	0.10	0.07	0.26
	30-60	0.12	0.12	0.11	0.07	0.26	0.11	0.19	0.21

Formatted: English (United States)

Formatted: English (United States)

Formatted: Font: Not Bold, English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Table.3. Nutrient content of soil (from 0 to 60 cm) in barren land of Maitha during the Rabi Session.

	Depth	Macro Nutrients (kgha ⁻¹)				Micro Nutrients (Mgha ⁻¹)			
		N	P	K	S	Cu	Fe	Mn	Zn
Min.	0-15	102.78	9.89	230.65	10.78	1.00	9.42	6.08	0.40
	15-30	43.39	7.78	207.98	10.30	0.98	6.72	7.07	0.19
	30-60	22.89	3.16	81.89	7.65	0.32	2.09	3.88	0.09
Max.	0-15	138.39	16.47	276.38	14.07	1.85	11.78	7.69	0.64
	15-30	64.68	10.98	285.98	13.13	2.07	9.33	8.07	0.42
	30-60	31.20	4.68	107.98	9.31	0.61	3.67	5.69	0.20
Mean	0-15	117.98	13.53	257.39	12.33	1.25	10.56	6.89	0.53
	15-30	56.10	8.79	236.91	11.85	1.41	7.97	7.33	0.30
	30-60	27.97	3.88	91.71	8.48	0.46	3.07	4.84	0.16
SD	0-15	13.88	2.45	17.67	1.26	0.35	0.88	0.75	0.09
	15-30	8.17	1.34	29.16	1.15	0.41	1.01	0.43	0.09
	30-60	3.76	0.76	10.44	0.73	0.13	0.62	0.66	0.04
CV	0-15	0.12	0.18	0.07	0.10	0.28	0.08	0.11	0.18
	15-30	0.15	0.15	0.12	0.10	0.29	0.13	0.06	0.30
	30-60	0.13	0.20	0.11	0.09	0.28	0.20	0.14	0.26

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: Font: Not Bold, English (United States)

Formatted: English (United States)

|

UNDER PEER REVIEW

