

# Diversity of Chrysanthemum (*Dendranthema grandiflora* T.) varieties under open field condition in Prayagraj

## ABSTRACT

The field experiment was carried out during September 2022 to January 2023, in Horticulture Research Farm, Department of Horticulture, Naini Agriculture Institute, SHUATS. The experiment was laid out in Randomized Block Design with three replications. The experiment comprised of fifteen varieties of chrysanthemum viz. Kanadee, Flood, Winter Queen, Basanti, White Bonsai, Local Yellow, Ravi Kiran, Button Type, Rani, UBC 12, Wall Street, Bidhan Rajat, Bidhan Antara, British gold and Vijay. It is clear from the experimental analysis that all characters were significantly affected by different varietal treatments. From the experimental findings, it was found that maximum height was found in the variety Bidhan Antara (40.79 cm), plant spread (30.55 cm), primary branches (6), whereas earliness in flowering was found in the variety Bidhan Rajat, flower yield per plant (176.76 g) was found maximum in the variety Bidhan Rajat, average flower weight was recorded more in the variety Rani (7.19 g), duration of flowering (71.2 days) and benefit-cost ratio (4.32:1) was found maximum in the variety Bidhan Rajat.

## INTRODUCTION

Chrysanthemum (*Dendranthema grandiflora* T.) belongs to the Asteraceae family. It is believed to be native to the northern hemisphere chiefly Europe and Asia and was believed to have been originated in China. The basic chromosome number of chrysanthemum is 9, while  $2n$  ranges from 36 to 75 though most of them are hexaploid. It is the national flower of Japan. It is commonly called as “Queen of the East”, “Autumn Queen”, “Guldaudi” in India and “Mum” in America. The word chrysanthemum is derived from the Greek word “chryos” means gold and “anthemon” or “anthos” means flower.

In the trade of global flower market, chrysanthemum is the second largest cut flower after rose and holds fifth rank as pot plant. It is commercially grown in different parts of the world. Netherlands, Italy, Colombia, Spain, Germany and USA are the important countries where it is mainly grown under greenhouse conditions. Japan is the largest producer of chrysanthemum in the world. In India, it is commercially grown in Karnataka, Tamil Nadu and Maharashtra. Chrysanthemum covers 20,090 ha area with production of 1,85,240 MT of loose flowers and 14,930 MT of cut flowers in India during 2016-2017. Karnataka is the most prominent chrysanthemum growing state with an area of 5100 ha and production of 61,200 MT of loose flowers during 2014-2015. In different states of India, it is grown with different names, Guldaudi in Hindi belt, Chandramalika in the eastern states, Samanti in the southern

states and Shevanti in the western states of India. Ease of cultivation, high returns and increasing market demand are the main reasons for the popularity of this crop.

In India, chrysanthemum occupies a place of pride both as a commercial crop and as a popular exhibition flower. The erect and tall growing cultivars are suitable for background planting in borders. The cultivars with the dwarf and compact growing habit, on the other hand, are suitable for front row plantation or pot culture. The decorative and fluffy bloomed smallflowered cultivars are ideal for garland making and hair decoration. The extra-large bloomed cultivars are used for exhibition value. Loose flowers are used for garlands, venis, worship etc. Long stem flowers or cut flowers are used for bouquet, vase etc. In North India various hues of red, yellow, white and purple chrysanthemums are grown in abundance for decorating the landscape either in the ground or in pots. But, in South India mostly the yellow coloured flowers are preferred and grown as loose flowers for trade. The cultivation of chrysanthemum is gaining importance in Gujarat due to its relative ease in cultivation, high returns and increasing market demand.

## **METHODS AND MATERIALS**

The field experiment was carried out at Horticulture Research Field, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences (SHUATS), Allahabad, during September 2022 to January 2023.

### **Geographical location of the experimental site:**

The experimental site is being located at a latitude of 25.41° North and longitude of 81.84 ° East, with an altitude of 98 meters above the mean sea level (MSL).

### **Climatic conditions of the experimental area:**

The area of Prayagraj comes under humid sub-tropical climate, which experiences warm humid monsoon, hot dry summer and cold dry winter. The annual mean temperature is 26.1°C while monthly mean temperatures are 18-29°C. The daily average maximum temperature is about 22°C and the minimum temperature is 9°C. The average annual rainfall received is 1042.2 mm. At this location, the temperature reaches upto 46°C-48°C and the minimum temperature recorded was 4°C-5°C. The relative humidity ranges in this location ranges between 20-94%.

## **RESULT AND DISCUSSION**

The present investigation was carried out for “Varietal evaluation of chrysanthemum (*Dendranthema grandiflora* L.) Under open field condition in Prayagraj” was made to find out best performing variety of Chrysanthemum in terms of flowering and yield, and to estimate the economics of different varieties. For this purpose, 15 cultivar were laid out in Randomized block Design (RBD) with three replications. The Varieties are Kanadee, Flood, Winter Queen, Basanti, White Bonsai, Local Yellow, Ravi Kiran, Button Type, Rani, UBC 12, Wall Street, Bidhan Rajat, Bidhan Antara, British gold and Vijay. These varieties of chrysanthemum were planted during rabi 2022-23. In present experiment data were recorded for various characters, viz., (A) Growth parameter: Plant height (cm) 30, 60, 90 DAP; Plant

spread(cm)30,60,90 DAP; Number of primary branches, (B) Floral parameter: . Days taken to first flower bud initiation, Days taken to 50% flowering, Number of flowers per plant, Flower form, Flower color, Flower diameter (cm), Disc color 8. Stalk length (cm), Flower yield per plant (g), Flower yield per hectare (q), Flower weight (g) , Duration of flowering (days); (D) Economic parameters: Cost of Cultivation, Gross Return, Net Return , Benefit Cost Ratio. The results of the present work are presented under following headings.

**TABLE:1- Mean Performance of Plant height**

TREATMENT	VARIETY	30DAP	60DAP	90DAP
V1	KANADEE	7.14	12.18	23.567
V2	FLOOD	9.067	15.807	29.953
V3	WINTER QUEEN	7.187	13.667	31.627
V4	BASANTI	7.647	15.467	28.307
V5	WHITE BONSAI	3.753	6.72	11.227
V6	LOCAL YELLOW	10.26	16.34	40.793
V7	RAVI KIRAN	6.287	11.053	23.12
V8	BUTTON TYPE	3.453	11.593	18.1
V9	RANI	8.393	16.78	36.953
V10	UBC 12	4.207	11.44	16.193
V11	WALL STREET	3.213	10.547	12.537
V12	BIDHAN RAJAT	11.273	19.953	28.653
V13	BIDHAN ANTARA	13.073	20.727	39.207
V14	BRITISH GOLD	5.667	10.853	22.493
V15	VIJAY	4.093	6.42	9.367
	C.D.	2.007	4.124	3.119
	SE(m)	0.689	1.416	1.071
	SE(d)	0.975	2.003	1.515
	C.V.	17.102	18.438	7.478

**TABLE:2- Mean Performance of Plant spread**

TREATMENT	VARIETY	30DAP	60DAP	90DAP
V1	KANADEE	4.433	8.067	16.033
V2	FLOOD	3.547	8.887	12.513

V3	WINTER QUEEN	3.66	7.68	11.927
V4	BASANTI	5.087	10.827	30.553
V5	WHITE BONSAI	3.827	8.507	13.627
V6	LOCAL YELLOW	4.453	9.8	22.293
V7	RAVI KIRAN	3.1	9.433	15.34
V8	BUTTON TYPE	2.487	7.333	14.293
V9	RANI	3.48	9.4	14.387
V10	UBC 12	2.38	8.78	14.26
V11	WALL STREET	2.081	11.693	16.907
V12	BIDHAN RAJAT	3.067	10.787	21.5
V13	BIDHAN ANTARA	5.747	12.52	24.307
V14	BRITISH GOLD	3.193	8.313	15.927
V15	VIJAY	2.147	6.673	9.513
	C.D.	0.673	1.91	4.308
	SE(m)	0.231	0.656	1.479
	SE(d)	0.327	0.927	2.092
	C.V.	11.398	12.284	15.169

**TABLE:3- Mean Performance of different varieties of chrysanthemum**

S.No	VARIETIES	NO OF PRIMARY BRANCHES	NO OF DAYS TO FIRST BUD INITIATION	DAYS TO 50%FLOWERING	FLOWER PER PLANT	FLOWER DIAMETER
1	KANADEE	4.133	80	115.8	19.133	5.693
2	FLOOD	4.4	76.6	107	23.667	<b>8.633</b>
3	WINTER QUEEN	3.333	87.2	110.8	14.6	8.12
4	BASANTI	5.133	75.467	102.867	37.867	4.627
5	WHITE BONSAI	5.867	70.4	106.333	39.867	4.193
6	LOCAL YELLOW	3.8	78.4	97.667	28	7.473
7	RAVI KIRAN	4.333	81.133	126.467	21.6	6.467
8	BUTTON TYPE	6	84.867	108.133	26.4	4.773
9	RANI	5.067	77.333	106.8	20.2	7.107

10	UBC 12	4.533	85.4	106.667	38.4	2.247
11	WALL STREET	3.667	67.067	84.467	26.467	4.467
12	BIDHAN RAJAT	10	71.467	102.2	25	8.093
13	BIDHAN ANTARA	10	72.733	96.933	41.467	4.74
14	BRITISH GOLD	8.133	88.267	124.8	22.4	5.067
15	VIJAY	5.467	77.533	115.733	27.8	3.833
	MEAN	5.591	78.2578	107.5111	27.5245	5.7022
	C.D.	0.795	5.145	5.433	1.701	0.862
	SE(m)	0.273	1.767	1.866	0.584	0.296
	SE(d)	0.386	2.499	2.638	0.826	0.419
	C.V.	8.453	3.911	3.006	3.676	8.993

**TABLE:4- Mean Performance of different varieties of chrysanthemum**

S.No	VARIETIES	STALK LENGTH	FLOWER WEIGHT	YEILD PER PLANT	YEILD PER HECTARE	FLOWERING DURATION
1	KANADEE	5.013	5.22	98.22	88.398	62.667
2	FLOOD	11.447	6.193	144.007	125.339	68.733
3	WINTER QUEEN	7.213	1.34	19.627	17.664	46.533
4	BASANTI	5.493	3.153	119.453	115.651	71.067
5	WHITE BONSAI	4.64	1.107	44.207	39.786	48.4
6	LOCAL YELLOW	10.62	4.213	117.993	106.194	68.267
7	RAVI KIRAN	5.093	4.153	89.72	80.748	48.333
8	BUTTON TYPE	5.133	1.14	29.913	26.922	42.4
9	RANI	7.06	7.193	148.353	130.818	52.667
10	UBC 12	5.113	0.64	24.407	21.966	52.933
11	WALL STREET	5.84	2.073	55.093	49.584	48.867
12	BIDHAN RAJAT	8.833	7.073	176.76	159.084	61.2

<b>13</b>	<b>BIDHAN ANTARA</b>	5.487	2.453	101.767	91.601	63.267
<b>14</b>	<b>BRITISH GOLD</b>	7.167	2.433	54.393	48.954	56
<b>15</b>	<b>VIJAY</b>	3.42	1.053	29.313	26.388	41.8
	<b>MEAN</b>	6.5048	3.2958	83.5484	75.2731	55.5422
	<b>C.D.</b>	0.429	0.073	7.863	10.404	2.899
	<b>SE(m)</b>	0.147	0.025	2.701	3.573	0.996
	<b>SE(d)</b>	0.208	0.035	3.819	5.053	1.408
	<b>C.V.</b>	3.923	1.312	5.598	8.221	3.105

**TABLE:5- GROSS RETURN,NET RETURN, BENEFIT COST RATIO**

<b>S.No</b>	<b>VARIETY</b>	<b>Yeild(q/ha)</b>	<b>Selling price/q</b>	<b>Gross return (Rs./ha)</b>	<b>Cost of cultivation (Rs./ha)</b>	<b>Net return (Rs./ha)</b>	<b>Benefit cost ratio</b>
<b>1</b>	<b>KANADEE</b>	88.398	18000	1591164	1037450	553714	0.533
<b>2</b>	<b>FLOOD</b>	125.339	18000	2256102	1037450	1218652	1.174
<b>3</b>	<b>WINTER QUEEN</b>	17.664	40000	706560	537450	169110	0.314
<b>4</b>	<b>BASANTI</b>	115.651	18000	2081718	537450	1544268	2.873
<b>5</b>	<b>WHITE BONSAI</b>	39.786	35000	1392510	1037450	355060	0.342
<b>6</b>	<b>LOCAL YELLOW</b>	106.194	18000	1911492	537450	137450	0.255
<b>7</b>	<b>RAVI KIRAN</b>	80.748	18000	1453464	1037450	416014	0.4
<b>8</b>	<b>BUTTON TYPE</b>	26.922	35000	942270	537450	404820	0.753
<b>9</b>	<b>RANI</b>	130.818	18000	2354724	1037450	1317274	1.269
<b>10</b>	<b>UBC 12</b>	21.966	40000	878640	537450	341190	0.634
<b>11</b>	<b>WALL STREET</b>	49.584	35000	1735440	1037450	697990	0.672
<b>12</b>	<b>BIDHAN RAJAT</b>	159.084	18000	2863512	537450	2326062	4.327
<b>13</b>	<b>BIDHAN ANTARA</b>	91.601	18000	1648818	537450	1111368	2.067
<b>14</b>	<b>BRITISH GOLD</b>	48.954	35000	1713390	1037450	675940	0.651
<b>15</b>	<b>VIJAY</b>	26.388	40000	1055520	1037450	18070	0.017

## CONCLUSION

It is concluded from the present investigation that the 15 chrysanthemum varieties showed significant variation in all the parameters observed. The variety Bidhan Rajat(V-12) showed the best performance in most of the parameters like number of primary branches, minimum days taken for first flower bud initiation, duration of flowering, flower yield per plant, gross returns, net returns and benefit-cost ratio, at par with the variety Bidhan Antara, followed by Basanti. While, the variety winter queen didn't perform well and stands at last place in each parameter.

## REFERENCE

- Bala, M. (2015). Evaluation of chrysanthemum (*Chrysanthemum morifolium* Ramat.) genotypes for morphological traits. *Journal of Horticultural Science*, 10(2):242- 244
- Cantor, M., Hitter, T., Varga, Z.S. and Buta, E. (2020). Studies Regarding a Varietal Assortment of Potted Chrysanthemum. *Romanian Journal of Horticulture*, 1:159- 166.
- Chawla, S.L., Patel, R.B., Dhaduk, B.K., Sudha, P. and Dipal, B. (2021). Varietal assessment of Chrysanthemum (*Dendranthema grandiflora*) under South Gujarat agro-climatic conditions. *Current Horticulture*, 9(2): 64-67.
- Fatmi, U., Deepanshu, Singh, D. and Shohe, M. (2017). Evaluation of Chrysanthemum (*Dendranthema grandiflora* Tzvelev) Cultivars under Allahabad agro-climatic conditions. *Trends in Biosciences*, 10(24):5115-5119.
- Guddaraddi, A. and Mishra, A. (2021). Varietal evaluation and genetic variability of chrysanthemum (*Dendranthema grandiflora* Tzvelev). *The Pharma Innovation Journal*, 10(12): 563-566.
- Kireeti, A., Ravindrababu, M., Prasad, J. and Ramadevi, P. (2017). Evaluation of chrysanthemum (*Dendranthema grandiflora* Tzvelev) varieties in humid coastal zone of Andhra Pradesh. *International Journal of Chemical Studies* 5(3): 370- 372.
- Kumar, A.S. and Polara, N. D. (2017). Evaluation of Chrysanthemum Varieties on Growth and Quality under South Saurashtra Region. *International Journal of Pure Applied Biosciences*, 5 (4): 1989-1997.
- Kumar,r.,Prasad,V.M.,Singh,D.(2021). Varietal evaluation of chrysanthemum (*Dendranthema grandiflora* L.) under Prayagraj agro-climatic conditions. ISSN (E): 2277- 7695 ISSN (P): 2349-8242 NAAS Rating: 5.23 TPI 2021; 10(12): 248-245.
- Kumar, S., Kumar, M., Malik, S., Singh, M.K. and Kumar, S. (2014). Evaluation of Chrysanthemum (*Dendranthema grandiflora* Tzvelev) Genotypes using Morphological Characters Under Climatic Conditions of Western UP. *Annals of Horticulture* 7 (2): 162-165.
- Madhumathi, C., Bhargav, V., Reddy, D.S., Kameshwari, P.L., Sreedhar, D. and Lakshmi, T.N. (2018). Assessment of chrysanthemum (*Chrysanthemum morifolium*

Ramat.) germplasm for commercial cultivation under Rayalaseema region of Andhra Pradesh. *Journal of Applied Horticulture*, 20(3): 213-218.

- Negi, R., Dhiman, S.R. and Gupta, Y.C. (2019). Studies on Growth and Flowering Behavior of Newly Evolved Genotypes of Chrysanthemum (*Dendranthema grandiflora* Tzvelev) for Loose Flower Production. *International Journal of Current Microbiology and Applied Sciences*, 8(11): 341-346.

UNDER PEER REVIEW