

POTENTIAL EFFECT OF SOYBEAN ON MUSCLE FATIGUE AND LACTIC ACID LEVELS IN WISTAR RATS

ABSTRACT

Muscle fatigue is a condition resulting from prolonged muscle contractions. Synthetic materials are often used to restore muscle fatigue, although they can have side effects. Soybean is a natural product that contains complex compounds, including thiamine, which has the potential to replace synthetic compounds. The current study was aimed to analyze the potential effect of *Glycine max* on the structure of muscle tissue/muscle fatigue and lactic acid levels in Wistar rats. The study used an experimental approach, the treatment of giving soybean powder and pure thiamine compounds to 24 male Wistar white rats aged 3 months totaling 24 rats, 150-200 g for 15 days. Data analysis techniques were carried out on the frequency of muscle fatigue through the struggling test, measurement of lactic acid levels in the blood, and structure of the biceps femoris muscle tissue in HE staining. The results showed struggling time, Fobs 16.398 > Ftable 3.554 and lactic acid levels Fobs 17.621 > Ftable 3.467 and changes in the muscle tissue structure of white rats. The conclusion that soybean seeds have the potential as a natural ingredient in overcoming muscle fatigue and can reduce lactic acid levels.

Keywords: soybean, muscle fatigue, lactic acid, struggling, Wistar rats

1. INTRODUCTION (ARIAL, BOLD, 11 FONT, LEFT ALIGNED, CAPS)

Muscle fatigue is a condition of strong muscle contractions for a long time. Muscle fatigue results in decreased oxygen levels and increased lactic acid levels (Lamb, G. D., & Stephenson, 2006);(Wan, J. J., Qin, Z., Wang, P. Y., Sun, Y., & Liu, 2017). Muscle fatigue occurs in anaerobic activities due to high-intensity activities, requiring fast energy and a short time. Muscle fatigue can result in a decrease in body capacity and condition (Pethick, J., Winter, S. L., & Burnley, 2021).

Reducing muscle fatigue can be done using vitamins B1, B6, and B12 (Wan, J. J., Qin, Z., Wang, P. Y., Sun, Y., & Liu, 2017); (Martel, J. L., Kerndt, C. C., Doshi, H., & Franklin, 2021). These vitamins are usually consumed in the form of synthetic compounds that are packaged and sold freely on the market. Various food and beverage supplements that are circulating in the market are often consumed by humans to reduce muscle fatigue. Synthetic chemicals are often used as dominant compounds, because they give a very fast effect and can restore conditions (Holdt, S. L., & Kraan, 2011); (Margot, J., Rossi, L., Barry, D. A.,

& Holliger, 2015);(Peng, H., Saunders, D. M., Sun, J., Jones, P. D., Wong, C. K., Liu, H., & Giesy, 2016); (Mardiana, M., Rachmawati, L., Sari, N. P., & Al Amien, 2022). Supplements are needed to reduce muscle fatigue because the mechanism of action is very short and immediately gives the expected effect (Peng, H., Saunders, D. M., Sun, J., Jones, P. D., Wong, C. K., Liu, H., & Giesy, 2016); (Bowtell, J., & Kelly, 2019);(Aćimović, M. G., Tešević, V. V., Smiljanić, K. T., Cvetković, M. T., Stanković, J. M., Kiprovski, B. M., & Sikora, 2020).

Various synthetic compounds of thiamine and pyridoxine are often used to treat muscle fatigue (Wan, J. J., Qin, Z., Wang, P. Y., Sun, Y., & Liu, 2017); (Titcomb, T. J., & Tanumihardjo, 2019). There are side effects caused by the use of synthetic materials, including tissue damage that can be caused by the use of synthetic materials, for example the occurrence of necrosis (Nisar, B., Sultan, A., & Rubab, 2018); (Huang, K., Hu, S., & Cheng, 2019). Various natural ingredients with thiamine and pyridoxine compounds are still not widely known, so they have not been widely used to treat muscle fatigue (Choi, S. K., Baek, S. H., & Choi, 2013); Wan, et al., 2017 ; (Hu, X., Zhang, Q., Zhang, Q., Ding, J., Liu, Y., & Qin, 2022). The use of natural ingredients with various compound complexities in them can provide more moderate results (Thomford, N. E., Senthane, D. A., Rowe, A., Munro, D., Seele, P., Maroyi, A., & Dzobo, 2018). Natural materials have a variety of complex compounds that have the potential to be used for health (Yuan, H., Ma, Q., Ye, L., & Piao, 2016);(Koparde, A. A., Doijad, R. C., & Magdum, 2019).

Thiamine is most commonly found in vegetable materials, especially legumes (Leguminoceae), one of which is soybeans (Annor, G. A., Ma, Z., & Boye, 2014); (Erbersdobler, H. F., Barth, C. A., & Jahreis, 2017). The value of thiamine in soybeans is quite large among other legumes, reaching 1.0 mg (Erbersdobler, et al., 2017). The use of soybeans to

reduce lactic acid levels and muscle fatigue (Watanabe, N., & Sakuda, 2010); (Ding, J. F., Li, Y. Y., Xu, J. J., Su, X. R., Gao, X., & Yue, 2011);(Mardiana, M., Kartini, A., Sutiningsih, D., Suroto, S., & Muhtar, 2023). The current study was aimed to analyze the potential effect of Glycine max on the structure of muscle tissue/muscle fatigue and lactic acid levels in Wistar rats

2. MATERIAL AND METHODS

Treatment of experimental animals

Wistar rats, 3 months old, 150-200 g, 24 heads. Acclimatization was carried out for 1 week in the rearing cage. Mice were divided into 3 treatment groups. Eight control group rats (P1), eight group rats (P2) were given soybeans, and eight group rats (P3) were given synthetic thiamine. The treatment was carried out for 15 days. One hour after the treatment was continued with a swimming test (struggling).

Determine the dose of soy

Soybeans are washed and finely ground. 100 g of soybeans contains 0.93 g of thiamine. The thiamine value of 0.93 g per 100 g soybeans is adjusted to the normal thiamine requirement per day. The normal thiamine requirement dose is converted to the normal thiamine requirement for rats, body weight 150-200 g and 0.3 mg/day is obtained. Giving soy 3 mg/day for 15 days (P2).

The struggling test

Conducted a swimming test on the 16th day for all rat treatment groups. Data collection was carried out 2 times with an interval of 15 minutes. The struggling test was carried out to analyze the swimming ability of the rats (Alver, 2014; Li, et al., 2016)

Surgery, and preparation

Surgery and removal of the biceps femoris was carried out on the 17th day Preparation of muscle tissue preparations stained with hematoxylin eosin (HE).

Data analysis

Data on lactic acid levels were analyzed using one way Anova SPSS 19 and

changes in muscle tissue structure were analyzed using a dyno eye and optilab microscope.

after administration of soy and thiamine compounds showed changes in the structure of the biceps femoris muscle tissue (Figure 1).

3. RESULTS AND DISCUSSION

Analysis of the muscle contraction activity of the Wistar rats by the strunggling test




Figure 1. Myofibril structure of the biceps femoris muscle, HE, 400X

- A. Mice after carrying out a strunggling test: Myofibrils are disorganized, the nucleus is at the edges, the arrangement is disorganized
- B. Rats after carrying out the strunggling test were then treated with soybeans: Miofibrils arranged in an orderly manner, the nuclei at the edges in an orderly arrangement
- C. After carrying out a strunggling test, rats are treated with synthetic thiamine: Myofibrils are in order, the nucleus is on the edge in an orderly arrangement

The results of observations of changes in the structure of muscle tissue in Figure 1 can be correlated with the results of lactic acid examination and strunggling time (Table 1).

Table 1. Lactic acid levels and strunggling time

| treatment | lactic acid levels (mmol/l) | | strunggling time (minutes) |
|----------------|-----------------------------|-------|----------------------------|
| | before | After | |
| P ₀ | 6.7 | 8.2 | 16.4 |
| P ₁ | 6.8 | 7.4 | 20.8 |
| P ₂ | 6.7 | 6.9 | 26.5 |

P₀ : Mice after carrying out a strunggling test

P₁ : Rats after carrying out the strunggling test were then

treated with soybeans
P₂ : After carrying out a strunggling test, rats are treated with synthetic thiamine

Table 2. One way Anova test table

| | | Sum of squares | df | Mean square | F | Sig. |
|---------------------|----------------|----------------|----|-------------|--------|------|
| LACTIC_ACID | Between groups | 6.947 | 2 | 3.474 | 16.398 | .000 |
| | Within groups | 4.449 | 21 | 0.212 | | |
| | Total | 11.396 | 23 | | | |
| STRUNGG_TIME | Between groups | 209.083 | 2 | 104.541 | 17.621 | .000 |
| | Within groups | 106.790 | 18 | 5.933 | | |
| | Total | 315.873 | 20 | | | |

Based on Table 2 shows Fobs > Ftable on lactic acid levels and struggling time, the Fobs value of lactic acid is 16.398 > Ftable 3.554 and the Fobs value of lactic acid is 17.621 > Ftable 3.467 with each value of Sig .000. Conclusion of the data in Table 2 that there is an effect of soy thiamine compounds on muscle fatigue and muscle tissue structure.

Muscle activity is a process of muscle contraction related to the use of adenosine triphosphate (ATP) as contraction energy (Li, J., King, N. C., & Sinoway, 2003); (Casas, M., Buvinic, S., & Jaimovich, 2014) ;(Sahlin, 2014). Contraction of muscle cells requires energy in the form of adenosine triphosphate (ATP), then ATP is hydrolyzed into adenosine diphosphate (ADP) and energy is used for contraction, this process can continue as long as the supply of intracellular ATP is still there.(Li, J., King, N. C., & Sinoway, 2003); (Casas, M., Buvinic, S., & Jaimovich, 2014); (Grassi, B., Rossiter, H. B., & Zoladz, 2015). The total supply of ATP in the body is very limited, further energy needs are obtained from ATP synthesis through the oxidative pathway of creatine phosphate, which is highly dependent on the availability of oxygen and glycogen reserves derived from glucose (Baker, J. S., McCormick, M. C., & Robergs, 2010); (Grassi, B., Rossiter, H. B., & Zoladz, 2015);(Hargreaves, M., & Spriet, 2020). The energy obtained from creatine phosphate is only for the needs of muscle contraction for a few moments, and then ATP will be fulfilled through non-oxidative (anaerobic) phosphorylation processes (Li, J., King, N. C., & Sinoway, 2003); (Casas, M., Buvinic, S., & Jaimovich,

2014); (Grassi, B., Rossiter, H. B., & Zoladz, 2015); (Bartlett, 2019); (Fitzgerald, 2020). Anaerobic metabolism utilizes glucose and glycogen through the process of glycolysis without oxygen to produce ATP and metabolic waste in the form of lactic acid (Grassi, B., Rossiter, H. B., & Zoladz, 2015);(Hargreaves, M., & Spriet, 2020).

Continuous muscle contraction can cause muscle fatigue, because ATP supplies are very limited. Muscle fatigue is a condition where muscle cells are no longer able to contract due to a lack of ATP (Casas, M., Buvinic, S., & Jaimovich, 2014); (Hargreaves, M., & Spriet, 2020). The neuromuscular junction is unable to transmit impulses and is accompanied by an accumulation of lactic acid (Allen, D. G., Lamb, G. D., & Westerblad, 2008), (Girard, O., Millet, G. P., Micallef, J. P., & Racinais, 2012);(Mukund, K., & Subramaniam, 2020). Muscle fatigue will cause pain due to ischemia of muscle tissue (Queme, L. F., Ross, J. L., & Jankowski, 2017);(Wirth, K. J., & Scheibenbogen, 2021). Muscle fatigue occurs due to a decrease in the percentage of muscle strength, recovery time for muscle fatigue, and the time it takes for fatigue to occur (Ament, W., & Verkerke, 2009);(Taylor, J. L., Amann, M., Duchateau, J., Meeusen, R., & Rice, 2016); (Bartlett, 2019).

The strunggling activity carried out by rats is strenuous activity so that the energy required will greatly increase when compared to the resting state, this will activate anaerobic metabolism in muscle cells to produce energy and will increase intracellular lactic acid levels. Accumulation of lactic acid in muscle cells

will cause intracellular acidosis and cause fatigue (Allen, D. G., Lamb, G. D., & Westerblad, 2008);(Mukund, K., & Subramaniam, 2020). Lactic acid in muscle cells will diffuse into the blood and increase blood plasma lactic acid levels (van Ginneken, V., Boot, R., Murk, T., van den Thillart, G., & Balm, 2004); (Seheult, J., Fitzpatrick, G., & Boran, 2017). The more strenuous physical activity will increase anaerobic metabolic processes, so that lactic acid levels also increase (Grassi, B., Rossiter, H. B., & Zoladz, 2015); (Bartlett, 2019);(Fitzgerald, 2020). Thiamine is capable of carrying out oxidative decarboxylation of pyruvate to acetyl CoA and can cause the entry of substrates into the Krebs cycle to form energy (Depeint, F., Bruce, W. R., Shangari, N., Mehta, R., & O'Brien, 2006); (Tylicki, A., Łotowski, Z., Siemieniuk, M., 2018). Glucose deficiency can be overcome by administering vitamin B6 which plays a role in the form of pyridoxal phosphate (PLP) as a coenzyme, especially in decarboxylation and other reactions related to protein metabolism (McAllister, 2014); (Stach, K., Stach, W., & Augoff, 2021); (Calderón-Ospina, C. A., & Nava-Mesa, 2020); (Ito, T., Ogawa, H., Hemmi, H., Downs, D. M., & Yoshimura, 2022). As a coenzyme for phosphorylation, PLP helps release glycogen from the liver and muscles so that it can be broken down into glucose which can serve as an energy source during activities (Calderón-Ospina, C. A., & Nava-Mesa, 2020). Vitamin B1 (thiamine) which functions to convert carbohydrates into energy, strengthens the nervous system, where thiamin will stimulate neurotransmitter work (Campos-Bedolla, P., Walter, F. R., Veszelska, S., & Deli, 2014); (Mikkelsen, K., & Apostolopoulos, 2019); (Tardy, A. L., Pouteau, E., Marquez, D., Yilmaz, C., & Scholey, 2020). (Boldrini, G. G., Martin Molinero, G., Perez Chaca, M. V., Ciminari, M. E., Moyano, F., Córdoba, M. E., ... & Gómez, 2022). Thiamine is useful in helping to overcome symptoms of fatigue because this vitamin can improve carbohydrate metabolism which is used to

produce energy and can reduce the buildup of lactic acid in muscles that are experiencing fatigue (Roosterman, D., & Cottrell, 2021).

Based on research data, the use of soybeans is very important for health. People in Indonesia consume soybeans as tempeh, tofu. Use of thiamine (vitamin B1) in Indonesia use of thiamine to treat beriberi, type 2 diabetes, immune disorder, anti-oxidant, anti-proliferative, anti-obesity and anti-inflammatory (Adaikalakoteswari, A., Rabbani, N., Waspadji, S., Tjokroprawiro, A., Kariadi, S. H., Adam, J. M., & Thornalley, 2012); (Hassan, M., Rahman, H., Yasmeen, B. N., Mukti, A., Haque, H., Khan, M., ... & Kabir, 2018); (Dukariya, G., Shah, S., Singh, G., & Kumar, 2020).

4. CONCLUSION

The use of soybeans in rats as experimental animals can reduce muscle fatigue as shown in the structure of muscle tissue, and can reduce lactic acid levels. Soybean can reduce lactic acid levels 0.6 mmol/l while thiamine reduces lactic acid levels 0.2 mmol/l

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript

Ethical Approval:

Animal Ethic committee approval has been collected and preserved by the author(s)

REFERENCES

1. Aćimović, M. G., Tešević, V. V., Smiljanić, K. T., Cvetković, M. T., Stanković, J. M., Kiproviski, B. M., & Sikora, V.S. (2020) 'Hydrolyates: By-products of essential oil

distillation: Chemical composition, biological activity and potential uses. *Advanced technologies*, 9(2), pp. 54–70.

2. Adaikalakoteswari, A., Rabbani, N., Waspadji, S., Tjokroprawiro, A., Kariadi, S. H., Adam, J. M., & Thornalley, P.J. (2012) (2012) 'Disturbance of B-vitamin status in people with type 2 diabetes in Indonesia—Link to renal status, glycemic control and vascular inflammation. *Diabetes Research and Clinical Practice*', 95(3), pp. 415–424.
3. Allen, D. G., Lamb, G. D., & Westerblad, H. (2008) 'Skeletal muscle fatigue: cellular mechanisms. *Physiological reviews*.'
4. Ament, W., & Verkerke, G.J. (2009) 'Exercise and fatigue. *Sports medicine*', 39, pp. 389–422.
5. Annor, G. A., Ma, Z., & Boye, J.I. (2014) 'Crops–legumes. *Food processing: principles and applications*', pp. 305–337.
6. Baker, J. S., McCormick, M. C., & Robergs, R.A. (2010) 'Interaction among skeletal muscle metabolic energy systems during intense exercise. *Journal of nutrition and metabolism*, 2010'.
7. Bartlett, M.F. (2019) 'Mechanisms that Limit Oxidative Phosphorylation during High-Intensity Muscle Contractions In Vivo.'
8. Boldrini, G. G., Martin Molinero, G., Perez Chaca, M. V., Ciminari, M. E., Moyano, F., Córdoba, M. E., ... & Gómez, N.N. (2022) 'Glycine max (soy) based diet improves antioxidant defenses and prevents cell death in cadmium intoxicated lungs. *Biometals*', 35(2), pp. 229–244.
9. Bowtell, J., & Kelly, V. (2019) 'Fruit-derived polyphenol supplementation for athlete recovery and performance. *Sports Medicine*', 49((Suppl 1)), pp. 3–23.
10. Calderón-Ospina, C. A., & Nava-Mesa, M.O. (2020) 'B Vitamins in the nervous system: Current knowledge of the biochemical modes of action and synergies of thiamine, pyridoxine, and cobalamin. *CNS neuroscience & therapeutics*', 26(1), pp. 5–13.
11. Campos-Bedolla, P., Walter, F. R., Veszelka, S., & Deli, M.A. (2014) 'Role of the blood–brain barrier in the nutrition of the central nervous system. *Archives of medical research*', 45(8), pp. 610–638
12. Casas, M., Buvinic, S., & Jaimovich, E. (2014) 'ATP signaling in skeletal muscle: from fiber plasticity to regulation of metabolism. *Exercise and sport sciences reviews*', 42(3), pp. 110–116.
13. Choi, S. K., Baek, S. H., & Choi, S.W. (2013) 'The effects of endurance training and thiamine supplementation on anti-fatigue during exercise. *Journal of exercise nutrition & biochemistry*', 17(4), p. 189.
14. Depeint, F., Bruce, W. R., Shangari, N., Mehta, R., & O'Brien, P.J. (2006) 'Mitochondrial function and toxicity: role of the B vitamin family on mitochondrial energy metabolism. *Chemico-biological interactions*', 163((1-2)), pp. 94–112.
15. Ding, J. F., Li, Y. Y., Xu, J. J., Su, X. R., Gao, X., & Yue, F.P. (2011) 'Study on effect of jellyfish collagen hydrolysate on anti-fatigue and anti-

oxidation. *Food Hydrocolloids*, 25(5), pp. 1350–1353.

16. Dukariya, G., Shah, S., Singh, G., & Kumar, A. (2020) 'Soybean and its products: Nutritional and health benefits. *J Nut Sci Heal Diet*', 1(2), pp. 22–29.

17. Erbersdobler, H. F., Barth, C. A., & Jahreis, G. (2017) 'Legumes in human nutrition. Nutrient content and protein quality of pulses. *Ernahrungs Umschau*', 64(9), pp. 134–139.

18. Fitzgerald, L.F. (2020) 'BIOENERGETIC MECHANISMS OF AGE-RELATED DIFFERENCES IN MUSCLE FATIGUE.'

19. van Ginneken, V., Boot, R., Murk, T., van den Thillart, G., & Balm, P.. (2004) 'Blood plasma substrates and muscle lactic-acid response after exhaustive exercise in common carp and trout: indications for a limited lactate-shuttle. *Animal biology*', 54(2), pp. 119–130.

20. Girard, O., Millet, G. P., Micalef, J. P., & Racinais, S. (2012) 'Alteration in neuromuscular function after a 5 km running time trial. *European journal of applied physiology*', 112, pp. 2323–2330.

21. Grassi, B., Rossiter, H. B., & Zoladz, J.A. (2015) 'Skeletal muscle fatigue and decreased efficiency: two sides of the same coin?. *Exercise and sport sciences reviews*', 43(2), pp. 75–83.

22. Hargreaves, M., & Spriet, L.L. (2020) 'Skeletal muscle energy metabolism during exercise. *Nature Metabolism*', 2(9), pp. 817–828.

23. Hassan, M., Rahman, H., Yasmeen, B. N., Mukti, A., Haque, H., Khan, M., ... & Kabir, A.L. (2018) 'Thiamine

deficiency-Beriberi—A forgotten disease. *Northern International Medical College Journal*', 10(1), pp. 351–354.

24. Holdt, S. L., & Kraan, S. (2011) 'Bioactive compounds in seaweed: functional food applications and legislation. *Journal of applied phycology*', 23, pp. 543–597.

25. Hu, X., Zhang, Q., Zhang, Q., Ding, J., Liu, Y., & Qin, W. (2022) 'An updated review of functional properties, debittering methods, and applications of soybean functional peptides. *Critical Reviews in Food Science and Nutrition*', pp. 1–16.

26. Huang, K., Hu, S., & Cheng, K. (2019) 'A new era of cardiac cell therapy: opportunities and challenges. *Advanced healthcare materials*', 8(2), p. 1801011.

27. Ito, T., Ogawa, H., Hemmi, H., Downs, D. M., & Yoshimura, T. (2022) 'Mechanism of pyridoxine 5'-phosphate accumulation in pyridoxal 5'-phosphate-binding protein deficiency. *Journal of Bacteriology*', 204(3), pp. e00521-21.

28. Koparde, A. A., Doijad, R. C., & Magdum, C.S. (2019) 'Natural products in drug discovery. In *Pharmacognosy-medicinal plants*. IntechOpen.'

29. Lamb, G. D., & Stephenson, D.G. (2006) 'Point: Counterpoint: Lactic acid accumulation is an advantage/disadvantage during muscle activity. *Journal of Applied Physiology*', 100(4), pp. 1410-1412.

30. Li, J., King, N. C., & Sinoway, L.I. (2003) 'ATP concentrations and muscle tension increase linearly with

- muscle contraction. *Journal of Applied Physiology*, 95(2), pp. 577–583.
31. Mardiana, M., Kartini, A., Sutningsih, D., Suroto, S., & Muhtar, M.S. (2023) 'Literature review: nutrition supplementation for muscle fatigue in athletes. *Jurnal Keolahragaan*', 11(1), pp. 10–23.
 32. Mardiana, M., Rachmawati, L., Sari, N. P., & Al Amien, T.N. (2022) 'WHEY PROTEIN, DAUN KELOR, KURMA, DAN KELELAHAN OTOT. *Kesehatan Masyarakat*', 2.
 33. Margot, J., Rossi, L., Barry, D. A., & Holliger, C. (2015) 'A review of the fate of micropollutants in wastewater treatment plants. *Wiley Interdisciplinary Reviews: Water*', 2(5), pp. 457–487.
 34. Martel, J. L., Kerndt, C. C., Doshi, H., & Franklin, D.S. (2021) 'Vitamin B1 (thiamine). In *StatPearls* [Internet]. StatPearls Publishing.'
 35. McAllister, C.H. (2014) 'Analysis of the biochemical and genetic properties of alanine aminotransferase enzymes.
 36. Mikkelsen, K., & Apostolopoulos, V. (2019) 'Vitamin B1, B2, B3, B5, and B6 and the Immune System. *Nutrition and immunity*', pp. 115–125.
 37. Mukund, K., & Subramaniam, S. (2020) 'Skeletal muscle: A review of molecular structure and function, in health and disease. *Wiley Interdisciplinary Reviews: Systems Biology and Medicine*', 12(1), p. e1462.
 38. Nisar, B., Sultan, A., & Rubab, S.L. (2018) 'Comparison of medicinally important natural products versus synthetic drugs-a short commentary. *Nat. Prod. Chem. Res*', 6(2), p. 308.
 39. Peng, H., Saunders, D. M., Sun, J., Jones, P. D., Wong, C. K., Liu, H., & Giesy, J.P. (2016) 'Mutagenic azo dyes, rather than flame retardants, are the predominant brominated compounds in house dust. *Environmental Science & Technology*', 50(23), pp. 12669–12677.
 40. Pethick, J., Winter, S. L., & Burnley, M. (2021) 'Physiological complexity: Influence of ageing, disease and neuromuscular fatigue on muscle force and torque fluctuations. *Experimental Physiology*', 106(10), pp. 2046–2059.
 41. Queme, L. F., Ross, J. L., & Jankowski, M.P. (2017) 'Peripheral mechanisms of ischemic myalgia. *Frontiers in cellular neuroscience*', 11, p. 419.
 42. Roosterman, D., & Cottrell, G.S. (2021) 'Rethinking the citric acid cycle: Connecting pyruvate carboxylase and citrate synthase to the flow of energy and material. *International Journal of Molecular Sciences*', 22(2), p. 604.
 43. Sahlin, K. (2014) 'Muscle energetics during explosive activities and potential effects of nutrition and training. *Sports medicine*', 44((Suppl 2)), pp. 167–173.
 44. Seheult, J., Fitzpatrick, G., & Boran, G. (2017) 'Lactic acidosis: an update. *Clinical Chemistry and Laboratory Medicine (CCLM)*', 55(3), pp. 322–333.
 45. Stach, K., Stach, W., & Augoff, K. (2021) 'Vitamin B6 in health and disease. *Nutrients*', 13(9), p. 3229.
 46. Tardy, A. L., Pouteau, E., Marquez, D., Yilmaz, C., & Scholey, A. (2020) 'Vitamins and minerals for energy,

fatigue and cognition: a narrative review of the biochemical and clinical evidence. *Nutrients*, 12(1), p. 228.

47. Taylor, J. L., Amann, M., Duchateau, J., Meeusen, R., & Rice, C.L. (2016) 'Neural contributions to muscle fatigue: from the brain to the muscle and back again. *Medicine & Science in Sports & Exercise*', 48(11), pp. 2294–2306.
48. Thomford, N. E., Senthebane, D. A., Rowe, A., Munro, D., Seele, P., Maroyi, A., & Dzobo, K. (2018) 'Natural products for drug discovery in the 21st century: innovations for novel drug discovery. *International journal of molecular sciences*', 19(6), p. 157.
49. Titcomb, T. J., & Tanumihardjo, S.A. (2019) 'Global concerns with B vitamin statuses: biofortification, fortification, hidden hunger, interactions, and toxicity. *Comprehensive reviews in food science and food safety*', 18(6), pp. 1968–1984.
50. Tylicki, A., Łotowski, Z., Siemieniuk, M., & R. (2018) 'A Thiamine and selected thiamine antivitamin—biological activity and methods of synthesis. *Bioscience Reports*', 38(1).
51. Wan, J. J., Qin, Z., Wang, P. Y., Sun, Y., & Liu, X. (2017) 'Muscle fatigue: general understanding and treatment. *Experimental & molecular medicine*', 49(10), pp. e384-e384.
52. Watanabe, N., & Sakuda, S. (2010) 'Effect of dietary fermented soybean on physical endurance in mice. *Cellulose*', 11(1.97).
53. Wirth, K. J., & Scheibenbogen, C. (2021) 'Pathophysiology of skeletal muscle disturbances in myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS). *Journal of Translational Medicine*', 19(1), p. 162.
54. Yuan, H., Ma, Q., Ye, L., & Piao, G. (2016) 'The traditional medicine and modern medicine from natural products. *Molecules*', 21(5), p. 559.