

Enumeration of genetic parameters and genetic diversity of morpho-physiological traits in CIMMYT bread wheat accessions [*Triticum aestivum* (L.) em. Thell]

Abstract

Fifty genotypes of CIMMYT bread wheat were evaluated at Agricultural Research Farm, BHU, Varanasi during 2019-2020. The analysis of variance revealed significant differences among genotypes for all traits. High Phenotypic coefficient of variation was recorded compared to the genotypic coefficient of variation. However, high genotypic coefficients of variation were found particularly for: grain yield per plot (8.71), harvest index (9.22), test weight (8.9), normalized difference vegetative index (9.59) and chlorophyll content (9.79), suggesting that these traits are having ample genetic potential for selection amongst genotypes, in breeding programs. The highest broad sense heritability manifested for harvest index (91.61%); remaining traits showed moderate estimates of heritability. Low to moderate genetic advance as percent mean was estimated for all the traits studied. This suggests the existence of variability for agronomic traits in the studied wheat genotypes which, should be exploited during future breeding programmes. Fifty genotypes were divided into six non-overlapping distinct clusters using tocher's method based on Euclidean distances. Thirty-one genotypes were classified in the first cluster accounting 62% of total genotypes followed by 15 genotypes categorized in the second cluster. The remaining four clusters have one genotype each. Divergence and cluster mean show that, crossings between genotypes of clusters (II, III), and VI could lead to recovery of good transgressive segregants for maximum heterosis in wheat varietal improvement.

Keywords: PCV, GCV, Heritability, Genetic advance as percent mean, Genetic diversity, Genetic variability

1. Introduction

Wheat (*Triticum aestivum* (L.) em. Thell; $2n = 6x = 42$) is one of the most important cereals of the world, belonging to the Poaceae family. It, accounts for one-sixth of all agricultural land (Kumar, 2014). Wheat is cultivated on about 220.83 mha with an annual yield of about 769.31 million tonnes (Giraldo et al., 2019). It is anticipated that by 2050, demand for wheat will have expanded by 50%, from current levels. There is no way to expand the area under production (CIMMYT, 2022). The only remaining option therefore, is to boost productivity by developing better high yielding wheat varieties and better management of crop production.

The Global Wheat Program of the International Maize and Wheat Improvement Centre (CIMMYT), is one of the most important public sources of nutritious, high-yielding, disease- and climate-resilient wheat varieties for Africa, Asia, and Latin America. CIMMYT is therefore, a central pillar for more resilient agri-food systems in the above said continents countries (Dreisigacker et al., 2021). CIMMYT breeding lines can be traced in the pedigree of varieties sown on over 60 mha, around the world (CIMMYT, 2022).

Grain yield is a highly complex polygenic trait that is influenced by many component traits and the environment (Kumar et al., 2017). Increasing yield through direct selection is difficult thus, component traits should be considered while selecting to increase yield. Understanding the genetic makeup of various yield traits, is crucial for this purpose (Kumar et al., 2017). Variability can be additionally partitioned into phenotypic coefficient of variation (PCV) and genotypic coefficient of variation (GCV) which, further aids in the selection of potential genotypes from amongst the available germplasm. The coefficient of variation simply reveals the level of variability existing for traits, but it provided no information on the heritable portion of that existing variability (Nzuve et al., 2014 *Incomplete reference in the list*). In order to analyse the relative contributions of genetic and non-genetic components to the overall phenotypic variance in a population, it is important to consider both, heritability, and genetic variability. Genetic advance is the estimation of expected gain resulting from selection pressure on the breeding material. High heritability associated with high genetic advance for different yield components has a better scope for selecting high-yielding genotypes (Yadaw et al., 2015 *Absent in the reference list*). Hence heritability and genetic advance are critical selection parameters.

Crop genetic diversity provides an opportunity to develop varieties that are widely adapted to a specific environment (Govindaraj et al., 2015 *Incomplete reference in the list*). In order to address changing end-user demands, increase productivity and adaptability to unpredictable climate change are desirable in a crop. Crop genetic diversity is regarded as a source of novel alleles for crop development (Tester and Langridge, 2010). The greater the genetic separation between parents, the more is the possibility to produce heterosis in progeny (Joshi and Dhawan, 1966). Therefore, the present investigation was carried out to achieve information on various genetic parameters and genetic diversity existing in important morpho-physiological traits in CIMMYT bread wheat accessions.

2. Material and methods

2.1. Plant materials and phenotypic evaluations

Material used for the experiment was sown during *Rabi* season 2019-20 at the Agriculture Research Farm, Banaras Hindu University, Varanasi, consisting of 49 diverse genotypes of bread wheat collected from CIMMYT, Mexico along with local check variety DBW 187 (Supplementary Table 1). The material was sown in randomized block design (RBD) with 2 replications by maintaining inter-plot gap of 50 cm and within each replication, genotypes were sown in 6 rows of 5 m length with an inter-row spacing of 20 cm and plant to plant spacing of 5 cm. Recommended agronomic practices and plant protection measures were followed to raise a healthy crop. Weather conditions during the crop season are given in Supplementary Figure 1.

The observations were recorded on thirteen traits like germination percentage (GNP), days to 50 % flowering (DTFF), days to maturity (DTM), chlorophyll content (CFC), normalized difference vegetative index (NDVI), canopy temperature (CNTP) in °C, plant height (PTH) in cm, spike length (SKL) in cm, tillers per square metre (TPSM), test weight (TWT) in grams, biological yield per plot (BYPP) in kg, grain yield per plot (GYPP) in kg, harvest index (HI) in %. With a Minolta SPAD-502 Chlorophyll metre, CFC was measured at the heading and anthesis phases, while CNTP was measured from the vegetative stage to dough stages using, a hand-held infrared thermometer.

2.2 Statistical analysis

Descriptive statistics and analysis of variance (Panse and Sukhatme (1969) model) were carried out using INDOSTAT software. GCV and PCV were calculated as per the standard formula suggested by Searle (1961). The formula provided by Allard (1960) was used to calculate genetic advance as percentage of mean (GAM) and broad sense heritability (h^2_b). Mahalanobis (1936) D^2 statistic was used to measure the genetic divergence among 50 genotypes.

3. Results and discussion

3.1 Phenotypic evaluation

ANOVA showed significant differences among the genotypes indicating presence of sufficient variability in the germplasm for the studied traits suggesting, ample scope for further improvement (Table 1). Similar results were also reported by Arya et al. (2017 Absent in the reference list), and Upadhyay et al. (2020).

Table 1: Analysis of variance for 13 morpho-physiological traits in 50 bread wheat accessions

Sl.No.	Source	Mean Sum of Squares (MSS)		
		Replication	Treatment	Error
	Degrees of freedom	1	49	49

1	GNP	4.0760	48.499**	20.953
2	DTFF	11.4580	39.868**	20.3
3	CNTP	6.6820	9.11*	4.656
4	NDVI	0.8850	103.58**	31.696
5	DTM	128.6180	188.278**	85.935
6	CFC	22.5630	47.931**	12.449
7	TPSM	0.3270	155.782**	62.202
8	PTH	13.3740	100.39*	51.31
9	SKL	0.6270	2.526**	1.286
10	BYPP	0.0180	1.817**	0.923
11	TWT	14.3110	63.024**	32.037
12	HI	1.610	18.455**	0.807
13	GYPP	0.0010	0.231**	0.118

* Significant at P = 0.05, ** Significant at P = 0.01

Where, GNP- Germination Percentage; DTFF- Days to 50 % Flowering; DTM- Days to Maturity; CFC- Chlorophyll Content; NDVI- Normalized Difference Vegetative Index; CNTP- Canopy Temperature; PTH- Plant Height (cm); SKL- Spike Length(cm); TPSM- Tillers Per Square Metre; TWT- Test Weight (g); BYPP- Biological Yield Per Plot (Kg); GYPP- Grain Yield Per Plot (Kg); HI- Harvest Index (%).

The range and mean value of all the traits have been provided in Supplementary table 2. TPSM showed a wider variation of 61(ACC-12) to 103 (ACC-30) with a mean value of 85. The lowest and highest value of GNP was observed in ACC-29 (78%) and ACC-47 (110%) respectively with a mean value of 92%. The lowest and highest values for DTFF were observed in ACC-47 (70 days) and ACC-46 (88 days) respectively with an average value of 78 days. The lowest and highest values of CNTP were observed in ACC-42(22.13°C) and ACC-27 (32.74°C) respectively, with a population mean of 25.36°C. The lowest and highest values of NDVI recorded in ACC-22 (46.09) and ACC-37(82.14) respectively with an average value of 62.49. DTM exhibited a range from 103 (ACC-10) to 138 days (ACC-32,33) with a mean value of 117 days. CFC ranged from 34.67(ACC-23) to 56.25(ACC-17) with a mean of 42.99. For PTH lowest and highest values were recorded in ACC-18 (82.7 cm) and ACC-38 (117.1 cm) respectively with an average of 98.8 cm. SKL ranged from 9.4(ACC-9,11,31) to 14.9cm (ACC-14,9) with a mean of 11.34 cm. BYPP ranged from 6.85(ACC-18) to 11.66kg (ACC-32, ACC-33) with a population mean of 8.58 kg. GYPP ranged from 1.71(ACC-41) to 3.57kg (ACC-36) with a population mean of 2.72 kg. TWT ranged from 28.65 (ACC-41) to 55.03 g (ACC-6) with a population mean of 44.22 g. HI ranged from 24.41(ACC-41) to 39.11%(ACC-36) with a population mean of 32.19%. GYPP ranged from 1.71(ACC-41) to 3.57kg (ACC-36) with a population mean of 2.74.

3.2 Genotypic and phenotypic variability

The study revealed that, PCV were higher than their corresponding GCV for all the traits among the genotypes (Figure1). This indicates that, characters were influenced by the environment. Selection based on phenotype alone, can be effective for the traits where variation between PCV and GCV were less, means such traits are less influenced by the environment. These findings are in agreement with a previous study in wheat (Upadhyayet al.,2020).

Higher values of GCV were recorded for CFC (9.79) followed by NDVI (9.59), HI (9.22), TWT (8.9), and GYPP (8.71) suggesting that, these traits possess ample genetic potential for selection among genotypes in breeding programs^{The values less than 10 per cent comes under the low PCV and GCV}. Lower values however, were recorded for CNTP (5.88), DTM (5.79), GNP (4.04), DTFF (4.026).

Higher PCV values were observed for TWT (15.58), GYPP (15.27), BYPP (13.64), NDVI (13.16) while^{All are having moderate values}, lower values were recorded for GNP (6.42), DTFF (7.05) among the traits. A wide difference between GCV and PCV was observed for TWT (6.689), GYPP (6.566), SKL (5.234), and BYPP (5.847) respectively, indicating the influence of environment on these traits. However, the least difference was observed in HI (0.413), GNP(2.38), and CFC(2.98) indicating that, these traits have a high heritable genetic variation and are less influenced by the environment. The current results are in agreement with those of Arya et al. (2017Absent in the reference list), Kumar et al. (2019Either Absent or mismatch in the year in the reference list), and Upadhyayet al. (2020).

3.3 Heritability and genetic advance

Estimates of heritability and GAM are presented in Supplementary Table2. The highest value of heritability was revealed by HI (91.61), while all remaining traits showed moderate estimates of heritability. Heritability estimations aid plant breeders in choosing superior genotypes from a variety of genetic groups. Similar results were also observed by Mohsin et al. (2019Mismatch in the year mentioned in the list), Bayisa et al. (2020), and Bhanu et al. (2018).The highest GAM was recorded for HI (18.19) and lowest for GNP (5.25). Similar findings have also been reported by Mohsin et al. (2019Mismatch in the year mentioned in the list), Upadhyayet al. (2020), Roy et al. (2021), and Yadav et al. (2021).

A character with high heritability does not necessarily also have high genetic advance, according to Johnson et al. (1955). A character may be assessed more precisely when heritability and genetic advance are investigated together, (Dudley and Moll, 1969). Consequently, Heritability estimates and GAM are typically more useful in finding gene action that is involved in the expression of different polygenic traits along with, estimating the

gain. High heritability coupled with moderate GAM was observed for HI (91.61 and 18.19) while moderate heritability with moderate GAM was recorded for NDVI (53.13 and 14.40), TPSM (42.93 and 10.89), CFC (58.76 and 15.47), TWT (32.59 and 10.46), and GYPP (32.51 and 10.23). Remaining traits exhibited moderate heritability coupled with, low GAM. Estimates of heritability, genetic advance, PCV, GCV for all the traits are displayed in Figure 1.




Figure 1: Graphical representation of genetic parameters of variation (GCV, PCV, Genetic Advance (5%), Genetic advance as percent of mean (5%), h^2_b for 13 traits among 50 bread wheat accessions

*Abbreviations: GCV-Genotypic coefficient of variation, PCV-phenotypic coefficient of variation, h^2_b -Broad Sense heritability

3.4 Genetic divergence (D^2) Analysis

One effective method of determining genetic divergence is D^2 analysis (Mahalanobis, 1936). To calculate the relative contribution of each component trait to the overall divergence and to measure the degree of differentiation at the intra- and inter-cluster levels, D^2 statistics is useful. Fifty genotypes were divided into six non-overlapping distinct clusters using Tocher's method based on, Euclidean distances so that, the genotypes within each cluster had lower D^2 values than the genotypes between the clusters (Figure 2). Similar findings have been reported by Kumar et al. (2013), Tsegaye et al., (2014), Arya et al., (2017 Absent in the reference list), Singh et al., (2017 Only two authors mentioned in the reference list) Santosh et al., (2019 Absent in the reference list), and Devesh et al, (2019) Tocher's method for genotypes clustering. Table 2 shows the percent contribution of each trait towards total genetic divergence. The more a trait contributes to diversity, the more times it ranked first. The traits like GYPP ranked a greater number of times (196) in first position and showed the

highest percent contribution (15.65%) towards total genetic divergence followed by, TWT (13.55), and PTH (13.34). Lowest percent contribution was reported by the trait CNTP (0.24) followed by, DTFF (1.71).

Table 2: Percent contribution of each trait towards total genetic divergence

Sl.No.	Source	Contribution(%)	Timesranked1 st
1	GNP	4.24	53
2	DTFF	1.71	21
3	CNTP	0.24	3
4	NDVI	8.5	106
5	DTM	6.54	82
6	CFC	4.98	62
7	TPSM	2.38	30
8	PTH	13.34	167
9	SKL	9.62	120
10	BYPP	6.9	86
11	TWT	13.55	169
12	HI	12.35	154
13	GYPP	15.65	196

Cluster I had the most genotypes (31), followed by cluster II (15), while clusters III, IV, V, VI had just one genotype each (Table 3). Even though a particular cluster was produced, the genotypes within it were obtained from various locations. The genotype clustering pattern showed that, the genotypes acquired from the same location did not make up a single cluster. This shows that genetic and geographic diversity are not always related. Similar results were also reported by Upadhyayet al. (2020).

Table 3: Clustering of 50 bread wheat accessions by tocher's optimization method

ClusterGroup	No. OfGenotypes	ListofGenotypes
1Cluster	31	ACC-32, ACC-33, ACC-50, ACC-26, ACC-40, ACC-44, ACC-49, ACC-35, ACC-34, ACC-4, ACC-9, ACC-8, ACC-15, ACC-31, ACC-13, ACC-36, ACC-23, ACC-39, ACC-27, ACC-45, ACC-14, ACC-42, ACC-43, ACC-46, ACC-48, ACC-24, ACC-1, ACC-30, ACC-19, ACC-28 & ACC-3
2Cluster	15	ACC-2, ACC-5, ACC-16, ACC-21, ACC-7, ACC-25, ACC-20, ACC-18, ACC-12, ACC-6, ACC-17, ACC-11, ACC-22, ACC-10 & ACC-29

3Cluster	1	ACC-38
4Cluster	1	ACC-41
5Cluster	1	ACC-37
6Cluster	1	ACC-47




Figure 2: Dendrogram of 50 bread wheat accessions for 13 traits

In the current study, intra-cluster distances were lower than inter-cluster distances indicating the presence of a high degree of genetic diversity (Figure 3). Cluster II recorded the largest intra-cluster distance (553.67), followed by cluster I (500.53). While clusters III, IV, V and VI had only one genotype, no intra-cluster distance was thus, found (Table 4). It may therefore be concluded that, genotypes inside a cluster with a high degree of divergence would offer more suitable breeding materials for crop improvement (Dobariya et al., 2006).

The inter-cluster distance values ranged from 553.13 to 1908.08 (Table 4). Cluster II and VI had the largest inter-cluster distance (1908.08), followed by cluster II and III (1774.78). The genotypes found in these clusters had a wide range of genetic diversity and might be employed in a wheat hybridization programme to recover better transgressive segregants, aiding the production of high yielding varieties. The smallest inter-cluster distances were found between clusters III and V (553.13), demonstrating a close association between these clusters and indicating that, they would not lead to satisfying outcomes.

Table 4: Average intra cluster (diagonal) – inter cluster distances

	Cluster1	Cluster2	Cluster3	Cluster4	Cluster5	Cluster6
Cluster1	500.53	970.94	796.76	814.50	862.76	1100.85
Cluster2	970.94	553.67	1774.78	1463.05	1180.62	1908.08
Cluster3	796.76	1774.78	0.00	1557.04	553.13	895.55
Cluster4	814.50	1463.05	1557.04	0.00	1228.44	1474.54
Cluster5	862.76	1180.62	553.13	1228.44	0.00	1083.11
Cluster6	1100.85	1908.08	895.55	1474.54	1083.11	0.00




Figure 3: Cluster diagram of 50 bread wheat accessions using tocher method

The cluster means of all studied traits are presented in Table 5. The differences between the clusters for each trait were revealed by the values of the cluster means for those traits. GNP cluster mean values ranged from 88.13 (cluster IV) to 110.75 (cluster VI). Cluster means of DTFF varied from 70.65 (cluster II) to 84.22 cm (cluster V). For CNTP cluster means ranged from 24.53 (cluster VI) to 27.57 (cluster IV). For NDVI cluster means ranged from 59.50 (cluster II) to 82.14 (cluster V). DTM cluster mean values ranged from 111.76 (cluster II) to 133.02 (cluster III and IV). CFC cluster mean values ranged from 38.44 (cluster III) to 48.47 (cluster VI). TPSM cluster mean values ranged from 77.71 (cluster II) to 96.92 (cluster VI). For PTH cluster means ranged from 84.53 (cluster IV) to 117.12 (cluster III). For SKL cluster means ranged from 10.42 (cluster IV and VI) to 12.55 (cluster III). For BYPP cluster means ranged from 7.16 (cluster IV) to 9.72 (cluster V). Cluster means of TWT varied from 28.65 (cluster IV) to 48.42 (cluster II). Cluster means of HI varied from 24.41 (cluster IV) to 33.79 (cluster II). GYPP Cluster means varied from 1.71 (cluster IV) to 3.20 (cluster I). The highest mean value for DTM and the lowest for GYPP were observed among all

clusters. Results on divergence and cluster mean reveal that, crossings between genotypes of clusters II, III, and VI could lead to achievement of maximum heterosis and recovery of better transgressive segregants; for wheat improvement (Arya et al. 2017 Absent in the reference list).

Table 5: Cluster Means for 13 morpho -physiological traits in 50 bread wheat accessions

	GN P	DT FF	CN TP	ND VI	DT M	C F C	TP SM	PT H	S K L	BY PP	T W T	HI	GY PP
Cluster1	91. 48	78. 97	25. 28	62. 20	128. 48	42 .6 1	87. 39	100 .39	11. 59	8.8 9	42. 69	31. 69	2.7 9
Cluster2	91. 15	74. 64	25. 44	59. 50	111. 76	43 .3 2	77. 71	94. 78	10. 88	7.9 0	48. 42	33. 79	2.6 5
Cluster3	93. 09	83. 09	24. 54	76. 45	133. 02	38 .4 4	89. 91	117 .12	12. 55	9.1 0	44. 79	30. 29	2.7 0
Cluster4	88. 13	79. 13	27. 57	66. 96	133. 02	44 .7 1	86. 41	84. 53	10. 42	7.1 6	28. 65	24. 41	1.7 1
Cluster5	88. 14	84. 22	26. 05	82. 14	116. 83	46 .9 7	88. 41	105 .35	10. 96	9.7 2	42. 19	32. 95	3.2 0
Cluster6	110 .75	70. 65	24. 53	78. 35	133. 01	48 .4 7	96. 92	100 .83	10. 42	8.9 0	45. 76	32. 48	2.8 9

Conclusion

Wide range of variability for most of the traits were observed among the germplasm accessions evaluated, indicating a considerable scope for the selection of desired genotypes that contribute directly or indirectly towards crop improvement. High heritability with high genetic advance was recorded for grain yield per plot, representing the contribution of on additive gene effect and the probability of improving the traits, by selection. It is possible to determine the type of gene action involved in the manifestation of various polygenic traits as well as, estimate the genetic advance, under selection, by using the derived heritability estimates and genetic advance. Divergence and cluster mean show that genotype crosses among clusters II, III, and VI may lead to higher heterosis and, recovery of best transgressive segregants for the development of better wheat varieties/hybrids.

References

- Allard, R.W. (1960). *Principles of Plant Breeding*. John Wiley and Sons. Inc, New York
- Bayisa, T., Tefera, H., & Letta, T. (2020). Genetic variability, heritability and genetic advance among bread wheat genotypes at South-eastern Ethiopia. *Agriculture, Forestry and Fisheries*, **9**(4), 128.
- Belderok, B., Mesdag, J., Mesdag, H., & Donner, D. A. (2000). *Bread-making quality of wheat: a century of breeding in Europe*. Springer Science & Business Media. This reference is Absent in the text
- Bhanu, A. N., Arun, B., & Mishra, V. K. (2018). Genetic variability, heritability and correlation study of physiological and yield traits in relation to heat tolerance in wheat (*Triticumaestivum* L.). *Biomedical Journal of Scientific & Technical Research*, **2**(1), 2112-2116.
- CIMMYT, 2022 <https://www.cimmyt.org/work/wheat-research/>
- Devesh, P., Moitra, P. K., Shukla, R. S., & Pandey, S. (2019). Genetic diversity and principal component analyses for yield, yield components and quality traits of advanced lines of wheat. *Journal of Pharmacognosy and Phytochemistry*, **8**(3), 4834-4839.
- Dobariya, K.L., Ribadia, K.H., Padhar, P.R. and Ponkia, H.P. (2006). Analysis of genetic divergence in some synthetic lines of bread wheat (*Triticumaestivum* L.). *Advances in Plant Sciences*. **19**(1): 221-225
- Dreisigacker S, Crossa J, Pérez-Rodríguez P, Montesinos-López OA, Rosyara U, Juliana P, Mondal S, Crespo-Herrera L, Govindan V, Singh RP, Braun HJ. Implementation of genomic selection in the CIMMYT global wheat program, findings from the past 10 years. *Crop Breeding, Genetics and Genomics*. 2021 Mar 26;**3**(2).
- Dudley JW, Moll RH. Interpretation and use of estimates of heritability and genetic variances in plant breeding 1. *Crop science*. 1969; **9**(3):257-262.
- Giraldo P, Benavente E, Manzano-Agugliaro F, Gimenez E. Worldwide Research Trends on Wheat and Barley: A Bibliometric Comparative Analysis. *Agronomy*. 2019; **9**(7):352. <https://doi.org/10.3390/agronomy9070352>
- Govindaraj, M., Vetriventhan, M., & Srinivasan, M. (2015). Importance of genetic diversity assessment in crop plants and its recent advances: an overview of its analytical perspectives. *Genetics research international*, 2015. **Incomplete reference**

- Gupta, R. K. (2004). Quality of Indian wheat and infrastructure for analysis. *A compendium of the training program (26–30 December, 2003) on wheat improvement in eastern and warmer regions of India: conventional and non-conventional approaches*. NATP project, (ICAR), BHU, Varanasi, India. This reference is Absent in the text
- Johnson, H. W., Robinson, H. F., & Comstock, R. E. (1955). Estimates of genetic and environmental variability in soybeans 1. *Agronomy journal*, **47**(7), 314-318.
- Kumar, B., Singh, C. M., & Jaiswal, K. K. (2013). Genetic variability, association and diversity studies in bread wheat (*Triticumaestivum* L.). *The Bioscan*, **8**(1), 143-147.
- Joshi, A. B., & Dhawan, N. L. (1966, January). Genetic improvement in yield with special reference to self-fertilizing crops. In *Indian Journal of Genetics and Plant Breeding* (pp. 101-+). PO BOX 11312, INDIAN AGRICULTURE RES INST, NEW DELHI 110012, INDIA: INDIAN SOC GENET PLANT BREEDING. Delete
- Kumar, B., Singh, C. M., & Jaiswal, K. K. (2013). Genetic variability, association and diversity studies in bread wheat (*Triticumaestivum* L.). *The Bioscan*, **8**(1), 143-147. Repeated.
- Kumar, S. (Ed.). (2014). *The Role of Bioinformatics in Agriculture* (1st ed.). Apple Academic Press. <https://doi.org/10.1201/b16568>
- Kumari, M., Kumar, M., Singh, V., Kumar, S. V., & Rathi, M. (2017). Trait association and morphological diversity in wheat (*Triticumaestivum* L.) genotypes. *Electronic Journal of Plant Breeding*, **8**(2), 534-540.
- Mahalanobis, P. C. (1936). On the generalized distance in statistics. National Institute of Science of India.
- Mohsin, T., Khan, N., & Naqvi, F. N. (2009 Mismatch in the year mentioned in the text). Heritability, phenotypic correlation and path coefficient studies for some agronomic characters in synthetic elite lines of wheat. *Journal of Food, Agriculture and Environment*, **7**(3-4), 278-282.
- Nzuve, F., Githiri, S., Mukunya, D. M., & Gethi, J. (2014). Genetic variability and correlation studies of grain yield and related agronomic traits in maize. Complete the reference
- Panse VG, Sukhatme PV. *Statistical methods for agricultural workers*. Indian Council of Agricultural Research, New Delhi, 1969.

- Roy, A., Kumar, A., Singh, A., Mandi, A., & Barman, M. (2021). Analysis of genetic diversity and correlation studies on grain yield and its component characters in bread wheat (*Triticumaestivum* L. emThell) genotypes. *The Pharma Innovation Journal*, **10**(5), 341-345.
- Jaiswal, J. P., Singh, A., &Gahatyari, N. C. (2019). Assessment of genetic diversity for terminal heat tolerance in bread wheat (*Triticumaestivum* L. em. Thell.). *Journal of Pharmacognosy and Phytochemistry*, **8**(1), 2572-2579. This reference is Absent in the text
- Searle, S.R. (1961). Phenotypic, genotypic and environmental correlations. *Biometrics*. 17: 474-480.
- Singh, G., & Kumar, R. (2017). Genetic Diversity for yield and yield traits in wheat. *Journal of Pharmacognosy and Phytochemistry*, **6**(5), 2062-2066.
- Tester, M., &Langridge, P. (2010). Breeding technologies to increase crop production in a changing world. *Science*, **327**(5967), 818-822.
- Upadhyay, P., Krishna, S., Thakur, P., Agrawal, N., Yadav, P., Prasad, L. C., & Mishra, V. K. (2020). Identification of genetic variability and diversity in selected wheat (*Triticumaestivum* L.) germplasm under three different dates of sowing. *Journal of Pharmacognosy and Phytochemistry*, **9**(2), 82-86.
- Yadav, N., Dashora, A., Choudhary, U., Dave, M., &Rathore, J. (2021). Assessment of genetic variability, heritability and genetic advance in bread wheat (*Triticumaestivum* L.) Genotypes,*ThePharma Innovation Journal* 2021; **10**(9): 528-530

Supplementary Table 1: List of accessions procured from CIMMYT and their codes

S.no	Accession	Name
1	ACC-1	DBW 187
2	ACC-2	NADI #2
3	ACC-3	QUAIU #1/SUP152
4	ACC-4	WBLL1*2/BRAMBLING/4/BABAX/LR42//BABAX*2/3/SHAMA*2/5/PBW343*2/KUKUNA*2//FRTL/PIFED
5	ACC-5	KACHU #1/KIRITATI//KACHU/3/BECARD/FRNCLN
6	ACC-6	BABAX/LR42//BABAX*2/3/KUKUNA/4/BACEU #1/5/BECARD/6/KACHU/KINDE
7	ACC-7	BABAX/LR42//BABAX/3/ER2000/4/KACHU #1/KIRITATI//KACHU
8	ACC-8	KACHU*2/SUP152
9	ACC-9	SAUAL/YANAC//SAUAL/3/2*KFA/2*KACHU
10	ACC-10	KANCHAN*2/JUCHI//2*BORL14
11	ACC-11	WBLL1*2/BRAMBLING//WBLL1*2/BRAMBLING/3/2*BORL14
12	ACC-12	FRANCOLIN #1*2/HAWFINCH #1//2*MUCUY
13	ACC-13	PBW65/2*PASTOR//TACUPETO F2001*2/BRAMBLING/3/TACUPETO F2001*2/BRAMBLING/6/2*SHORTENED SR26 TRANSLOCATION/4/ATTILA/3*BCN//BAV92/3/PASTOR/5/MUNAL
14	ACC-14	MUU/KBIRD//2*KACHU/KIRITATI
15	ACC-15	MUU/KBIRD//2*KACHU/KIRITATI
16	ACC-16	WBLL1*2/BRAMBLING//VORB/FISCAL/3/BECARD/4/MUCUY/5/MUCUY


17	ACC-17	SUP152/BLOUK #1//BECARD/FRNCLN
18	ACC-18	ALTAR 84/AE.SQUARROSA (221)//3*BORL95/3/URES/JUN//KAUZ/4/WBLL1/5/MUTUS/6/SUP152/BAJ #1
19	ACC-19	BECARD/FRNCLN//BORL14
20	ACC-20	SNTL/3/KACHU//WBLL1*2/BRAMBLING
21	ACC-21	BORL14*2//BECARD/QUAIU #1
22	ACC-22	BORL14*2//BECARD/QUAIU #1
23	ACC-23	BORL14*2//BECARD/QUAIU #1
24	ACC-24	WBLL1*2/BRAMBLING/4/BABAX/LR42//BABAX*2/3/SHAMA*2/5/BECARD/QUAIU #1
25	ACC-25	NADI#1*2/3/MUTUS/AKURI #1//MUTUS
26	ACC-26	SHAKTI/6/WHEAR/KUKUNA/3/C80.1/3*BATAVIA//2*WBLL1/4/T.DICOCCON PI94625/AE.SQUARROSA (372)//SHA4/CHIL/5/WHEAR/KUKUNA/3/C80.1/3*BATAVIA//2*WBLL1
27	ACC-27	CHIPAK/3/SWSR22T.B./2*BLOUK #1//WBLL1*2/KURUKU
28	ACC-28	BORL14*2/MISR 1
29	ACC-29	SAUAL/3/ACHTAR*3//KANZ/KS85-8-4/4/SAUAL/5/2*BAJ #1/3/KIRITATI//ATILA*2/PASTOR
30	ACC-30	SUP152/BLOUK #1//2*BECARD/FRNCLN
31	ACC-31	MUTUS*2/MUU//2*MUCUY
32	ACC-32	BECARD/FRNCLN*2//BORL14
33	ACC-33	KACHU/BECARD//WBLL1*2/BRAMBLING*2/3/ONIX/KBIRD
34	ACC-34	KACHU/BECARD//WBLL1*2/BRAMBLING*2/3/FRNCLN*2/TECUE #1
35	ACC-35	KACHU/BECARD//WBLL1*2/BRAMBLING*2/3/KACHU/KINDE
36	ACC-36	KACHU/BECARD//WBLL1*2/BRAMBLING*2/8/REH/HARE//2*BCN/3/CROC_1/AE.SQUARROSA

		(213)//PGO/4/HUITES/5/T.DICOCCON PI94624/AE.SQUARROSA (409)//BCN/6/REH/HARE//2*BCN/3/CROC_1/AE.SQUARROSA (213)//PGO/4/HUITES/7/MUTUS
37	ACC-37	KACHU #1/3/T.DICOCCON PI94624/AE.SQUARROSA (409)//BCN/4/2*KACHU*2/5/KACHU/KINDE
38	ACC-38	NADI#2*2/6/BECARD #1/5/KIRITATI/4/2*SERI.1B*2/3/KAUZ*2/BOW//KAUZ
39	ACC-39	MEX94.27.1.20/3/SOKOLL//ATTILA/3*BCN/4/2*SUP152*2/TECUE #1
40	ACC-40	MACE/5/TILILA/JUCHI/4/SERI.1B//KAUZ/HEVO/3/AMAD/6/KACHU/BECARD//WBLL1*2/BRAMBLING
41	ACC-41	KACHU//KIRITATI/2*TRCH/3/KFA/2*KACHU
42	ACC-42	KACHU//KIRITATI/2*TRCH/3/KFA/2*KACHU
43	ACC-43	MUU/FRNCLN//FRANCOLIN #1/3/BORL14
44	ACC-44	WBLL1*2/BRAMBLING*2//BAVIS/3/KACHU #1/KIRITATI//KACHU
45	ACC-45	WBLL1*2/BRAMBLING*2//BAVIS/3/KFA/2*KACHU
46	ACC-46	KENYA SUNBIRD/2*KACHU/3/WBLL1*2/BRAMBLING*2//BAVIS
47	ACC-47	SUNCO.6/FRAME//PASTOR/3/NAVJ07/4/1447/PASTOR//KRICHAUFF/5/BORL14
48	ACC-48	BABAX/LR42//BABAX*2/3/SHAMA/4/TACUPETO F2001*2/BRAMBLING/5/BORL14
49	ACC-49	KACHU*2/3/ND643//2*PRL/2*PASTOR/4/KIDEA
50	ACC-50	BORL14*2/FITIS


Supplementary Table 2: Genetic variability parameters for 13 morpho -physiological traits in 50 bread wheat accessions

Sl. No.	Traits	Minimum	Maximum	mean	V _g	GCV	V _p	PCV	h ² _b	GA(5 %)	GAM(5 %)
1	GNP	78	110	92	13.773	4.049	34.726	6.429	39.662	4.815	5.252
2	DFFF	70	88	78	9.784	4.026	30.084	7.059	32.522	3.675	4.729
3	CNTP	22.13	32.74	25.36	2.227	5.885	6.883	10.345	32.359	1.749	6.896
4	NDVI	46.09	82.14	62.49	35.942	9.594	67.638	13.161	53.139	9.003	14.407
5	DTM	103	138	117	51.171	5.792	137.106	9.481	37.322	9.003	7.289
6	CFC	34.67	56.25	42.99	17.741	9.798	30.19	12.782	58.764	6.651	15.473
7	TPSM	61	103	85	46.79	8.073	108.992	12.322	42.93	9.233	10.897
8	PTH	82.7	117.1	98.8	24.54	5.012	75.85	8.812	32.353	5.804	5.873
9	SKL	9.4	14.9	11.34	0.62	6.945	1.906	12.179	32.524	0.925	8.16
10	BYPP	6.85	11.66	8.58	0.447	7.795	1.37	13.642	32.653	0.787	9.176
11	TWT	28.65	55.03	44.22	15.493	8.9	47.531	15.589	32.596	4.629	10.468
12	HI	24.41	39.11	32.19	8.824	9.229	9.631	9.642	91.618	5.857	18.197
13	GYPP	1.71	3.57	2.74	0.057	8.711	0.175	15.277	32.518	0.28	10.233

Where, GNP- Germination Percentage; DFFF- Days to 50 % Flowering; DTM- Days to Maturity; CFC- Chlorophyll Content; NDVI- Normalized Difference Vegetative Index; CNTP- Canopy Temperature; PTH- Plant Height (cm); SKL- Spike Length (cm); TPSM- Tillers Per Square Metre; TWT- Test Weight (g); BYPP- Biological Yield Per Plot (Kg); GYPP- Grain Yield Per Plot (Kg); HI- Harvest Index (%), V_g-Genotypic variance, GCV- Genotypic coefficient of variation, V_p- Phenotypic variance, PCV-Phenotypic coefficient of variation, h²_b -Broad sense heritability, GA-Genetic advance, GAM-Genetic advance as percent mean



Supplementary Figure 1: Mean weekly meteorological data (rainfall in mm, temperature maximum in°C , temperature minimum°C, RH% morning, RH% evening, sunshine hours, evaporation in mm) during 2019-20 *Abbreviations- RH-Relative humidity



Supplementary Figure 2: Pie chart showing the Percent contribution of each trait towards total genetic divergence among 50 bread wheat accessions

UNDER PEER REVIEW