

# **Morpho-molecular characterization of carrot soft rot incitant, *Pectobacterium carotovorum* subsp *carotovorum***

## **ABSTRACT :**

Carrot is an important root vegetable which plays an important role in human health. Globally, Post harvest diseases are the major constraint in carrot production, especially soft rot which resulting in severe yield loss. Early diagnosis of these post-harvest diseases paves a way for reducing the economic losses. Carrot samples showing typical rotting symptoms were collected from markets of four different districts of Tamil Nadu and the pathogen involved were isolated. Severe carrot soft rot incidence (66.74%) was observed in samples collected from Ooty area of The Nilgiris district and the least disease incidence of (16.21%) was recorded in Perundurai of Erode district. Pathogenicity of soft rot pathogen were conducted and the virulent isolates were identified based on Percent Disease Index (PDI) showing >50% rotting falls under disease grade 9 using disease grade scale (0 to 9). The bacterial isolates (KPB-7 and OCB-5) causing soft rot were characterized using various biochemical assays where in they showed positive response for methyl red, H<sub>2</sub>S gas production, KOH and catalase tests besides showing negative response for gram's reaction. Furthermore, molecular characterization of 16s rRNA region revealed the soft rot isolate (KPB-7) as *Pectobacterium carotovorum* subsp *carotovorum* (with an accession number OR251119).

Keywords : Carrot ; Soft rot ; Morphological ; Molecular characterization

## **1. INTRODUCTION**

Carrot is a biennial flowering plant of Apiaceae family and cultivated worldwide for its fleshy edible root and its nutritional status. It is a rich source of alpha and beta carotene which also contain Vitamins (A, K and B6) and minerals helps to improve eye vision and widely used for culinary purposes. In India, it covers an area of 110 thousand hectares with a production of 386.39 thousand tonnes (APEDA,2022) and Haryana is the leading producer of carrot. In Tamil Nadu, major carrot growing districts are Nilgiris, Dindigul and Krishnagiri (2022). Post harvest loss is a major constraint in carrot cultivation and around 20-60% postharvest losses was observed in vegetables Kitinoja *et al.* [1]. Even 50-100% economic losses were recorded due to post harvest infections from field to storage Bhat *et al.* [2]. Chances of infection on harvested products would be high during harvest, transportation and storage and spreads through wounds of damaged plants causing economic damage to fleshy vegetables Whitehead *et al.* [3]. In vegetables, greater loss occurs mainly due to the soft rot and sour rot diseases Bhat *et al.* [2]. Soft rot caused by *Pectobacterium carotovorum* subsp *carotovorum* produces cell wall degrading enzymes such as cellulase, pectinase and polygalacturonase which act as a virulent factor for disease development. Early days, identification of microbes were done based on phenotypic character and biochemical tests De Boer and Kelman [4]. In recent years, molecular characterization of organisms is essential for confirming their identity wherein PCR (Polymerase Chain reaction) has been used and it is based on the amplification of

target DNA sequence Kang *et al.* [5]. *Pectobacterium* are necrotrophic plant pathogens responsible for inducing diseases like wilting, rotting, and blackleg in crucial agricultural crops, notably potatoes, carrots, tomatoes, onions, pineapples, corn, rice, hyacinths, chrysanthemums, and calla lilies. These infections lead to substantial reductions in crop yield, as documented in previous studies by Adeolu *et al.* [6] ; Charkowski [7] ; Charkowski *et al.* [8] ; Perombelon and Kelman [9]. Soft rot of carrot caused by *Pectobacterium* sp was previously reported by Michalik *et al.* [10] ; Parthiban [11] ; Tang *et al.* [12] ; Wasendorf *et al.* [13] . Since, the current study is focused on isolation and characterization of soft rot pathogen *Pectobacterium carotovorum* subsp *carotovorum* infecting carrot through morphological and molecular level analysis.

## 2. MATERIALS AND METHODS

### 2.1 Sample collection and isolation of pathogen

Diseased carrot root samples from major carrot growing areas of Ooty and local markets of Coimbatore were collected based on symptoms. The disease severity of the soft rot was observed by using Percent disease index (PDI) and calculated by using this formula given by Rose [14] .

$$\text{Per cent Disease Index (PDI)} = \frac{\text{Sum of individual ratings} \times 100}{\text{Total number of plants/ leaves observed} \times \text{Maximum disease grade}}$$

Grading the root vegetable by using 0-9 scale TNAU [15] and assessing the disease severity.

**Chart 1. List of grade scale and its description**

| Grade scale | Description                                     |
|-------------|---|
| 0           | No infection                                    |
| 1           | Less than 1% lesion covering the root vegetable |
| 3           | 1-10%lesion covering the root vegetable         |
| 5           | 11-25% lesion covering the root vegetable       |
| 7           | 26-50% lesion covering the root vegetable       |
| 9           | Lesion covering more than 50% of root vegetable |

The infected portion were cut from the carrot root and macerated in sterile pestle and mortar using 1-2 ml sterile distilled water and kept for 15 mins for oozing of bacterial cells. Then the bacterial suspension was serially diluted upto  $10^{-6}$  dilution and plated on Nutrient Agar medium using pour plate method in order to get uniform colonies. The single colonies were picked and streaked on NA medium and incubated at  $28^{\circ}\text{C}$  for 48 hours. The isolates were also streaked on CVP (Crystal violet pectate) a selective medium for *Erwinia* species bacteria which formed cavities or deep pits Cupples and Kelman [16]. The isolates were named based on location and serially numbered as GPB-1, MPB-2, UDB-3, PPB-4, OCB-5, GMB-6, KPB-7, GKB-8, OTB-9, ITB-10, SMB-11 and PDB-12.

## 2.2 Morphological characterization

The isolates of soft rot pathogen were characterized using various biochemical assays. The size, shape and arrangement of bacterial cells were identified through Gram staining Schaad *et al.* [17]. Other biochemical tests such as Methyl red test Mc Devitt [18], Catalase test Hayward [19], Gelatin hydrolysis test Clarke [20], KOH test Schaad *et al.* [17], H<sub>2</sub>S gas production Sendilvel *et al.* [21] and Potato soft rot test Muturi *et al.* [22] were also performed.

## 2.3 Pathogenicity tests

The pathogenicity of soft rot pathogen was confirmed in two different ways as furnished below. In the first method, surface disinfected healthy carrots were cut into slices of 5mm thickness and placed in petri plates. A volume of 150-200  $\mu\text{l}$  bacterial suspension ( $1 \times 10^8$  cfu/ml) was inoculated by injecting them onto the slices Dadasogolu and Kotan [24]. In the second method, 200-300  $\mu\text{l}$  of bacterial suspension ( $1 \times 10^8$  cfu/ml) was injected on disinfected healthy whole carrots using sterile syringe, later covered with wet cotton and placed in polythene bags Chandrashekar *et al.* [25]. Then the inoculated carrots were incubated for 3-4 days at  $28^{\circ}\text{C}$  for symptom expression and sterile distilled water is served as control in both the methods.

## 2.4 Molecular characterization

Bacterial DNA was isolated from the virulent isolates using lysis method Chen *et al.* [26] and they were amplified using 16s rRNA gene of bacterial universal primers 27F (5'-AGAGTTTGATCCTGGCTCAG-3') and 1492R (5'-GGTTACCTTGTTACGACTT-3'). PCR reaction was carried out using 10  $\mu\text{l}$  reaction mixture containing 1  $\mu\text{l}$  template DNA, 5  $\mu\text{l}$  Smart Prime 2X PCR Master Mix, 1  $\mu\text{l}$  forward primer, 1  $\mu\text{l}$  reverse primer and 2  $\mu\text{l}$  sterile water. DNA Amplification parameters were fixed as follows: initial denaturation  $95^{\circ}\text{C}$  for 3 min followed by 35 cycles of denaturation at  $95^{\circ}\text{C}$  for 30 sec, annealing at  $55^{\circ}\text{C}$  for 30 sec, initial extension at  $72^{\circ}\text{C}$  for 1 min and final extension at  $72^{\circ}\text{C}$  for 5 min. The amplified DNA was quantified through 1.2% agarose gel electrophoresis along with 1 Kb ladder. Resolved gel was documented in gel documentation unit (Bio rad) and the PCR products were partially sequenced and submitted in NCBI GenBank Lazaro *et al.* [27].

## 2.5 Statistical Analysis

Experimental data were analyzed statistically using Analysis of Variance (ANOVA) and the mean difference of all the treatment in Duncan's Multiple Range Test at 5% level of significance Gomez and Gomez [28]. All the data were analyzed using SPSS software (version 16) and interpreted.

## 3. RESULTS AND DISCUSSION

### 3.1 Sample collection and Isolation of the Pathogen

The diseased samples of carrot were collected from carrot growing areas of Ooty and from different local markets in Coimbatore, Erode and Dindigul districts. Percent disease index was calculated for the randomly selected carrots (100) based on the symptoms observed and characterized using disease score chart (0 to 9). Maximum disease incidence of soft rot incidence (66.74%) was noticed in Kettipalada of The Nilgiris district followed by Ottanchatrum of Dindigul district (45.55%) and the least disease incidence of 16.21 per cent was recorded in Perundurai of Erode district (Table 1). Carrot showing water soaked lesions with depressed and discoloured symptoms were isolated and the colonies with creamy white, slimy appearance were streaked in Nutrient Agar (NA) Medium. Similar type of results were obtained by Rahman *et al.* [29] observed the symptoms started with water soaked lesions later developing soft, watery and decay. Snehalatharani and Khan [30] reported that bacterial colonies with cream to white raised colonies with mucoid. Muturi *et al.* [22] isolated *Pectobacterium* from infected potato tubers and observed creamy, white colonies with mucoid.

### 3.2 Morphological characterization

Colony morphology studies of the bacterial soft rot pathogen revealed that the bacteria produced raised, mucoid, cream to white coloured colonies (Table 2, Fig 1). Furthermore, biochemical characterization revealed that the bacterial isolates OCB-5 and KPB-7 recorded a positive reaction for Methyl red, Growth at 36-37°C, H<sub>2</sub>S gas production, Gelatin liquefaction, Catalase test, KOH test and Potato soft rot test and negative for gram's reaction (Table 3). Gram staining indicate that pink colour bacterial cells results shows that the isolates were gram negative. Maximum growth of bacteria was recorded after 48 hrs incubation at 37°C in the isolate KPB-7 and MPB-2 (OD value @ 620 nm 1.733 and 1.627) respectively. De Boer and Kelman,2001 also reported that *Pectobacterium carotovorum* can grow at 37°C. Potato soft rot test showing softening of the tissue with rotting symptom Muturi *et al.* [22]. From the antibiotic resistance tests, inhibition zone was observed around the paper disc containing streptomycin and no such zone in paper disc containing erythromycin and found that the isolates were resistant to erythromycin and susceptible to streptomycin antibiotic. Akbar *et al.* [31] reported that *Pectobacterium carotovorum* isolates resistant to erythromycin and susceptible to streptomycin. The results of biochemical characterization are in agreement with similar results of Rahman *et al.* [29] and Ragavi [32]. In addition, Muturi *et al.* [22] found that the pathogen showed pectolytic activity to degrade the plant cell wall.

### 3.3 Pathogenicity test

Pathogenicity test were established for all the 12 isolates and the carrot inoculated artificially with each isolates with three replications. The pathogen able to produce symptoms was reisolated and proving the Koch's postulates. Intensity of disease was calculated by using Percent Disease Index (PDI) under disease grade scale (0 to 9). All the tested 12 isolates showed typical rotting symptoms within 1-3 days. The isolates, OCB-5 and KPB-7 were found to be the virulent isolates which showed severe infection of > 50% rotting under grade scale 9 (Table 4, Fig. 2). Tang *et al.* [12] also reported that the artificially inoculated carrot showing soft rot symptoms. Chandrashekar *et al.* [25] reported that bacterial suspension artificially inoculated into whole carrot showing water soaked lesions after 24hrs and complete rotting after 72hrs of incubation. In carrot slices, after 24hrs of incubation showing water soaked lesions and extending complete rotting.

### 3.4 Molecular characterization

The DNA extracted from the virulent bacterial isolates (KPB-7) produced DNA fragments corresponding to the 16S region of the rRNA gene when subjected to PCR amplification with 16S rRNA universal primers. From the gel electrophoresis, it is evident that the isolates produced DNA fragments at the amplicon size of 1500 bp (Fig. 3). The partial sequences of the isolates (KPB-7) were obtained and they were submitted in NCBI GeneBank with an accession number (OR251119). The isolate is identified as *Pectobacterium carotovorum* subsp *carotovorum* through NCBI BLAST search which showed 99% identity with other isolates of *P. carotovorum* subsp *carotovorum* in NCBI Database. Caruso *et al.* [33] reported that the isolates from tomato belongs to *Pectobacterium carotovorum* subsp *carotovorum* and *P.carotovorum* subsp *brasiliensis*. In a similar study , Muturi *et al.*,2018 stated that the analysis of 16S rRNA gene sequence revealed that the strain KPM17 was *Pectobacterium carotovorum* and 98% identity with *P.carotovorum* strain cc303. Ragavi [32] reported that rhizome rot of banana was *Pectobacterium carotovorum* subsp *carotovorum* had 99% similarity with existing isolates. Wasendorf [13] reported that the analysis of 16S region of the rRNA gene sequence was *Pectobacterium* strains isolated from carrot.

Table 1 : Collection of isolates from different district of Tamil Nadu infecting carrot

| S.No | Isolate No | Location             | District   | Latitude ( <sup>0</sup> N)-Longitude( <sup>0</sup> E) | Percent Disease Index (PDI)*<br>Soft rot |
|------|------------|----------------------|------------|---|--|
| 1    | GPB-1      | Gandhipuram          | Coimbatore | "11.01-76.96"   | 42.57 <sup>c</sup><br>(40.70)            |
| 2    | MPB-2      | Mettupalayam         |            | "11.30-76.93"   | 38.27 <sup>bc</sup><br>(38.14)           |
| 3    | UDB-3      | Ukkadam              |            | "10.99-76.96"   | 27.52 <sup>abc</sup><br>(31.35)          |
| 4    | PPB-4      | Periyanaickenpalayam |            | "11.14-76.94"   | 18.27 <sup>ab</sup><br>(24.83)           |
| 5    | OCB-5      | Ottanchatrum         | Dindigul   | "10.36-77.96"   | 45.55 <sup>cd</sup><br>(42.41)           |
| 6    | GMB-6      | Gandhi Market        |            | "10.48- 77.75"  | 34.77 <sup>abc</sup>                     |

|    |        |                |              |               |                                 |
|----|--------|----------------|--------------|---------------|---------------------------------|
|    |        |                |              |               | (35.96)                         |
| 7  | KPB-7  | Kethipalada    | The Nilgiris | "11.35-76.73" | 66.74 <sup>d</sup><br>(55.57)   |
| 8  | GKB-8  | Gandhikandi    |              | "11.30-76.62" | 40.23 <sup>c</sup><br>(39.12)   |
| 9  | OTB-9  | Ooty           |              | "11.41-76.69" | 36.64 <sup>abc</sup><br>(37.13) |
| 10 | IRB-10 | Ithalar        |              | "11.30-76.65" | 24.47 <sup>abc</sup><br>(29.16) |
| 11 | SMB-11 | Sathyamangalam | Erode        | "11.50-77.23" | 41.63 <sup>c</sup><br>(39.94)   |
| 12 | PDB-12 | Perundurai     |              | "11.27-77.58" | 16.21 <sup>a</sup><br>(23.66)   |

\*Mean of three replications and the mean followed by a common letter as superscript does not differ significantly at 5% level by DMRT. Values in parentheses are arc sine transformed value

**Table 2: Cultural characteristics of different isolates of soft rot pathogen associated with carrot**

| Isolate No                     | Colony colour   | Appearance             |
|--------------------------------|-----------------|------------------------|
| GPB-1, UDB-3, GMB-6 and IRB-10 | White           | Slimy                  |
| MPB-2, OCB-5, GKB-8, IPD-12    | Creamy white    | Slimy                  |
| PPB-4, KPB-7                   | Yellowish white | Slimy, Raised colonies |
| ISM-11                         | Yellowish white | Slimy                  |
| OTB-9                          | Creamy white    | Slimy, Raised colonies |

**Table 3 : Biochemical characterization of different isolates of soft rot pathogen associated with carrot**

| S.No | Biochemical test              | GPB-1 | MPB-2 | UDB-3 | PPB-4 | OCB-5 | GMB-6 | KPB-7 | GKB-8 | OTB-9 | IRB-10 | SMB-11 | PDB-12 |
|------|-------------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|--------|--------|--------|
| 1.   | Gram staining                 | -     | -     | -     | -     | -     | -     | -     | -     | -     | -      | -      | -      |
| 2.   | H <sub>2</sub> S Production   | +     | -     | +     | -     | +     | -     | +     | -     | -     | +      | -      | -      |
| 3.   | Catalase test                 | -     | +     | -     | +     | +     | +     | +     | -     | +     | -      | +      | -      |
| 4.   | KOH test                      | -     | +     | +     | -     | +     | +     | +     | -     | +     | -      | -      | +      |
| 5.   | Gelatin hydrolysis            | -     | -     | -     | +     | +     | -     | +     | +     | -     | +      | -      | +      |
| 6.   | Growth at 36-37°C             | +     | +     | +     | +     | +     | +     | +     | +     | +     | +      | +      | +      |
| 7.   | Potato soft rot test          | +     | +     | +     | -     | +     | +     | +     | -     | +     | +      | +      | -      |
| 8.   | Pits formation in CVP Medium  | +     | +     | -     | -     | +     | -     | +     | +     | -     | -      | -      | -      |
| 9.   | Methyl red test               | -     | +     | -     | -     | +     | +     | +     | -     | +     | +      | -      | +      |
| 10.  | Erythromycin sensitivity test | -     | -     | -     | +     | +     | -     | +     | +     | +     | +      | -      | -      |

+: Positive reaction  
 -: Negative reaction

**Table 4: Severity level of different isolates of Carrot soft rot pathogen associated with carrot**

| Score<br>chart/<br>Isolates | GPB-1 | MPB-2  | UDB-3 | PPB-4  | OCB-5  | GMB-6 | KPB-7 | GKB-8 | OTB-9 | IRB-10 | SMB-11 | PDB-12 | Severity                                    |
|-----------------------------|-------|--------|-------|--------|--------|-------|-------|-------|-------|--------|--------|--------|---|
| 0                           | -     | -      | -     | -      | -      | -     | -     | -     | -     | -      | -      | -      | No infection                                |
| 1                           | <1%   | -      | -     | -      | -      | -     | -     | -     | -     | <1%    | <1%    | <1%    | Water soaked lesion                         |
| 3                           | -     | -      | 1-10% | -      | -      | -     | -     | 1-10% | 1-10% | -      | -      | -      | Water soaked lesion<br>with initial rotting |
| 5                           | -     | 11-25% | -     | 11-25% | -      | -     | -     | -     | -     | -      | -      | -      | 25% rotting of the<br>vegetable             |
| 7                           | -     | -      | -     | -      | 26-50% | -     | -     | -     | -     | -      | -      | -      | 50% rotting of the<br>vegetable             |
| 9                           | -     | -      | -     | -      | -      | -     | >50%  | -     | -     | -      | -      | -      | Complete rotting of<br>the vegetable        |

**Fig 1. Pure cultures of different isolates of soft rot pathogen associated with carrot**




GPB-1




MPB-2



UDB-3



PPB-4



OCB-5



GMB-6



KPB-7



GKB-8




OTB-9



ITB-10



SMB-11



PDB-12

**Fig 2: Pathogenicity test**


Soft rot pathogen : *Pectobacterium carotovorum* subsp *carotovorum*



A- Control , B- Inoculated carrot showing soft rot symptoms

**Fig 3 . Molecular characterization of *Pectobacterium carotovorum* subsp *carotovorum***

**(KPB-7 Isolate)**



- L1 : 1 Kb Ladder
- L2 : GPB 1
- L3 : OCB 5
- L4 : KPB 7
- L5 : Control

#### **4. CONCLUSION**

In this current study, soft rot pathogen infecting carrot were isolated and they were identified through morphological and molecular characterization. Highest Soft rot incidence (66.74%) was observed in samples collected from Ooty area of The Nilgiris district followed by Ottanchatrum of Dindigul district with 45.55% incidence and the least disease incidence of (16.21%) was observed in Perundurai of Erode district. All the isolates of soft rot pathogen expressed pathogenic nature calculated based on the Percent Disease Index (PDI) wherein OCB-5 and KPB-7 were found to be highly virulent by using

disease grade chart (0 to 9) showing more than 50% rotting under disease grade 9. Biochemical characterization of 12 isolates revealed that the isolate OCB-5 and KPB-7 were positive for Catalase test, Methyl red, H<sub>2</sub>S gas production, Gelatin liquefaction, KOH test, Growth at 36-37°C, and Potato soft rot test and negative for gram's reaction. The soft rot bacterial isolate (KPB-7) was identified as *Pectobacterium carotovorum* subsp *carotovorum* with an accession number (OR251119) showing 99% identity with other *Pectobacterium* isolates in NCBI Database. Earlier detecting of the pathogen through morpho-molecular characterization helps in effective management strategies.

UNDER PEER REVIEW

## REFERENCES

1. Kitinoja, L., Tokala, V. Y., & Brondy, A. Challenges and opportunities for improved postharvest loss measurements in plant-based food crops. *Journal of Postharvest Technology*, 2018;6(4), 16-34.
2. Bhat, K. A., Masood, S. D., Bhat, N. A., Bhat, M. A., Razvi, S. M., Mir, M. R., Akhtar, S., Wani, N. and Habib, M. Current status of postharvest soft rot in vegetables: a review. *Asian Journal of Plant Sciences*, 2010;9(4), 200-208.
3. Whitehead, N. A., Byers, J. T., Commander, P., Corbett, M. J., Coulthurst, S. J., Everson, L., Harris, A.K., Pemberton, C.L. The regulation of virulence in phytopathogenic *Erwinia* species: quorum sensing, antibiotics and ecological considerations. *Antonie Van Leeuwenhoek*, 2002; 81, 223-231.
4. De Boer, S. H. Gram-negative bacteria, B-2 *Erwinia* soft rot group. Laboratory guide for identification of plant pathogenic bacteria, 2001;56-72.
5. Kang, H.W., Kwon, S. W., & Go, S. J. PCR-based specific and sensitive detection of *Pectobacterium carotovorum* ssp. *carotovorum* by primers generated from a URP-PCR fingerprinting-derived polymorphic band. *Plant pathology*, 2003; 52(2), 127-133.
6. Adeolu M, Alnajjar S, Naushad S, Gupta RS. Genome-based phylogeny and taxonomy of the 'Enterobacteriales': proposal for *Enterobacterales* ord. nov. divided into the families *Enterobacteriaceae*, *Erwiniaceae* fam. nov., *Pectobacteriaceae* fam. nov., *Yersiniaceae* fam. nov., *Hafniaceae* fam. nov., *Morganellaceae* fam. nov., and *Budviciaceae* fam. nov. *Int. J. Syst. Evol. Microbiol.* 2016;66: 5575–5599.
7. Charkowski AO. The changing face of bacterial soft-rot diseases. *Annu. Rev. Phytopathol.* 2018;56: 269–288.
8. Charkowski A, Blanco C, Condemine G, Expert D, Franza T, Hayes C, Hugouvieux-Cotte-Pattat N, López Solanilla E, Low D, Moleleki L, Pirhonen M, Pitman A, Perna N, Reverchon S, Rodríguez Palenzuela P, San Francisco M, Toth I, Tsuyumu S, van der Waals J, van der Wolf J, van Gijsegem F, Yang C-H, Yedidia I. The role of secretion systems and small molecules in soft-rot *Enterobacteriaceae* pathogenicity. *Annu. Rev. Phytopathol.* 2012;50:425–449.
9. Perombelon MCM, Kelman A. Ecology of the soft rot *Erwinias*. *Annu. Rev. Phytopathol.* 1980;18:361–387.
10. Michalik B, Simon PW, Gabelman WH. Assessing susceptibility of carrot roots to bacterial soft rot. *HortScience*. 1992 Sep 1;27(9):1020-1022.
11. Parthiban, V. K. Studies on the bacterial soft rot of carrot caused by *Erwinia carotovora* var. *carotovora* (Jones) Ph.D. Thesis. Tamil Nadu Agricultural University, Coimbatore. 2004. p. 114
12. Tang WQ, Chang CY, Lee YJ, Chu CC. First report of *Pectobacterium aroidearum* causing bacterial soft rot of carrot in Taiwan. *Plant Disease*. 2021 Mar 15;105(3):695.
13. Wasendorf, C., Schultz, D. L., Schmitz-Esser, S., & Peters, N. T. Genome sequences of soft rot-causing *Pectobacterium* isolates from different vegetables. *Microbiology Resource Announcements*, 2022;11(1):01066-2
14. Rose DH. Diseases of apple fruits in the market. *Bulletin, United States Department of Agriculture*. 1974; 1253:23- 24.
15. TNAU, Score chart for crop diseases. Tamil Nadu Agricultural University, Coimbatore. 1980.
16. Cuppels, D., & Kelman, A. Evaluation of selective media for isolation of soft-rot bacteria from soil and plant tissue. *Phytopathology*, 1974;64(4), 468-475.
17. Schaad, N. W., Jones, J. B., & Chun, W. Laboratory guide for the identification of plant pathogenic bacteria (No. Ed. 3). American Phytopathological society (APS press). 2001.
18. McDevitt, S. Methyl red and Voges-Proskauer test protocols. American Society for Microbiology, 2009, 8.
19. Hayward, A. C. Identification of *Pseudomonas solanacearum*. In: SAVERNET Bacterial wilt Training Course held on October 5 to November 16. AVRDC, Taiwan. 1992.p. 101.
20. Clarke, P. H., & Cowan, S.T. Biochemical methods for bacteriology. *Microbiology*, 1952; 6(1-2), 187-197.

21. Sendhilvel V, Marimuthu T, Raguchander T and Prabakar K. Biochemical methods for the detection of *Erwinia carotovora* var *carotovora* from Onion seeds, Madras Agric. J. 2005; 92(4-6): 234- 237.
22. Muturi, P., Yu, J., Li, J., Jiang, M., Maina, A. N., Kariuki, S., ... & Wei, H. Isolation and characterization of pectolytic bacterial pathogens infecting potatoes in Nakuru County, Kenya. Journal of applied microbiology, 2018;124(6), 1580-1588.
23. De Hoog, G. S., Smith, M. T., & Guého, E. *Galactomyces* Redhead & Malloch. In The Yeasts Elsevier. 1988; 209-213.
24. Dadasoglu.F and Kotan.R Identification and characterization of *Pectobacterium carotovorum* subsp *carotovorum* J.Animal and Plant Sciences 2017 ; 27(2): 647-654.
25. Chandrashekar, B. S., PrasannaKumar, M. K., Buela, P. P., Pramesh, D., Swathi, S. P., & Sahana, N. B. Association of *Acinetobacter baumannii* with soft rot disease of carrot in India. J Bacteriol Mycol, 2022;9(1), 1194.
26. Chen, W. P., & Kuo, T. T. A simple and rapid method for the preparation of gram-negative bacterial genomic DNA. Nucleic acids research, 1993;21(9), 2260.
27. Lázaro-Silva, D. N., De Mattos, J. C. P., Castro, H. C., Alves, G. G., & Amorim, L. M. F. The use of DNA extraction for molecular biology and biotechnology training: a practical and alternative approach. Creative Education, 2015;6(08), 762.
28. Gomez, K. A., & Gomez, A. A. Statistical procedures for agricultural research. John wiley & sons.1984;680.
29. Rahman, M. M., Ali, M. E., Khan, A. A., Hashim, U., Akanda, A. M., & Hakim, M. A. Characterization and identification of soft rot bacterial pathogens in Bangladeshi potatoes. African Journal of Microbiology Research, 2012;6(7), 1437-1445.
30. Snehalatharani, A., & Khan, A. N. A. Biochemical and physiological characterisation of *Erwinia* species causing tip-over disease of banana. Archives of Phytopathology and Plant Protection, 2010;43(11), 1072-1080.
31. Akbar, A., Ahmad, M., Khan, S. Z., & Ahmad, Z. Characterization of the causal organism of soft rot of tomatoes and other vegetables and evaluation of its most aggressive isolates. American Journal of Plant Sciences,2015; 6(04), 511.
32. Ragavi, G., Muthamilan.M, Nakkeeran.S, Kumaravadiel.N, Sivakumar.U and Suganthi, A. Molecular Detection of the Causative Agent of Soft Rot (*Pectobacterium carotovorum* subsp *carotovorum*) in banana (*Musa* sp.). Int.J.Curr.Microbiol.App.Sci. 2019; 8(11): 1854-1868.
33. Caruso, A., Licciardello, G., La Rosa, R., Catara, V., & Bella, P. Mixed infection of *Pectobacterium carotovorum* subsp. *carotovorum* and *P. carotovorum* subsp. *brasiliensis* in tomato stem rot in Italy. Journal of Plant Pathology,2016; 98(3), 661-665.