

Response of China aster to bio-inoculant amendments under pot experiment

A B S T R A C T

1. The present investigation was layout in Completely Randomized design (CRD) with 09 treatments and each treatment replicated thrice. The treatments consist of different combinations of bio-inoculants (Azospirillum, PSB, VAM and KSB). The treatment T₈ (75%RDF+Azospirillum+PSB) was found significantly higher compared to other treatment combination, which recorded highest plant height (27.28 cm), Number of leaves (27.44), plant spread (15.28 cm²), Days to bud emergence (47.11 days), days of first bud break (54.67 days), opening first flower(62.33 days), number of flowers per plant(14.12), stalk length (13.00 cm), flower diameter (4.92cm), Vase life (12.22 days), Leaf area (14.11 cm²). The economics viz. Gross return (Rs. 16,800), Net return (Rs. 8928) and Benefit cost ratio (2.13) was found highest in the same treatment.

Key words: China aster, Azospirillum, PSB, KSB, VAM.

Introduction

China aster (*Callistephus chinensis* (L.) Nees) belongs to family 'Asteraceae' and is native to China. The genus has only a single species, namely, *Callistephus chinensis*. The genus derives its name from two Greek words 'kalistos', most beautiful, and 'stephos', a crown, referring to the flower. Cassini described the China aster as *Callistephus hortensis*. It was first named by Linnaeus as *Aster chinensis*, and Nees subsequently changed this name to *Callistephus chinensis* (Janakiram, 2006). China aster plants range in height from 15 cm to about 1.0 m with pompon flowers about the size of a button to large flower heads having single, double, anemone-flowered, peony-flowered, incurved, quilled or shaggy flower types. The colour range is so great that today China aster is one of the most valuable garden flowers. In all the different types, the colours include pure white, many shades of pink, primrose, pale blue, mauve, purple, dark blue, and scarlet (Randhawa and Mukhopadhyay, 1986).

In view of China aster adapting well to varying soil and climatic conditions, it can be grown successfully under Indian climatic conditions. China aster flowers last longer which are used as cut flower, loose flower, bedding plant, for flower decoration, preparation of bouquets and garlands. In India, it is being grown in Karnataka, Tamil Nadu, Andhra Pradesh, Maharashtra and West Bengal.

Biofertilisers are defined as preparations containing living cells or latent cells of efficient strains of microorganisms that help

crop plants' uptake of nutrients by their interactions in the rhizosphere when applied through seed or soil. They accelerate certain microbial processes in the soil, which augment the extent of availability of nutrients in a form easily assimilated by plants. Very often microorganisms are not as efficient in natural surroundings as one would expect them to be and, therefore, artificially multiplied cultures of efficient selected microorganisms play a vital role in accelerating the microbial processes in the soil (Murugesan R and Prasad G, 2006).

China aster responds very well to the application of inorganic and organic fertilizers. Bio-fertilizer usually consists of live or latent cells of microorganisms which include biological nitrogen fixers, P-solubilizing, mineralization of nitrogen and transformation of several elements into available forms. VAM, Azotobacter, Azospirillum and phosphate solubilizing bacteria are commonly applied biofertilizers in horticultural crops.

Use of bio-fertilizers reduces per unit consumption of inorganic fertilizers and increases the quality and quantity of flowers.

Bioagents influences plant development by several mechanisms, such as production of growth hormones, solubilization of insoluble minor nutrients in soil and increased uptake and translocation of less available minerals.

Bioinoculants are the microbial inoculants which can be usually defined as a preparation containing live or dormant cells of efficient strains of nitrogen fixing, phosphate solubilizing, and cellulytic microorganisms, etc.

In contrast to chemical fertilizers, biofertilizers are viable microorganisms which are not the source of nutrients but provide help to plants in accessing the nutrient availability in rhizospheric region.

These microbial formulations are used to enhance certain microbial process to increase the availability of nutrients in a form which can be assimilated by plant. Biofertilizers are low-cost, renewable sources of plant nutrients.

This study aimed to evaluate the most suitable bio-inoculant for plant growth, yield and flower quality of China aster and estimate the economics of various treatments.

Materials and Methods

The present investigation was carried out at Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences (SHUATS), Prayagraj (U.P) in the month of November to March during the winter season of the year 2022.

The different treatment manipulated as follows T₁- 100%RDF, T₂- 75%RDF + Azospirillum, T₃- 75%RDF + KSB, T₄- 75%RDF + VAM, T₅- 75%RDF + PSB, T₆- 75%RDF + Azospirillum + KSB, T₇- 75%RDF + Azospirillum + VAM, T₈- 75%RDF + Azospirillum + PSB, T₉- 75%RDF + Azospirillum + KSB + VAM. The treatments were arranged in a Completely Randomized Design (CRD) with 9 treatments in 3 replications.

Results and Discussion

Growth attributes

Vegetative parameters viz., plant height, number of leaves per plant, plant spread and leaf area were recorded at different stages of plant growth from 20, 40 and 60 days after transplanting and the results from the observations made are as follows.

Plant height (cm)

Significant difference was observed due to different combination of bioinoculants for plant height, at 60 DAT. The Maximum Plant height at 60 days (27.28 cm) was recorded in the T₈ (75%RDF+ Azospirillum + PSB), followed by T₆ (75%RDF+ Azospirillum + KSB) with (25.67 cm) and the minimum Plant Height at 60 days (23.78 cm) was recorded in T₁(100%RDF).

Number of leaves per plant

Significant difference was observed due to different combination of bioinoculants for number of leaves per plant, at 60 DAT. The Maximum number of leaves per plant at 60 days (27.44) was recorded in the T₈ (75%RDF+ Azospirillum + PSB), followed by T₄ (75%RDF+ VAM) with (26.11) and the minimum number of leaves per plant at 60 days (23.00) was recorded in T₁ (100%RDF).

Plant spread (cm²)

Significant difference was observed due to different combination of bioinoculants for plant spread (cm²), at 60 DAT. The Maximum plant spread (cm²) at 60 days (15.28 cm²) was recorded in the T₈ (75%RDF+ Azospirillum + PSB), followed by T₉ (5%RDF+ Azospirillum +KSB +VAM) with (14.83 cm²) and the minimum plant spread (cm²) at 60 days (12.33 cm²) was recorded in T₁ (100%RDF).

The deviation in leaf area due to different combination of bioinoculants was significant. The maximum leaf area was in (14.11 cm²) T₈ (75% RDF + Azospirillum + PSB) which was statistically at par with the other treatments like (13.93 cm²) T₉ (75% RDF + Azospirillum +KSB+VAM). However, minimum leaf area (10.67 cm²) was recorded in T₁ (100% RDF).

Flowering attributes

Days taken for bud emergence

Significant difference was observed due to different combination of bioinoculants for days of bud emergence, at DAT. The minimum days of bud emergence (47.11 days) was recorded in the T₈ (75%RDF+ Azospirillum + PSB), followed by T₂

(75%RDF+Azospirillum) with (47.56 days) and the Maximum days of first bud emergence (55.44 days) was recorded in T₁ (100%RDF).

Days taken for bud break

Significant difference was observed due to different combination of bioinoculants for days taken to bud break, at DAT. The minimum days taken for bud break (54.67 days) was recorded in the T₈ (75%RDF+ Azospirillum + PSB), followed by T₂ (75%RDF+Azospirillum) with (55.00 days) and the Maximum days taken for bud break (61.11 days) was recorded in T₁ (100%RDF).

Chart1: Various treatments of bud emergence, bud break, Plant height and plant spread.

Treatment Symbol	Treatments combination	Plant height (cm)	Plant Spread (cm)	Leaf area per plant (cm ²)	Number of Leaves per plant	Days taken for bud emergence	Days taken for bud break	Days taken for opening first flower
T ₁	100%RDF	23.28	12.33	10.67	23.00	55.44	61.11	69.22
T ₂	75%RDF+ Azospirillum	25.22	13.67	11.88	24.44	47.56	55.00	64.00
T ₃	75%RDF+ PSB	24.28	13.56	13.72	23.67	51.44	58.44	66.67
T ₄	75%RDF+ VAM	25.06	12.33	12.73	26.11	48.44	55.67	65.33
T ₅	75%RDF+ KSB	24.89	14.03	12.28	25.33	53.89	60.44	68.33
T ₆	75%RDF+ Azospirillum + PSB	25.67	13.23	13.33	24.11	49.56	57.11	67.33
T ₇	75%RDF+ Azospirillum + VAM	24.67	13.57	13.32	25.44	51.33		

							56.11	65.56
T ₈	75%RDF+ Azospirillum + KSB	27.28	15.28	14.11	27.44	47.11	54.67	62.33
T ₉	75%RDF+ Azospirillum + PSB+VAM	25.56	14.83	13.93	26.00	48.44	55.44	63.00
S. Ed. (±)		1.427	0.792	1.205	2.359	3.592	3.81	3.731
C. D. at 5 %		N/A	1.677	N/A	N/A	N/A	N/A	N/A

Table.1 Response of bioinoculants on Growth, Flowering, yield and post-harvest attributes of china aster.

Treatment Symbol	Treatments combination	Stalk length (cm)	Flower diameter (cm)	Number of flowers per plant	Number of buds fully opened	Chlorophyll content	Vase life (days)
T ₁	100%RDF	11.3	3.89	11.56	11.11	40.98	11.11
T ₂	75%RDF+ Azospirillum	12.3	4.36	13.67	13.00	40.32	11.56
T ₃	75%RDF+ PSB	12.1	4.48	12.56	11.78	40.79	11.56
T ₄	75%RDF+ VAM	11.8	4.29	12.78	12.11	39.11	11.78
T ₅	75%RDF+ KSB	11.6	4.63	14.00	12.67	39.72	11.67
T ₆	75%RDF+ Azospirillum + PSB	11.7	4.07	12.89	11.67	43.43	11.56
T ₇	75%RDF+ Azospirillum + VAM	11.7	4.07	13.78	13.44	44.12	11.89
T ₈	75%RDF+ Azospirillum + KSB	13.0	4.92	14.56	14.00	46.99	12.22
T ₉	75%RDF+ Azospirillum + PSB+VAM	11.4	4.68	13.11	12.44	41.73	11.33
S. Ed. (±)		0.570	0.227	1.079	0.796	2.549	0.343
C. D. at 5 %		N/A	0.481	N/A	N/A	N/A	N/A

Flower stalk length (cm)

Significant difference was observed due to different combination of bioinoculants for flower stalk diameter, at DAT. The Maximum flower stalk length (13.00 cm) was recorded in the T₈ (75%RDF+ Azospirillum + PSB), followed by T₂ (75%RDF+Azospirillum) with (12.32 cm)

and the minimum flower stalk length (11.30 cm) was recorded in T₁ (100%RDF).

Flower diameter (cm)

Significant influence of different treatments on flower diameter was observed. (Table 1). Highest Flower diameter was registered in T₃ (7.80 cm) and it was closely followed by T₁

(7.50 cm), T₆ (7.40 cm) and T₉ (7.10 cm). The treatment T₁₀ (5.83) recorded lowest flower diameter.

Yield parameters

Number of flowers per plant

The treatments significantly influenced the number of flowers per plant. Among the treatments, the maximum number of flowers per plant (14.12) were observed in T₈ i.e. 75% RDF + Azospirillum + PSB with other treatments like T₅ (100% RDF + PSB) with 14.00 number of flowers per plant, respectively. The minimum number of flowers per plant (22.51) was observed in the treatment T₁ (100%RDF).

Number of buds fully opened

An inspection of the data presented in Table 1 reveals the significant results for number of buds fully opened. The maximum number of buds fully opened (14.00) were observed in T₈ i.e. 75% RDF + Azospirillum + PSB with other treatments like T₇ (75%RDF+ Azospirillum + VAM) with (13.44) number of buds fully opened, respectively. The minimum number of buds fully opened (11.11) was observed in the treatment T₁ (100%RDF).

Chlorophyll content

Significant difference was observed due to different combination of bioinoculants for chlorophyll content (days), at DAT. The Maximum chlorophyll content (12.22 days) was recorded in the T₈ having (75% + Azospirillum + PSB) and it was found at par with T₇ (75% RDF + Azospirillum + KSB+VAM), was (11.89 days) while, the

minimum chlorophyll content (3.89 cm) was noticed in T₁ (100%RDF).

Post-harvest attributes

Vase life of cut flower

Application of various dose of different combinations of biofertilizers produced significant effects on the vase life of China aster cut flowers (Table 1). Significant difference was observed due to different combination of bioinoculants for vase life (days), at DAT. The Maximum vase life (12.22 days) was recorded in the T₈ having (75% + Azospirillum + PSB) and it was found at par with T₇ (75% RDF + Azospirillum + KSB+VAM), was (11.89 days) while, the minimum vase life (3.89 cm) was noticed in T₁ (100%RDF).

Discussion

The possible reason for increase in plant height no. of leaves ,plant spread ,bud break,vase life and economics is that combined application of biofertilizers with 75% of recommended dose of RDF and resulted in better nutrition which leads to increased photosynthesis activity, enhanced cell division and enlargement as nitrogen is important constituent of nucleic acid and it might have increased the synthesis of carbohydrate, amino acids etc. from which the phytohormones like auxins, gibberellins, cytokines have been synthesized and phosphorus being an essential component of protoplasm and chlorophyll, caused conversion of photosynthates into phospholipids resulting in adequate vegetative growth thus increased plant height. Biofertilizers produce several growth

promoting hormones (auxins, cytokinins and gibberellins etc.) in addition to increasing the availability of nitrogen and phosphorus to the plants resulting in better plant growth. Similar results of increased plant height due to combined application of biofertilizers with reduced dose of NPK have been reported by Chaitra and Patil (2007), Patil and Agasimani (2013) and Kirar et al.

Conclusion

On the basis of present investigation, it is concluded that the treatments T₈ (75% + Azospirillum + PSB) was found best treatment combinations which were significantly increased growth, flowering and quality parameters of China aster. The treatments promoted the plant height, bud emergence and for vase life. Azospirillum and PSB resulting in enhanced nutrient uptake which led to increased growth, yield and quality of China aster flowers.

References

- Ahmad M S, Beigh M A, Nanda A B, Lone R A and Hussain K. 2007. Effect of Azospirillum, VAM and inorganic fertilizers on growth and yield of African marigold cv. 'Pusa Narangi'. *Journal of Plant Science Research* **23**(1/2): 51- 53.
- Almeida, E.F.A. (2013). Soil Sustainable management in rose integrated production. *Acta Horticulturae*, 970: 355-360.
- Bhaskaran P, Ambrose G and Jayabalan N. 2002. Usefulness of biofertilizers ineconomising nitrogenous fertilizers in *Tagetes erecta* L. *Journal of Phytological Research* **15**(2): 155-160.
- Bose B. Subash and Chanda, (2019). Effect of integrated nutrient management on yield and quality of China aster (*Callistephus chinensis* (L.) Nees) cv. Pit and Pot. *Plant Archives* **19** (1): 518- 520.
- Dharma S. 2006. Effect of biofertilizers and biostimulants on growth and flowering of carnation (*Dianthus caryophyllus* L.) cv. 'Sunrise'. Ph. D. Thesis submitted to Dr. Yashwant Singh Parmar University of Horticulture and Forestry, Nauni, Solan, Himachal Pradesh, India.
- Johnson, CR., J.A. Menge and E.LV. Johnson, 1982. Effect of vesicular-arbuscular mycorrhizae on growth of *Chrysanthemum morifolium* Ramat. *Scientia Hort.*, **17**:265-9.
- Karuppaiah P. 2005. Effect of Azospirillum, phosphobacteria and organic manures on growth and yield of French marigold (*Tagetes patula* L.). *Plant Archives* **5**(2): 661-664.
- Kumar P, Raghava S P S and Misra R L. 2003b. Effect of biofertilizers on growth and yield of China aster. *Journal of Ornamental Horticulture New Series* **6**(2): 85-88
- Kumar V and Singh A. 2007. Effect of vermicompost and VAM inoculation on vegetative growth and floral attributes in China aster (*Callistephus chinensis* (L.) Nees). *Journal of Ornamental Horticulture* **10**(3): 190-192

- Kumar, P S. (2017). Effect of biofertilizer and organic manures on plant growth, flowering and tuber production of dahlia (*Dahlia variabilis* L.) cv S.P. Kamala. *International Journal of Pure and Applied Bioscience* 5(2): 549- 555.
- Latt, T.T., Nyunt, T.T., War, N.M., Shwe, S.S. and Myint, S.S. 2017. Role of Trichoderma in nutrient management of China aster [*Callistephus chinensis* (L.) Nees] with special emphasis on nitrogen input. *Sustainability in Myanmar* (Southeast Asian modernities, band 15), first edition, lit verlag.
- Mogal S A, Khiratkar S D, Chopde N K, Dalvi A M, Kuchanwar O D and Khobragade Y R. (2006). Effect of organic manures and biofertilizers with reduced doses of nitrogen on growth, yield and quality of China aster. *Journal of Soils and Crops* 16(1): 180-185.
- Mostafa M M. 2002. Effect of biofertilizer, salinity and magnetic technique on the growth of some annual plants. *Alexandria Journal of Agricultural Research* 47(2): 151-162.
- Munikrishnappa P.M. (2014). Integrated nutrient management studies in China aster (*Callistephus chinensis* (L.) Nees) - A review. *Trends in Biosciences*. 7 (12): 1217-1221.
- Nandre D R, Jogdande N D, Dalal S R, Bansode A B and Chaudhale B S. (2005). Effect of Azotobacter on growth and yield of China aster under reduced nitrogen doses. *Crop Research Hisar* 29(2): 272-274.
- Patanwar, M. (2014). Growth and development of chrysanthemum (*Dendranthema grandiflora*) as influenced by integrated nutrient management. *The Ecoscan*: 6: 459-462.
- Patil V.S. Agasimani A D. (2013). Effect of Integrated nutrient management on growth and yield parameters in China aster (*Callistephus chinensis*). *Mysore Journal of Agricultural Sciences*, 47(2): 267-272.
- Prabhat, K. (2003). Effect of biofertilizers on growth and yield of China aster. *Journal of Ornamental Horticulture*, 6 (2): 85-88.
- Ravindra S P, Hanumanth Appa M, Hegde J N, Maheshwar K J and Nagesha L. (2013). Effect of integrated nutrient management on growth, yield and vase life of China aster (*Callistephus chinensis* (L.) Nees) for coastal Karnataka. *Environment and Ecology* 31(2C): 1104-1106.
- Srivastava Ranjan et al., 2007. Influence of biofertilizers on growth and flowering in Gladiolus cv. American Beauty. *Acta Horticulturae*. 742: 183-188.
- Syamal M M, Dixit S K and Kumar S. 2006. Effect of bio-fertilizers on growth and yield in marigold. *Journal of Ornamental Horticulture* 9(4): 304-305.

UNDER PEER REVIEW