

Influence of Zinc and Boron application on growth and yield of French Bean
(*Phaseolus vulgaris* L.)

ABSTRACT

A field experiment was conducted during *Rabi* 2022 at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj (U.P), India. To find out the influence of Zinc and Boron on the growth and yield attributes of French bean. The experiment was laid out in Randomized Block Design with ten treatments including control each replicated thrice based on one year of experimentation. The application of Zinc 45kg/ha⁻¹ + Boron 0.5% @ 15DAS, recorded significantly higher Plant height (44.00 cm), Plant dry weight (29.31 g/plant). Significantly maximum pods/plant (17.40), Seeds/pod (7.00), Seed index (44.04 g), Seed yield (1.37 t/ha), stover yield (3.21 t/ha) were recorded with the treatment of Zinc 45kg/ha⁻¹ + Boron 0.5% @ 15DAS.

Keywords: Boron, Economics, French Bean, Growth parameter, Zinc.

INTRODUCTION

Kidney beans are the young, immature fruits of various types of kidney beans (*Phaseolus vulgaris*), but the immature or young pods of pole beans (*Phaseolus coccineus*), fava beans (*Sesquipedal` s*), and hyacinth beans (*Lablab purpureus*) are also used. increase. Kidney beans are known by many common names, including broad bean (French: haricot vert), kidney bean (although most modern varieties are "stringless"), broad bean, or simply "snaps." (Singh 2015) In the Philippines, it is also called 'bagoio bean' or 'habithuela' to distinguish it from the 1-meter-long bean (Maribehlla 2019). Unlike many other types of beans, green beans are harvested and consumed in pods before the bean seeds within them are fully mature. A similar practice is to harvest and consume immature pea pods, as is done with sugar snap peas and field peas.

More than 130 varieties (cultivars) of edible legumes are known. In home gardens, there are many varieties of green beans that are grown specifically for green pods, chosen for their juiciness and flavor, and there are many of them. Beans with different pod colors (green, purple, red, striped, Singh BK 2011) are collectively known as snap beans, while green beans are predominantly green. Shapes range from the thin 'fillet' type to the wide 'Romano' type and the more common type in between. The three most common types of kidney beans (*Phaseolus vulgaris*) species are round and flat. Stringless beans or bush beans, lacking the tough fibrous threads running the length of the pod. Runner beans belong to another species, kidney beans. Green beans sometimes have purple pods instead

of green pods that turn green when cooked. Green beans with yellow pods are also called wax beans. Wax bean varieties usually have a bushy or dwarf form.

Zinc is an essential micronutrient required by all living organisms and plays several important roles in life, growth and development. It plays an important role in the normal physiological activities of growth and development (Stanton et al., 2022; Wu et al., 2022). Zinc is an essential trace metal that has a structural role in protein regulation as an enzymatic cofactor (Sobczyk and Gaunt, 2022) and is involved in several enzymes involved in numerous processes such as DNA replication, protein synthesis, and lipid metabolism. acts as a cofactor for In addition, zinc plays an important role in many plant metabolic processes, including enzymatic activity, photosynthesis, chlorophyll synthesis, and other biochemical functions (Grüngreiff et al., 2020). Zinc is the second most abundant trace element required by all living organisms. It occurs as a divalent cation (Zn^{2+}) and has no redox activity under physiological conditions. This explains the performance of Zn in different physiological roles in different biological activities. Due to the high zinc content, a single application significantly reduced the yield of bean roots. There are several examples showing that the application of zinc fertilizers or zinc-enriched NPK fertilizers to crops not only improves productivity, but also improves plant zinc concentrations in grains. Zinc is involved in the formation of auxin and activation of dehydrogenase enzymes. Stabilization of the ribosomal fraction. Zinc deficiency is now listed as a major risk factor for human health and mortality worldwide.

Boron (B) is a micronutrient essential for the growth and health of all crops. It is a component of plant cell walls and reproductive structures. It is a mobile nutrient in the soil, meaning it can easily move through the soil. Boron is one of the essential micronutrients required for normal growth of most plants. Boron is required for proper development and differentiation of tissues and also helps reduce infertility and malformations of the reproductive organs [Singh 2008]. Boron aids in normal plant growth, nitrogen uptake from soil, sugar mobilization, cell wall synthesis, root elongation, and nucleic acid synthesis. Boron improves grain and straw yield, nutrient content, nutrient intake and quality in legume crops. Boron deficiency limits the production of legumes [Mani 1998]. Given the importance of phosphorus and boron, this study was conducted with the aim of investigating the response of boron and its interaction in maximizing the growth and yield performance of French beans.

MATERIAL AND METHODS

The experiment conducted to know the Influence of Zinc and Boron application on the growth and yield of French Bean (*Phaseolus vulgaris* L.) was carried out at the Crop Research Farm of Sam Higginbottom University, Prayagraj, Uttar Pradesh in 2022. The soil was sandy loam in texture, medium in available nitrogen (238.12 kg/ha), low in Phosphorous (38.3 kg/ha⁻¹), and medium in potassium (244.8 kg/ha⁻¹). The experiment was laid out in an RBD consisting of ten treatments including Control with 3 replications, viz., The treatments which are T₁: Zinc 15kg/ha⁻¹ + Boron 0.5% @ 15DAS, T₂: Zinc 15kg/ha⁻¹ + Boron 0.6% @ 25DAS, T₃: Zinc 15kg/ha⁻¹ + Boron 0.7% @ 55DAS, T₄: Zinc 30kg/ha⁻¹ + Boron 0.5% @ 15DAS, T₅: Zinc 30kg/ha⁻¹ + Boron 0.6% @ 25DAS, T₆: Zinc 30kg/ha⁻¹ + Boron 0.7% @ 55DAS, T₇: Zinc 45kg/ha⁻¹ + Boron 0.5% @ 15DAS, T₈: Zinc 45kg/ha⁻¹ + Boron 0.6% @ 25DAS, T₉: Zinc 45kg/ha⁻¹ + Boron 0.7% @ 55DAS, T₁₀: Control are used. The French bean seeds were sown at a spacing of 35 cm x 10 cm with a seed rate of 50 – 75 kg/ha⁻¹, growth-like plant height and dry weight and yield-attributing characteristics, such as the number of seeds per pod, have considerably increased. The yield contributing characters such as the number of pods per plant, number of seeds per pod, seed yield, and stover yield were recorded at the time of harvest and averages were calculated and the data were statistically analyzed using the ANOVA technique in Randomized Block design.

RESULTS AND DISCUSSION

Influence of zinc and boron on the growth of French bean: The perusal of the data of growth attributes recorded at growth stage, is presented in Table 1.

Plant height: At 80 DAS, there was a significant difference among the treatments. However, the highest plant height (44.00 cm) was recorded with the application of Zinc 45kg/ha⁻¹ + Boron 0.5% @ 15DAS, whereas the minimum plant height (36.90 cm) was recorded with the treatment Control 120:60:50 (N:P:K) kg/ha⁻¹ and Zinc 30kg/ha⁻¹ + Boron 0.5% @ 15DAS (33.30 cm), Zinc 15kg/ha⁻¹ + Boron 0.5% @ 15 DAS (32.97 cm) were statistically at par with T₇. This might be due to the quick availability of boron to crop during the entire growing season. Boron plays an important role in tissue differentiation and carbohydrate metabolism. It is also a constituent of cell membrane and essential for cell division, and maintenance of conducting tissue with regulatory effect on other element. Similar results were shown by Alam *et al.* (2014).

Plant dry weight: highest dry weight (29.31 g) was recorded with the application of Zinc 45kg/ha⁻¹ + Boron 0.6% @ 15DAS, whereas the minimum plant dry weight (24.65 g) was recorded with the treatment Control 120:60:50 (N:P:K) kg/ha⁻¹ and Zinc 30kg/ha⁻¹ + Boron 0.5% @ 15DAS (29.05 g) was statistically at par with T₇. This might be due to the quick

availability of boron to crop during the entire growing season. Boron plays an important role in tissue differentiation and carbohydrate metabolism. It is also a constituent of the cell membrane and essential for cell division, and maintenance of conducting tissue with a regulatory effect on another element (Rahman *et al.*, 2013).

Influence of zinc and boron on yield attributes of French bean

Number of Pods/plants: The perusal of the data of the number of Pods/plants recorded at harvest, is presented in Table 2. The data reveals that there was a significant effect among different treatments on the Number of Pods/plants.

The maximum number of Pods/plant (17.40) was recorded with the treatment of the application of Zinc 45kg/ha⁻¹ + Boron 0.5 % @ 15DAS over all the treatments, and the minimum was recorded in Control 120:60:50 (N:P:K) kg/ha⁻¹ (10.00). However, the treatments Zinc 30kg/ha⁻¹ + Boron 0.5 % @ 15DAS (17.07), and Zinc 15kg/ha⁻¹ + Boron 0.5% @15DAS (16.80) which was found to be statistically at par with T7. This might be due to the role of zinc in the production of biomass and that iron is necessary for chlorophyll synthesis and has many essential roles in plant growth and development (Abdollahi *et al.*, 2010).

The number of Seeds/Pod: Significantly Maximum Number of seeds/pods (7.00) was recorded with the treatment of the application of Zinc 45kg/ha⁻¹ + Boron 0.5% @ 15DAS over all the treatments, and the minimum was recorded in Control 120:60:50 (N:P:K) kg/ha⁻¹ (4.07). However, the treatments Zinc 30kg/ha⁻¹ + Boron 0.5% @ 15DAS (6.00), and Zinc 15kg/ha⁻¹ + Boron 0.5% @ 15DAS (6.40) which was found to be statistically at par with T7. The improvement might be due to an increase in the germination percentage of seed inside the pod and may be due to boron making stigma receptive and sticky and making pollen grain fertile and enhancing the pollination. Thus, increased fruit setting reduces the sterility of the flowers and the number of grains per pod increases. Similar results were reported by Zaman *et al.* (1996).

Seed index (g): Highest seed index (44.04 g) was recorded with the treatment of the application of Zinc 45kg/ha⁻¹ + Boron 0.5% @ 15DAS over all the treatments and the minimum was recorded in Control 120:60:50 (N:P:K) kg/ha⁻¹ (33.90 g). However, the treatments Zinc 30kg/ha⁻¹ + Boron 0.5% @ 15DAS (43.85 g), Zinc 15kg/ha + Boron 0.5% @ 15DAS (42.56 g) which was found to be statistically at par with T7. This improvement in the test weight of green gram may be due to boron, which affects cell division, carbohydrate metabolism, sugar and starch formation, which increased the size and weight of the grain. Similar results were reported by Padbhushan and Kumar (2014).

Seed yield (t/ha): Significantly Maximum seed yield (1.37 t/ha) was recorded with the treatment of the application of Zinc 45kg/ha + Boron0.5% @15DAS over all the treatments, and the minimum was recorded in Control 120:60:50 (N:P:K) kg/ha (1.22 t/ha). However, the treatments Zinc 30kg/ha + Boron0.5% @15DAS (1.34 t/ha), Zinc 15kg/ha + Boron0.5% @15DAS (1.33 t/ha) which was found to be statistically at par with T7. Yield enhancement meets plant needs through increased assimilation and translocation of photosynthesis from leaves to seeds, increased boron to the process of somatic to reproductive tissue differentiation, meristematic activity and flower development. may be due to the B application associated with The original strain gained weight and produced more flowers and a higher seed yield. Similar results were reported by Kaiser et al. report. (2010).

Stover yield (t/ha): Significantly Maximum stover yield (3.21 t/ha) was recorded with the treatment of the application of Zinc 45kg/ha⁻¹ + Boron0.5% @15DAS over all the treatments, and the minimum was recorded in Control 120:60:50 (N:P:K) kg/ha⁻¹ (2.95 t/ha). However, the treatments Zinc 30kg/ha⁻¹ + Boron0.5% @15DAS (3.20 t/ha⁻¹), Zinc 15kg/ha⁻¹ + Boron0.5% @15DAS (3.19 t/ha⁻¹) which was found to be statistically at par with T7.

CONCLUSION

It is concluded that the application of Zinc 45kg/ha⁻¹ + Boron 0.5% at 15DAS recorded higher seed yields as compared to other treatments and was found suitable for obtaining higher yield in French beans.

REFERENCES

- Alam Md., Sarowar and Islam Md., Faridul (2016) Effect of zinc and boron on seed yield and yield contributing traits of mungbean in acidic soil Journal of Bioscience and Agriculture Research 11 941-946.
- Abdollahi M, Eshghi S, Tafazoli E. Interaction of Paclobutrazol, Boron and Zinc on Vegetative Growth, Yield and Fruit Quality of Strawberry (*Fragaria* × *Ananassa* Duch. cv. Selva). Journal Biology Environmental Science. 2010; 4:67-75.
- "Baguio Beans". Maribehlla. January 25, 2011. Retrieved October 20, 2019.
- Grüngreiff, K., Gottstein, T., and Reinhold, D. (2020). Zinc deficiency—An independent risk factor in the pathogenesis of haemorrhagic stroke? *Nutrients*. 12:3548. doi: 10.3390/nu12113548.
- Kaiser, M.S., Rahman, MA., Amin, MHA., Amanullah, ASM and Ashanullah, ASM (2010) Effect of sulphur and boron on the seed yield and protein content of mungbean Bangladesh Research Publication 3,1181-1136.

- Mani PK, Haldar M. Effect of dolomite on boron transformation in acid soil in relation to nutrition of green gram. *Journal of Indian Society of Soil Science*.1996;44:458-461.
- Padbhushan, Rajeev and Kumar, Dinesh (2014) Influence of soil and foliar applied boron on green gram in calcareous soils. *International Journal of Agriculture, Environment and Biotechnology* 7, 129-136.
- Rahman, Inayat Ur, Afzal, Aftab, Zafar, Iqbal, Farhana, Ijaz, Sohail, Salma, Shad, Shafiul, Manan and Muhammad, Afzal. (2013) Response of common bean (*Phaseolus vulgaris*) to basal applied and foliar feeding of different nutrients application. *American-Eurasian J. Agric. and Environmental. Science* 14 (9): 851-854.
- Singh RN, Singh S, Kumar B. Interaction effect of sulphur and boron on yield and nutrient uptake and quality characters of soybean (*Glycine max L. Merrill*) grown in acidic upland soil. *Journal of the Indian Society of Soil Science*. 2006;54(4):516- 518.
- Singh BK and Singh B. 2015. Breeding perspectives of snap bean (*Phaseolus vulgaris* L.). *Vegetable Science* 42(1): 1-17.
- Sobczyk, M. K., and Gaunt, T. R. (2022). The effect of circulating zinc, selenium, copper and vitamin K1 on COVID-19 outcomes: a Mendelian randomization study. *Nutrients*.14:233. doi: 10.3390/nu14020233.
- Singh B K, Pathak K A, Ramakrishna Y, Verma V K and Deka B C. 2011. "Purple-podded French bean with high antioxidant content." *ICAR News: A Science and Technology Newsletter* 17 (3): 9.
- Stanton, C., Sanders, D., Krämer, U., and Podar, D. (2022). Zinc in plants: integrating homeostasis and biofortification. *Mol. Plant*. 15, 65–85. doi: 10.1016/j.molp.2021.12.008.
- Zaman, A.K.M.M., Alam, M.S., Roy, B. and Beg, A.H. (1996) Effect of B and Mo application on mungbean. *Bangladesh Journal of Agricultural. Research* 21, 118-124.
- Wu, Q., Liu, C., Wang, Z., Gao, T., Liu, Y., Xia, Y., *et al.* (2022). Zinc regulation of iron uptake and translocation in rice (*Oryza sativa* L.): Implication from stable iron isotopes and transporter genes. *Environ. Pollut.* 297:118818. doi: 10.1016/j.envpol.2022.118818.

Table 1: influence of Zinc and Boron on growth parameters of French bean.

Treatments	Plant height (cm)	Plant Dry Weight (g/plant)
Zinc 15kg/ha + Boron0.5%at15DAS	42.37	27.89
Zinc 15kg/ha + Boron0.6%at25DAS	38.40	27.19
Zinc 15kg/ha + Boron0.7%at55DAS	37.40	26.89
Zinc 30kg/ha + Boron0.5%at15DAS	43.40	29.05
Zinc 30kg/ha + Boron0.6%at25DAS	38.40	27.52
Zinc 30kg/ha + Boron0.7%at55DAS	37.45	27.01
Zinc 45kg/ha + Boron0.5%at15DAS	44.00	29.31
Zinc 45kg/ha + Boron0.6%at25DAS	40.80	27.80
Zinc 45kg/ha + Boron0.7%at55DAS	38.03	27.10
Control (RDF) 120:60:50 Kg NPK/ha	36.90	24.65
Sem (\pm)	0.41	0.23
CD(p=0.05)	1.24	0.69

Table 2: influence of Zinc and Boron on yield parameters of French bean.

Treatments	Pods/plant (No)	Seeds/pod (No)	Seed index (g)	Seed yield (t/ha)	Stover yield (t/ha)
Zinc 15kg/ha + Boron0.5%at15DAS	16.80	6.40	42.56	1.32	3.19
Zinc 15kg/ha + Boron0.6%at25DAS	12.00	5.07	38.97	1.29	3.16
Zinc 15kg/ha + Boron0.7%at55DAS	11.00	3.93	35.78	1.26	3.05
Zinc 30kg/ha + Boron0.5%at15DAS	17.07	6.60	43.85	1.33	3.20
Zinc 30kg/ha + Boron0.6%at25DAS	14.00	4.93	40.78	1.31	3.17
Zinc 30kg/ha + Boron0.7%at55DAS	10.00	3.93	36.99	1.27	3.08
Zinc 45kg/ha + Boron0.5%at15DAS	17.40	7.00	44.04	1.37	3.21
Zinc 45kg/ha + Boron0.6%at25DAS	16.00	6.00	41.18	1.31	3.18
Zinc 45kg/ha + Boron0.7%at55DAS	11.00	3.93	38.25	1.28	3.09
Control (RDF) 120:60:50 Kg NPK/ha	10.00	4.07	33.90	1.22	2.95
Sem (\pm)	0.50	0.13	0.68	0.01	0.03
CD(p=0.05)	1.49	0.39	2.03	0.03	0.02