

Effect of pH on the growth of *Aspergillus niger* and *Aspergillus flavus*

ABSTARCT

Aspergillus fungus is well-known for causing a number of secondary plant and food rots that can result in the buildup of mycotoxins, despite their significant economic contribution to the fermentation industries. In the past, even the most fundamental influences on the development and inhibition of significant mycotoxigenic fungi were only partially understood. To overcome this issue, an investigation on the impact of pH on the mycelial development of *Aspergillus* species was conducted. Results revealed that, the mycelial growth of *A. ~~aspergillus~~niger* and *A. flavus* was increased with increase in pH level up to 6 and then became static up to pH level 9 and there was no mycelial growth in pH 4.

Key words: *Aspergillus niger*, *Aspergillus flavus*, pH, Mycelial growth.

INTRODUCTION

Aspergillus is a significant part of daily life, and the genus include the human pathogen *A. fumigatus*, the plant pathogen *A. flavus*, and the industrial workhorses *A. niger*, *A. terreus*, and *A. oryzae*. When compared to different fungal genera, the *Aspergillus* genus of filamentous fungi contains a disproportionately high number of diverse species. (Mogensen *et al.*, 2006; Perrone *et al.*, 2007). At various times, particularly before harvesting and after-harvest processing, they might contaminate agricultural goods. *Aspergillus* species can cause changes in sensory, nutritional, and quality traits, including pigmentation, discoloration, spoiling, the emergence of off-odours, and off-flavours. But the spoilage of feed and food with mycotoxin is one of the most remarkable effects of their existence. Due to their parasitic nature, the majority of them are found on plant items as storage moulds (Kozakiewicz, 1989). The ecology of spoilage fungi, particularly mycotoxigenic organisms, has been found to depend significantly on pH, temperature, and the activity of water (Abdel-Hadi & Magan, 2009). 'Hydrogen ion concentration' (pH) has been demonstrated to have an impact on fungal development, either directly by acting on the outside of cells or indirectly by affecting the availability of nutrients (Abubakare *et al.*, 2013). Studies on pH have shown that fungi may grow in a range of pHs, with the highest generation of dry mycelial weight and sporulation occurring in liquid medium at pH 5.5 and pH 6.5, respectively (Saha *et al.*, 2008; Deshmukh *et al.*, 2012). With this context, in vitro research was done to compare the mycelial growth of *A. niger* and *A. flavus* at various pH levels.

MATERIALS AND METHODS

Preparation of PDA medium with different pH

Formatted: Font: Italic

The following ingredients were used to make PDA media:

Potato - 200g

Dextrose - 20g

Agar - 20g

Water - 1000 ml

Peeled potato was cut into small pieces and it was boiled in 500ml of distilled water till the pieces got softened. The extract was sieved through a double-layered muslin cloth and collected in a beaker into which 20g dextrose was added. In another 500 ml of water, 20 g of agar was melted. Potato extract and melted agar were mixed together, and the final volume of the media was made up to 1000 ml by adding distilled water. 100 ml of PDA media was taken in seven separate conical flasks to adjust pH. Using 1 M NaOH or absolute glacial acetic acid, the pH of the medium was adjusted from 3 to 9. The pH level was tested using an electrical pH metre. The medium was sterilised for 15 mins at 15 psi in an autoclave.

Inoculation of *Aspergillus sp.*

Under aseptic conditions, about 20 ml of medium was poured into each sterile Petri plate. To inhibit the bacterial pathogens, a small amount of streptomycin sulphate was mixed into each plate. Mycelial discs (8mm diameter) of *A. niger* and *A. flavus* were placed in centre of the Petri plates. The Petri plates were incubated at 25±2°C for seven days, and fungus growth was monitored daily.

STATISTICAL ANALYSIS

Experiment was carried out by using completely randomised design. Data were analysed by using excel tool for CRD.

RESULT AND DISCUSSION

Effect of pH on mycelial growth of *A. niger*

In pH 3, agar did not solidify. Agar in a medium with a low pH gets hydrolyzed when autoclaved (Kurtzman *et al.*, 2011). Mycelial growth of *A. niger* was observed on seventh day after inoculation was shown in Plate 1. Results revealed that the highest mycelial growth was observed at pH 6 for *A. niger* (8.7cm) followed by pH 7 (8cm) (Fig. 1). No fungal growth was observed in pH 4. According to Zhao *et al.*, (2010), pH 5.0 to 8.0 is ideal for *Diplocarpon mali* conidial formation. Swe *et al.*, (2009) showed that the colony numbers per

ml decrease, when the medium's alkalinity increases. Growth began to slow down above pH 6 in certain cases.





Plate 1. Mycelial growth of *Aspergillus niger* on seventh day in different pH. (A) pH 4, (B) pH 5, (C) pH 6, (D) pH 7, (E) pH 8, (F) pH 9

Effect of pH on mycelial growth of *A. flavus*

Mycelial growth of *A. flavus* was observed on seventh day after inoculation was shown in Plate 2. The results indicated that pH 6 (8.6 cm) had the maximum mycelial growth for *A. flavus*. At pH 4, no mycelial development was seen. Table 1 shows that there was no discernible variation in *A. flavus* mycelial development between pH values of 5 and 9.






Plate 2. Mycelial growth of *Aspergillus flavus* on seventh day in different pH. (A) pH 4, (B) pH 5, (C) pH 6, (D) pH 7, (E) pH 8, (F) pH 9




Fig. 1. Mycelial growth of *Aspergillus sp.*

This was in contrast with the results of David *et al.*, (2005), who found that regardless of water activity (w_aAw) level, the development of *Aspergillus-A. carbonarius* obtained from wine and table grapes was impacted more favourably at pH 4.0 and 7.0 than at pH 2.6. However, the range of acidic and alkaline conditions needed for fungi to develop is somewhat broad, ranging from pH 3.0 to pH 8.0, with the ideal pH being about pH 5.0 if nutritional needs are fulfilled (Pardo *et al.*, 2006). In general, filamentous fungi are known to be tolerant of acidic pH, and the majority of them have an ideal pH range between 5.0 and 6.0 for cellular development and a number of metabolic processes (Rosfarizana *et al.*, 2000).

Table 1. *Aspergillus sp.* mycelial growth at various pH levels

pH	Mycelial growth (cm)	
	<i>Aspergillus niger</i>	<i>Aspergillus flavus</i>
4.0	0.8	0.8
5.0	6.9	8.3
6.0	8.7	8.6
7.0	8.0	8.4
8.0	7.6	8.2
9.0	7.3	8.1
Mean	6.5	7.1
SEm	0.198	0.116
CD @ 5%	0.611	0.358

Formatted: Not Superscript/ Subscript

CONCLUSION

The study identified the necessary pH for *Aspergillus niger* and *A. flavus* growth. Result showed that *Aspergillus sp.* will grow in acidic to weak alkaline medium (pH 5-9). Further, it is necessary to do research on how various factors affect *Aspergillus* growth.

REFERENCE

1. Abdel-Hadi, A., & Magan, N. (2009). Influence of physiological factors on growth, sporulation and ochratoxin A/B production of the new *Aspergillus ochraceus* grouping. *World Mycotoxin Journal*, 2(4), 429-434.
2. Abubakar A., Suberu H. A., Bello I. M., Abdulkadir R., Daudu O. A., Lateef A. A.. Effect of pH on Mycelial Growth and Sporulation of *Aspergillus parasiticus*. *Journal of Plant Sciences*. Vol. 1, No. 4, 2013, pp. 64-67. doi: 10.11648/j.jps.20130104.13
3. David, M., Neus, B., Sonia, M., David, A., Vicente, S., & Naresh, M. (2005). Water relations of germination, growth and ochratoxin 'A' production by *Aspergillus*

carbonarius isolates from wine and table grapes from the northern Mediterranean basin. Ecophysiology of ochratoxigenic moulds. *Journal of Applied Microbiology*, 98, 839-844.

4. Deshmukh, A. J., Mehta, B. P., Sabalpara, A. N., & Patil, V. A. (2012). In vitro effect of various nitrogen, carbon sources and pH regimes on the growth and sporulation of *Colletotrichum gloeosporioides* Penz. and *Sacc* causing anthracnose of Indian bean. *Journal of Biopesticides*, 5, 46.
5. Kozakiewicz, Z. (1989). *Aspergillus* species on Stored Products. *Mycological Papers* 161: 1–188.
6. Kurtzman, C. P., Fell, J. W., Boekhout, T., & Robert, V. (2011). Methods for isolation, phenotypic characterization and maintenance of yeasts. In *The yeasts* (pp. 87-110). Elsevier.
7. Mogensen, J., Nielsen, H. B., Hofmann, G., & Nielsen, J. (2006). Transcription analysis using high-density micro-arrays of *Aspergillus nidulans* wild-type and *creA* mutant during growth on glucose or ethanol. *Fungal Genetics and Biology*, 43(8), 593-603.
8. Pardo, E., Marin, S., Ramos, A. J., & Sanchis, V. (2006). Ecophysiology of ochratoxigenic *Aspergillus ochraceus* and *Penicillium verrucosum* isolates. Predictive models for fungal spoilage prevention—a review. *Food additives and contaminants*, 23(4), 398-410.
9. Perrone, G., Susca, A., Cozzi, G., Ehrlich, K., Varga, J., Frisvad, J. C., ... & Samson, R. A. (2007). Biodiversity of *Aspergillus* species in some important agricultural products. *Studies in mycology*, 59(1), 53-66.
10. Rosfarizan, M., Ariff, A. B., Hassan, M. A., & Karim, M. I. A. (2000). Influence of pH on kojic acid fermentation by *Aspergillus flavus*. *Pakistan Journal of Biological Sciences*, 3(6), 977-982.
11. Saha, A., Mandal, P., Dasgupta, S., & Saha, D. (2008). Influence of culture media and environmental factors on mycelial growth and sporulation of *Lasiodiplodiatheobromae* (Pat.) Griffon and Maubl. *Journal of Environmental Biology*, 29(3), 407.
12. Swe, K. H., Alimon, A. R. and Ramin, M. (2009). Effect of Delaying Sporulation by Addition of Ammonium Sulphate on the Fermentation of Palm Kernel Cake Based Substrate by *Aspergillus niger*. *American Journal of Agriculture and Biological Sciences*. 4(4):262 – 265.

13. Zhao, H., Huang, L., Xiao, C. L., Liu, J., Wei, J. and Gao, X. (2010). Influence of Culture Media and Environmental Factors on Mycelial Growth and Conidial Production of *Diplocarpon mali*. The Society for Applied Microbiology, Letters in Applied Microbiology 50: 639–644.

UNDER PEER REVIEW