

# ***Original Research Article***

## **Deciphering the genetic variability for seed yield components in sesame (*Sesamum indicum* L.)**

### **ABSTRACT**

The present study conducted an analysis of genetic variability for components traits of seed yield in 20 sesame (*Sesamum indicum* L.) genotypes at the Research Farm of S.K.N. College of Agriculture, Jobner. The experiment followed a ~~Randomized-randomized Block block Design-design~~ with three replications during *kharif*, 2018. The main objective of this study was to assess the genetic variability among the genotypes. The results of analysis of variance (ANOVA) showed significant genotypic variation for all traits, indicating their potential for breeding programs. Among the traits studied, plant height exhibited the highest variability, while test weight had the least variability. Furthermore, phenotypic coefficients of variation (PCV) were higher than genotypic coefficients of variation (GCV) for all the traits, suggesting the influence of environmental factors on them. Notably, traits such as capsules per plant, seeds per capsule, biological yield per plant, and seed yield per plant showed both high heritability and high genetic gain. This makes them promising for selection and ~~eventually, eventually~~; these traits can be efficiently explored in different breeding programs for genetic improvement of sesame.

*Keywords: Coefficient of variance, Genetic advance, Genetic variability, Heritability, Sesame*

### **1. INTRODUCTION**

Sesame (*Sesamum indicum* L.) is a diploid species ( $2n=2x=26$ ) with an autogamous breeding system. It belongs to the family Pedaliaceae. It has been known by various names such as Gingelly, Til, Beniseed, and Nuvvulu in different regions of the country [1], sesame holds a significant status as the "queen of oilseeds" [2]. Ancient records point to Ethiopia (Africa) as the likely place of origin for sesame, and it was later introduced to India and China. Over a period of time, it became a staple food in Southern Asia and various other parts of the continent [3]. The crop's widespread cultivation and utilization have made it an essential agricultural commodity with both culinary and economic significance. It is the 3<sup>rd</sup> major oilseed crop after groundnut and rapeseed-mustard [4]. Sesame seeds are rich source of energy, comprising approximately 50-60% oil, 18-25% protein and 13.5% carbohydrate [5]. Due to its cholesterol-free nature, sesame oil is often recommended for heart patients. Moreover, sesame seeds contain specific compounds that possess antioxidant as well as anti-pathogenic properties [6, 7]. These health benefits make sesame a valuable addition to various culinary and medicinal applications.

Genetic variability has utmost importance in offering fundamental understanding of the nature and scope of variation existing within a crop species. Genetic variability serves as a pioneer step for any crop improvement program [8]. The effective and systematized exploration of genetic diversity is vital for the success of crop improvement programmes [9]. Genetic variability assists in determining the level of parental diversity necessary to obtain superior genotypes in segregating populations. Moreover, it proves to be highly effective in predicting potential genetic gain [10]. It has been observed that, a genotype is influenced by

**Comment [H1]:** The study describe GCV and PCV for selected sesame varieties for yield attributing charaters. The study is important in deciding the characters for improvement in plant improvement programme; while repetition of varieties (plant material) selected for study will not be given any addition information. It is accepted that, authors have taken care to avoid such repetition of varieties for study and also have selected one or two varieties as a check for effective comparison. Variety wise data is not given.

**Formatted:** Left: 1.4", Right: 1.4", Bottom: 1.4"

**Comment [H2]:** Write the percent (quantifiable) variation due to phenotypic and genotypic characters for important parameter rather than general statement about "PCV > GCV"; Enlist few characters with their heritability or genetic gain values rather than enlisting all of them having highgenetic gain.

**Comment [H3]:** Some information need to be provided for the results of findings of earlier study rather than just citing references. Research gap should be specific to the MS not the generalize research gap (For e.g. Location, or type of plant material, or methods/ procedure used or parameters specification, etc.). Avoid the generalize statements as research gap.

environmental factors such as rainfall, day length, temperature etc. [11], and these factors contribute to the non-heritable variations. Therefore, genetic variability studies play a key role to categorize the overall variation into heritable and non-heritable components. Moreover, heritability and genetic advance are the key factors determining effective trait selection during breeding programs. The knowledge of heritability and genetic advance assist a plant breeder to make correct decisions during the selection process, leading to more effective crop improvement programs and the development of superior cultivars with desirable traits.

Previously, various studies have been undertaken to study genetic variability in sesame, targeting various agro-morphological [12], seed yield [13, 14] and oil [15] traits using diverse germplasm panel. Moreover, genetic divergence in sesame employing molecular [16] as well as phenotypic diversity [17, 18] have divergence in sesame employing molecular [16] as well as phenotypic diversity [17, 18] has also been studied. Despite the considerable efforts made by various researchers to improve the economic traits of sesame, organized research in this area still lags behind as compared to other oilseed crops [19, 20]. Therefore, present study has been planned with an objective of estimating genetic variability parameters for components traits of seed yield in sesame.

## 2. MATERIAL AND METHODS

### 2.1 Plant materials

The plant materials for present study consisted of twenty diverse genotypes of sesame viz., GT-10, RT- 372, RT- 125, RT- 46, RT- 383, RT- 103, RT-127, RT- 346, RT- 351, RT- 378, RMT- 486, PRAGATI, RT- 384, RT- 385, RMT- 425, TKG-22, RMT- 447, RMT- 479, RMT- 505 and RMT-450, acquired from Agriculture University, Jodhpur, Rajasthan.

### 2.2 Design of experiment

The current study was conducted during the *Khariif* season of 2018 at the research Research farm-Farm of S.K.N. Agriculture University, Jobner, Rajasthan. Twenty sesame genotypes were evaluated in 3 replications with randomized block design. Each individual plot size was 4.0 m x 0.6 m<sup>2</sup>, containing two rows of each genotype in each replication. The spacing between rows and plants was maintained at 30 cm and 10 cm, respectively. All other package and practices were carried out according to recommended standards.

### 2.3 Record of observations

In the current study, ten characters viz., days to 50 per cent flowering (DFF), days to maturity (DM), plant height (PH), branches per plant (BPP), capsules per plant (CPP), seeds per capsule (SPC), test weight (TW), biological yield per plant (BYPP), harvest index (HI) and seed yield per plant (SYPP) were studied and observations were recorded. To record observations for PH, BPP, CPP, SPC, TW, BYPP, HI and SYPP five plants were selected randomly from individual plot and their mean value was calculated, whereas in case of DFF and DM data was recorded on whole plot basis.

### 2.4 Data analysis

The analysis of variance (ANOVA) was done as per Panse and Sukhatme [21]. Genotypic ( $\sigma_g^2$ ) and phenotypic variance ( $\sigma_p^2$ ) assessed as per Johnson et al. [22]:

$$\sigma_g^2 = \frac{MSg - MSe}{r}$$

$$\sigma_p^2 = \sigma_g^2 + \sigma_e^2$$

**Comment [H4]:** Information crop cultivation/ growing practices need to be given at least in short. Information agroclimatic condition of the place of experimentation need to be given (Also weather condition during the period of experimentation need to be given). The reason or justification for selection of varieties need to be given along with name and source of varieties. Mention the type of seed selected (breeder, foundation or certified seed).

**Formatted:** Font: Italic

**Comment [H5]:** Check the notation for correctness.

Where,  $r, MSg, MSe, \sigma_g^2, \sigma_p^2$  and  $\sigma_e^2$  denoting number of replications, mean sum of square due to genotypes, mean sum of square due to error, genotypic variance, phenotypic variance and environmental variance, respectively.

GCV and PCV ~~was were~~ calculated as per Burton and de Vane [23]:

$$GCV = \frac{\sigma_g}{\bar{x}} \times 100$$

$$PCV = \frac{\sigma_p}{\bar{x}} \times 100$$

GCV, PCV and  $\bar{x}$  denotes genotypic coefficients of variation, phenotypic coefficients of variation and overall mean, respectively.

Heritability ( $h^2$ ) in broad sense was calculated as per Hanson et al. [24]:

$$h^2 = \frac{\sigma_g^2}{\sigma_p^2} \times 100$$

Heritability values were categorized according to their percentage range: low (<30%), moderate (30-60%), and high (>60%) [25].

Prediction of expected genetic gain was elucidated as per Johnson et al. [22]:

$$GA = i \cdot h^2 \cdot \sigma_p$$

$$GAM = \frac{i \cdot h^2 \cdot \sigma_p}{\bar{x}}$$

Where,  $i, h^2, \sigma_p, GA$  and  $GAM$  represent selection intensity, heritability, phenotypic standard deviation, genetic advance and genetic advance as percentage mean, respectively. Genetic advance was classified into three groups: low (0-10%), moderate (10-20%) and high (>20%) [25].

### 3. RESULTS AND DISCUSSION

#### 3.1 Analysis of variance (ANOVA)

ANOVA represents the extent of variability among the genotypes [17]. It also partitions the total variation into different components. In the present investigation, genotypes have shown significant genotypic variation for all the traits viz., DFF, DM, PH, BPP, CPP, SPC, TW, BYPP, HI and SYPP (Table 1). This indicates that all the studied genotypes have substantial variation for all the traits and these genotypes could be further utilized in different breeding programs. In support to the current findings, Anbanandan et al.[26] and Hika et al. [12] observed highly significant difference for different agro-morphological and seed traits.

Table 1. Analysis of variance showing mean squares for different genotypes in sesame

Sources of variation	df	Days to 50% flowering	Days to maturity	Plant height (cm)	Branches per plant	Capsules per plant	Seeds per capsule	Test weight (g)	Biological yield per plant (g)	Harvest index (%)	Seed yield per plant (g)

**Comment [H6]:** Result for Table 1 need to be explained with more details and also discussed in more elaborative form (reason of such variations and impact of weather and climate on performance).

The varetial variations was not given in result and discussion section. Give the information and table also for varieties also.

**Comment [H7]:** Why this sentence has been referenced with reference showing sesame variation study?

Replication	2	2.07	11.22	38.10	0.39	2.74	57.58	0.02	0.37	28.65	1.01
Genotypes	19	14.56**	13.88**	192.41**	0.50**	96.47**	148.97**	0.04**	26.17**	59.43**	3.48**
Error	38	3.72	4.23	32.83	0.14	4.98	23.37	0.01	4.18	15.33	0.41
CV		4.35	2.37	6.18	13.56	7.08	9.64	3.09	12.03	13.29	12.86
SEm ±		1.11	1.19	3.31	0.22	1.29	2.79	0.06	1.18	2.26	0.37
CD ( $\alpha=5\%$ )		3.18	3.41	9.48	0.63	3.69	7.99	0.17	3.38	6.47	1.06

\*, \*\* = significant at 5% and 1% levels, respectively

### 3.2 Genetic variability parameters, heritability and genetic advance

The present study has been undertaken on 20 sesame genotypes for ten different seed yield traits. The mean performance of different traits viz., DFF, DM, PH, BPP, CPP, SPC, TW, BYPP, HI and SYPP were 44.37 days, 86.68 days, 92.66 cm, 2.76, 31.53, 50.15, 3.24 g, 17.00 g, 29.47% and 4.98 g, respectively (Table 2). Moreover, these traits have shown diverse range; the maximum range was recorded in case of plant height (83.13-108.40 cm), showing more divergence for plant height, whereas test weight had minimum range (3.00-3.51 g) indicating least divergence (Table 2). Similarly, genotypic and phenotypic variance were found to be maximum for plant height and minimum for test weight. Genotypic variance indicates the true variation present in a genotype, whereas phenotypic variance is cumulative effect of genotypic and environmental variance [25].

The estimates of phenotypic coefficient of variation (PCV) were of higher magnitude than estimates of genotypic coefficient of variation (GCV) for all the characters. This showed that the apparent variation is not only due to genotypes but also due to the influence of environment. Similar observations were also made by Patil and Loksha [27], while assessing the genetic variability among mutant lines of sesame. In the current study, we observed that SYPP had maximum value of GCV and PCV with magnitude of 20.32 and 24.07%, respectively; indicating greater variability among the genotypes for seed yield, whereas, DM had minimum value of GCV and PCV i.e., 2.07 and 3.15%, respectively; indicating less variation for maturity among the genotypes (Table 2). Moreover, traits like DFF and TW had considerable low magnitude of variation, while remaining traits viz., PH, BPP, CPP, SPC, BYPP and HI had considerably high magnitude of variation. In line to this, similar results were observed by Bharathi et al.[28], where high GCV and PCV was observed for plant height, branches per plant, capsules per plant and seed yield.

Heritability is an important genetic parameter, which determines the effectiveness of selection [22, 29]). Based on the heritability, traits are characterized as follows, if  $h^2$  is <30% it is low heritable, 30-60% it is moderately heritable and >60% considered as high heritable [25]. In present investigation, DFF, DM, BPP, TW and HI were recorded as moderately heritable with the percentage magnitude of 49.25, 43.22, 46.15, 50.00 and 48.95, respectively (Table 2). On the other hand, five traits viz., PH, CPP, SPC, BYPP and SYPP were categorized as highly heritable traits with 61.84, 85.96, 64.18, 63.68 and 71.33% heritability, respectively (Table 2). Therefore, selection for these traits will be more effective as major portion of genetic variation is exhibiting additive gene effects.

**Table 2. Genetic variability parameters for yield and its contributing traits**

Characters	Mean	Range	$h^2$ (%)	$\sigma_g^2$	$\sigma_e^2$	$\sigma_p^2$	GCV (%)	PCV (%)	GA	GAM (%)
Days to 50% flowering	44.37	40.67-51.00	49.25	3.61	3.72	7.33	4.28	6.10	2.75	6.20
Days to maturity	86.68	84.00-93.00	43.22	3.22	4.23	7.45	2.07	3.15	2.43	2.80

Plant height (cm)	92.66	83.13-108.40	61.84	53.2	32.83	86.03	7.87	10.01	11.81	12.75
Branches per plant	2.76	2.00-3.60	46.15	0.12	0.14	0.26	12.60	18.42	0.49	17.75
Capsules per plant	31.53	21.40-41.60	85.96	30.5	4.98	35.48	17.52	18.89	10.55	33.46
Seeds per capsule	50.15	42.20-65.67	64.18	41.87	23.37	65.24	12.90	16.11	10.68	21.29
Test weight (g)	3.24	3.00-3.51	50.00	0.01	0.01	0.02	3.04	4.33	0.14	4.39
Biological yield per plant (g)	17.00	10.60-21.90	63.68	7.33	4.18	11.51	15.93	19.96	4.45	26.19
Harvest index (%)	29.47	19.98-36.93	48.95	14.7	15.33	30.03	13.01	18.60	5.53	18.76
Seed yield per plant (g)	4.98	3.17-6.75	71.33	1.02	0.41	1.43	20.32	24.07	1.76	35.35

In addition, genetic advance is another important genetic parameter for selection of different characters. Genetic advance is categorized as low (<10%), moderate (10-20%), high (>20%) [25]. In the present study, PH, CPP and SCP have shown moderate genetic advance (>10%), whereas, remaining traits have shown low genetic advance (<10%). Genetic advance shows the expected amount of genetic gain in the progeny through selection [30]. Interestingly, when genetic advance was explained in terms percentage of trait mean (GAM), CPP, SPC, BYPP and SYPP shown high genetic advance, with percentage genetic gain of 33.46, 21.29, 26.19 and 35.35, respectively (Table 2). Similar results were obtained by Bharathi et al.[28] for capsules per plant, seeds per capsules and seed yield. High GAM indicates the effectiveness of selection and rapid genetic improvement of the traits under study in successive generations[14].

#### 4. CONCLUSION

In the present investigation, efforts have been made to study genetic variability for twenty sesame genotypes. From this study, we conclude that genotypes have vast diversity among themselves, showing substantial variation for seed yield traits. Moreover, genetic variability studies are crucial for understanding variation within crop species and serve as a critical starting point for crop improvement programs. Based on the different genetic parameters, it was found that the germplasm panel was more diverse for branches per plant, capsules per plant, seeds per capsule, biological yield per plant, harvest index, and seed yield per plant. It was also noted that traits such as capsules per plant, seeds per capsule, biological yield per plant and seed yield per plant, have shown high heritability as well as high genetic gain. Therefore, selection would be more rewarding for them and these traits can be efficiently explored in different breeding programs for genetic improvement of sesame.

#### REFERENCES

1. Priyadarshini A, Umesha C, Meshram MR. influence of phosphorus and potassium levels on growth, yield and economics of sesame (*Sesamum indicum* L.) under eastern Uttar Pradesh condition. Biol. Forum Int. J 2021a;13:645-50.
2. Priyadarshini A, Umesha C, Meshram R. Effect of different levels of phosphorus and potassium on yield, yield attributes and oil content of Sesame (*Sesamum Indicum* L.). Environment Conservation Journal. 2021b;22(1-2):183-90.
3. Nehra A, Gothwal DK, Jeeterwal RC, Puniya SS, Gupta D, Ahmad S. The study of heterosis for seed yield and its attributes in sesame (*Sesamum indicum* L.) under normal environmental conditions of Rajasthan. Pharma Innov J. 2021;10(10):175-9.
4. Patel MK, Singh D, Sharma V, Tiwari D, Ola MP, Kumar S et al. Stability Analysis in Sesame (*Sesamum indicum* L.) under different Doses of Fertilizers. Biol. Forum Int. J. 2023; 15(2):365-70.

**Comment [H8]:** Conclusion should be concise and address the objective; Avoid writing it as result and discussion; Write name and percent GCV and PCV for important charactes and second stateemnt of of name of varieties showing higher GCV and PCV;

**Comment [H9]:** Considering th size of result and data presented in MS, number of fererences are more; reduce to 15—20 numbers.

5. Prasad N, Sanjay KR, Prasad DS, Vijay N, Kothari R. A review on nutritional and nutraceutical properties of sesame. *Nutrition & food. J Nutr.* 2012;2(2):127.
6. Jat L, Yadav SS, Dhayal BC, Yadav G, Choudhary KM, Bera M. Effect of sulphur fertilization and varieties on sulphur use efficiency, yield attributes and yield of sesame. *Journal of Pharmacognosy and Phytochemistry.* 2017;6(4):717-20.
7. Alshahrani S, Al Sreaya AA, Mashyakhi MY, Alqahtani S, Sivakumar SM, Alhazmi HA, Rehman Z, Alam F. Chemical characterization and antibacterial efficacy of Saudi sesame oil against human pathogenic bacteria. *Environment Conservation Journal.* 2020;21(1-2):19-29.
8. Ola MP, Patel MK, Kumar A, Ram M, Pant NC. Genetic variability for seed yield and its component characters in taramira (*Eruca sativa* Mill.). *Pharma Innov J.* 2023;12(3): 3102-05.
9. Allard RW. Principles of plant breeding. John Wiley and Sons. Inc. New York. 1960;485.
10. Harisha R, Bhadru D, Bhargava K, Vanisri S, Shankar VG, Balakrishnan AP, Gowda MM, Singhal S. Assessment of Variability and Genetic Diversity for Elite Rice (*Oryza sativa* L.) Genotypes of Telangana and Andhra Pradesh. *International Journal of Environment and Climate Change.* 2022;12(11):3612-22.
11. Valiki SR, Ghanbari S, Golmohammadzadeh S, Kiasari KR. Effect of different plant density on growth and yield of three cultivars of sesame (*Sesamum indicum* L.). *Biol. Forum Int. J.* 2015;7:1524-28.
12. Hika G, Geleta N, Jaleta Z. Genetic variability, heritability and genetic advance for the phenotypic traits in sesame (*Sesamum indicum* L.) populations from Ethiopia. *Science, Technology and Arts Research Journal.* 2015;4(1):20-6.
13. Ong'Injo EO, Ayiecho PO. Genotypic variability in sesame mutant lines in Kenya. *African crop science journal.* 2009;17(2):101-07.
14. Kiruthika S, Narayanan SL, Parameswari C, Mini ML, Arunachalam P. Genetic variability studies for yield and yield components in sesame (*Sesamum indicum* L.). *Electronic journal of plant breeding.* 2018;9(4):1529-37.
15. Pavani K, Lal Ahamed M, Ramana JV, Sirisha AB. Studies on genetic variability parameters in sesame (*Sesamum indicum* L.). *International Journal of Chemical Studies.* 2020;8(4):101-4.
16. Gebremichael DE, Parzies HK. Genetic variability among landraces of sesame in Ethiopia. *African Crop Science Journal.* 2011;19(1):1-13.
17. Tanwar A, Bisen R. Genetic diversity analysis in sesame (*Sesamum indicum* L.) germplasm based on morphological and quality traits. *Electronic Journal of Plant Breeding.* 2018;9(1):9-17.
18. Swapna B, Thippani S, Padmaja D, Nayak SS. Genetic Diversity for Yield and Yield Contributing Characters in Sesame (*Sesamum indicum* L.). *International Journal of Environment and Climate Change.* 2023;13(8):1717-25.
19. Solanki ZS, Rajpurohit BS, Gupta D. Phenotypic stability for seed yield in sesame (*Sesamum indicum* L.). *Indian Journal of Agricultural Sciences.* 2001; 71(3), 201-202.

20. Ahmed AS, Das S, Saharia DD, Sarma MK, Nath MP, Talukdar JC. Assessment of Genetic variability and Character Association for Morpho-physiological Attributes and Seed Yield in Sesame. Biol. Forum Int. J. 2022;14(2):425-30.
21. Panse VG, Sukhatme PV. Statistical methods for agricultural workers. I.C.A.R. New Delhi, 1985.
22. Johnson HW, Robinson HF, Comstock RE. Estimates of genetic and environmental variability in soybeans 1. Agronomy journal. 1955;47(7):314-18.
23. Burton GW, Devane DE. Estimating heritability in tall fescue (*Festuca arundinacea*) from replicated clonal material 1. Agronomy journal. 1953;45(10):478-81.
24. Hanson CH, Robinson HF, Comstock RE. Biometrical studies of yield in segregating populations of Korean lespedeza 1. Agronomy journal. 1956;48(6):268-72.
25. Yeshiwas Y, Negash B. Genetic variability, heritability and genetic advance of growth and yield components of garlic (*Allium sativum* L.) germplasms. Journal of Biology, Agriculture and Healthcare. 2017;7(21):2224-3208.
26. Anbanandan V. Genetic variability and heritability in sesame (*Sesamum indicum* L.). European Journal of Biotechnology and Bioscience. 2018;6(5):69-70.
27. Patil MK, Lokesha R. Estimation of genetic variability, heritability, genetic advance, correlations and path analysis in advanced mutant breeding lines of sesame (*Sesamum indicum* L.). J. Pharmacogn. Nat. Prod. 2018;4:1-5.
28. Bharathi D, Rao VT, Mohan YC, Bhadru D, Venkanna V. Genetic Variability Studies in Sesame (*Sesamum indicum* L.).Int. j. appl. biol. pharm. 2014;5(4):31-33.
29. Divya K, Rani TS, Babu TK, Padmaja D. Assessment of genetic variability, heritability and genetic gain in advanced mutant breeding lines of sesame (*Sesamum indicum* L.). Int. J. Curr. Microbiol. App. Sci. 2018;7(6):1565-74.
30. Saxena K., Bisen R. Genetic variability, heritability and genetic advance for the phenotypic traits in sesame (*Sesamum indicum* L.). Int. J. Pure App. Biosci. 2017; 5(2): 1126-31.