

Original Research Article

Effect of Fe, Zn and Bio-Fertilizers on Soil Quality and Nutrient Content of Urdbean (*Vigna mungo* L.) under Vertisols

ABSTRACT

An experiment was conducted on urdbean during *kharif* 2021 at Instructional Farm, College of Agriculture, Ummedganj, Kota (Rajasthan). The experiment comprised 10 treatments viz. (Control, 75 % RDF, 100 % RDF, 75 % RDF + *Rhizobium* @ 600 g ha⁻¹ seed inoculation, 75 % RDF + *PSB* @ 600 g ha⁻¹ seed inoculation, 75 % RDF + *Rhizobium* @ 600 g ha⁻¹ + *PSB* @ 600 g ha⁻¹, 75 % RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage, 75 % RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage + *Rhizobium* @ 600 g ha⁻¹, 75 % RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage + *PSB* @ 600 g ha⁻¹, 75% RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage + *Rhizobium* 600 g ha⁻¹ + *PSB* @ 600 g ha⁻¹) was carried out in randomized block design with three replications. Significantly higher nitrogen (3.87 and 1.84 %), phosphorus (0.34 and 0.154 %), potassium (0.56 and 1.82 %), Fe (45.15 and 73.89 mg kg⁻¹) and Zn (32.61 and 19.69 mg kg⁻¹) content in seed and straw of urdbean were recorded with application of 75% RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ foliar spray at pre flowering and pod formation stage + *Rhizobium* 600 g ha⁻¹ + *PSB* @ 600 g ha⁻¹ seed inoculation over rest of treatments. The significantly higher uptake of nitrogen (49.36, 27.84 and 77.20 kg ha⁻¹), phosphorus (4.32, 2.33 and 6.65 kg ha⁻¹), potassium (7.16, 27.44 and 34.60 kg ha⁻¹), Fe (573.25, 1120.36 and 1693.60 g ha⁻¹) and Zn (413.78, 297.40 and 711.18 g ha⁻¹) by seed, straw and total by urdbean crop with application of 75% RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ foliar spray at pre flowering and pod formation stage + *Rhizobium* 600 g ha⁻¹ + *PSB* @ 600 g ha⁻¹ seed inoculation over rest of treatments. Application of Fe, Zn foliar spray and bio fertilizers seed inoculation did not significantly influenced soil physico-chemicals properties during the experimentation. The maximum available nitrogen (285.21 kg ha⁻¹), phosphorus (25.41 kg ha⁻¹), potassium (252.66 kg ha⁻¹), Fe (5.75 mg kg⁻¹) and Zn (0.76 mg kg⁻¹) after urdbean crop harvest in soil were recorded with application of 75% RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ foliar spray at pre flowering and pod formation stage + *Rhizobium* 600 g ha⁻¹ + *PSB* @ 600 g ha⁻¹ seed inoculation over rest of treatments.

Key words: Bio-fertilizer, nutrient content, nutrient uptake, Fe, urdbean and Zn.

INTRODUCTION

Pulses are the integral part of Indian dietary system due to richest source of protein and other nutrients. Indian population is predominantly vegetarian and protein requirement for the growth and development of the human being is mostly met with pulses. The availability of pulses per capita per day has proportionately declined from 71 g to 52 g against the minimum requirement of 70 g day⁻¹/capita [1]. Pulses are improving soil fertility by atmospheric nitrogen fixation through *Rhizobium culture*. In India pulses are grown an area of 28.34 million ha with total production of 23.15 million tonnes and productivity of 817 kg ha⁻¹ [2]. Among the pulses, urdbean [*Vigna mungo* (L.) Hepper], is an important crop in India. Urdbean is the third most important pulse crop after chickpea and pigeonpea in India. It is highly nutritious containing 24-26% protein, 1.3% fat and 60% carbohydrates on dry weight basis and it is rich source of calcium, iron and vitamins [3].

The major constraints for low yield of urdbean are lack of micronutrient application and non-adoption of proper agronomic practices. Micronutrients are equally important in plant nutrition as major nutrients. The incidence of micronutrient deficiencies in crops has increased markedly in recent years due to intensive cropping, loss of top soil by

erosion, losses of micronutrients through leaching, liming of acid soils, decreased proportions of farmyard manure or other organic sources compared to chemical fertilizers and use of marginal lands for crop production [4]. The optimum supply of micronutrients under balanced condition is very important for achieving higher productivity of urdbean. Foliar application of micronutrients has been proved to be an important asset in fertilizer application with a specific aim of increasing nutrient availability at the time of need, especially at the later stage of plant growth. Foliar application of nutrients using water soluble fertilizers is one of the possible ways to enhance the productivity of pulses. The foliar fertilization of Zn and Fe at flowering stage improves growth and yield urdbean [5].

Bio-fertilizer plays a vital role in maintaining long term soil fertility and sustainability [7]. Incorporation of FYM alone or bio-fertilizers improves available nutrient status of the soil with enhanced soil biological activity which in turn provides a congenial physical condition and improved availability of nutrient in the rhizosphere, resulting in an improvement in the crop growth and providing a better source-sink relationship [6]. Among bio-fertilizers, *Rhizobium* inoculation is cheapest, easiest and safest method of

supplying nitrogen to legumes through well-known symbiotic nitrogen fixation process. Inoculation of appropriate strain enhances nodule formation resulting better nitrogen fixation. *Rhizobium* species in association with plant roots in urdbean

MATERIALS AND METHODS

A field experiment was conducted on urdbean during *kharif* 2021 at Instructional Farm, College of Agriculture, Umedganj, Kota (Rajasthan). Geographically, is situated at 25.11⁰ North latitude and 75.50⁰ East longitude at an altitude of 258 meters above mean sea level. In Rajasthan, this region falls under the Agro-Climatic Zone-V (Humid South Eastern Plain Zone). This zone possesses typical sub-tropical conditions with maximum temperature range in summer is 42.2- 43.0 °C and minimum 12 - 27°C. In this zone annual average rainfall is received 1275 mm. The soil of experimental site was clay loam in texture, slightly saline in reaction, medium in available nitrogen (217.35 kg ha⁻¹) and phosphorus (19.82 kg ha⁻¹) while high in potassium (236.31kg ha⁻¹) and sufficient in DTPA extractable micronutrients with pH (7.60) and EC (0.49 dS m⁻¹). Source of nutrients were applied urea for nitrogen, DAP for phosphorus and mutate of potash for potassium. The full dose of fertilizer 100 % RDF (20:40:30 NPK kg ha⁻¹) was applied as basal dose. Before sowing, seed

improved soil fertility by fixing atmospheric nitrogen and produces plant growth substances in the soil. *Rhizobium* inoculation can increase the grain yield of pulse crops [8].

of urdbean was inoculated with liquid *rhizobium* culture @ 10 ml kg⁻¹ seed and liquid *PSB* culture @ 10 ml kg⁻¹ seed as per treatment. Foliar fertilization of ZnSO₄ and FeSO₄ were applied at pre flowering and pod formation stage. Spraying was done with knapsack sprayer and the leaves were wetted thoroughly with fine mist. The absorption of solution by urdbean leaves, a sticker was added in the spray solution.

The experiment laid out in randomized block design with three replications. The experiment comprised 10 treatments viz. (control, 75 % RDF, 100 % RDF, 75 % RDF + *Rhizobium* @ 600 g ha⁻¹ seed inoculation, 75 % RDF + *PSB* @ 600 g ha⁻¹ seed inoculation, 75 % RDF + *Rhizobium* @ 600 g ha⁻¹ + *PSB* @ 600 g ha⁻¹, 75 % RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage, 75 % RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage + *Rhizobium* @ 600 g ha⁻¹, 75 % RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage + *PSB* @ 600 g ha⁻¹, 75% RDF + 0.1 % FeSO₄ +

0.5 % ZnSO₄ at pre flowering and pod formation stage + *Rhizobium* 600 g ha⁻¹+ PSB @ 600 g ha⁻¹) was carried out in randomized block design with three replications. For recording pre and post-harvest observations, five plants were randomly selected for each plot and tagged with labels for various observations on seed and straw. Soil samples were recorded plot wise as per standard procedures and statistically analysed by adopting appropriate method of standard analysis of variance [9].

Chemical analysis in Plant

Plant samples were randomly collected from labelled plants in each treatment. The samples were then cleaned with water, air dried and then dried in hot-air oven at 60 °C for 18 hours. The samples were oven dried, powdered and stored in polythene bags. These powdered samples were analysed for nitrogen, phosphorus, potassium, zinc and ferrous content. Nitrogen was determined by Kjeldahl distillation method. Plant sample (0.5 g)

was digested in digestion flasks using sulphuric acid and digestion mixture. After

complete digestion, the digested material was distilled in alkaline medium and the liberated ammonia was trapped in four per cent boric acid solution containing mixed indicator. The trapped ammonia was titrated against standard sulphuric acid [10]. Phosphorus content in the di-acid digested extract was estimated by Vanadomolybdo-phosphoric yellow colour method in nitric acid medium and the colour intensity was measured at 420 nm wave length [10]. Potassium in the plant sample was estimated by atomizing the diluted di-acid extract to a flame photometer [10]. After making suitable dilution of di-acid extract, the samples were fed to the atomic absorption spectrophotometer using appropriate hallow cathode lamps to estimate the Zn content in plant sample. After making suitable dilution of di-acid extract, the samples were fed to the atomic absorption spectrophotometer using appropriate hallow cathode lamps to estimate the Fe content in plant sample. Nitrogen, phosphorus and potassium uptake by seed and stover at harvest was calculated by using following formula:

$$\begin{aligned} \text{Nutrient uptake by seed (Kg ha}^{-1}\text{)} &= \frac{\text{Nutrient content in seed (\%)} \times \text{Seed Yield (Kg ha}^{-1}\text{)}}{100} \\ \text{Nutrient uptake by straw (Kg ha}^{-1}\text{)} &= \frac{\text{Nutrient content in straw (\%)} \times \text{Straw Yield (Kg ha}^{-1}\text{)}}{100} \end{aligned}$$

Zinc and Iron uptake by plants seed and stover at harvest was calculated by using following formula:

$$\text{Nutrient uptake by seed (mg ha}^{-1}\text{)} = \frac{\text{Nutrient content in seed (\%)} \times \text{Seed Yield (Kg ha}^{-1}\text{)}}{100}$$

$$\text{Nutrient uptake by straw (mg ha}^{-1}\text{)} = \frac{\text{Nutrient content in straw (\%)} \times \text{Straw Yield (Kg ha}^{-1}\text{)}}{100}$$

Soil analysis after crop harvest

Soil samples (0-15 cm) were collected from each plot after harvest of

urdbean crop. These were dried and passed through 2 mm sieve and prepared for subsequent analysis as per standard methods.

Table 1.0: Soil physico-chemical properties analysis methods.

S. No.	Particulars	Values	Method employed
A.	Mechanical composition		
i.	Sand (%)	23.80	Hydrometer method [11]
ii.	Silt (%)	36.00	
iii.	Clay (%)	40.20	
iv.	Textural class	Clay loam	
B.	Physical properties		
i.	Bulk density (Mg m ⁻³)	1.34	Core sampler method, [10]
ii.	Particle density (Mg m ⁻³)	2.39	Pycnometer or RD bottle method
iii.	Porosity (%)	43.94	Calculated by BD and PD
C.	Chemical composition		
i.	Organic carbon (%)	0.41	Determination by rapid titration method, [12]
ii.	Available nitrogen (kg ha ⁻¹)	217.35	Estimation by alkali potassium permanganate method [13]
iii.	Available phosphorus (kg ha ⁻¹)	19.82	Olsen's P, 0.5 M NaHCO ₃ method, pH 8.5 [14]
iv.	Available potassium (kg ha ⁻¹)	236.31	Extraction was done with 1 N neutral ammonium acetate at pH 7.0 and determined by flame photometer [15]
v.	Available Zn (mg kg ⁻¹)	0.57	Extraction by 0.005 M DTPA + 0.01 M CaCl ₂ + 0.1 M tri ethanol amine at pH 7.3 [16]
vi.	Available Fe (mg kg ⁻¹)	3.93	Extraction by 0.005 M DTPA + 0.01 M CaCl ₂ + 0.1 M tri ethanol amine at pH 7.3

			[16]
vii.	EC at 25 ⁰ C (dSm ⁻¹)	0.49	Measured in soil:water solution (1:2) [17]
viii.	pH (1 :2 soil water suspension)	7.60	Glass electrode pH meter in soil:water solution (1:2) [10]

RESULTS AND DISCUSSION

Nutrient content and uptake

The data pertaining to nutrient content and uptake of urdbean crop significantly influenced under application of Fe, Zn and Bio-fertilizers were presented in Table 2.0. Significantly higher nitrogen (3.87 and 1.84 %), phosphorus (0.34 and 0.154 %), potassium (0.56 and 1.82 %), Fe (45.15 and 73.89 mg kg⁻¹) and Zn (32.61 and 19.69 mg kg⁻¹) content in seed and straw of urdbean were recorded with application of 75% RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ foliar spray at pre flowering and pod formation stage + *Rhizobium* 600 g ha⁻¹ + PSB @ 600 g ha⁻¹ seed inoculation over control nitrogen (1.66 and 1.37 %), phosphorus (0.22 and 0.177 %), potassium (0.40 and 1.33 %), Fe (45.15 and 73.89 mg kg⁻¹) and Zn (32.61 and 19.69 mg kg⁻¹), 75 % RDF nitrogen (1.94 and 1.48 %), phosphorus (0.24 and 0.123 %), potassium (0.44 and 1.48 %), Fe (45.15 and 73.89 mg kg⁻¹) and Zn (32.61 and 19.69 mg kg⁻¹), 100 % RDF nitrogen (2.45 and 1.57 %), phosphorus (0.25 and 0.129 %), potassium (0.47 and 1.57 %), Fe (45.15 and 73.89 mg kg⁻¹) and Zn (32.61 and 19.69 mg kg⁻¹), 75 % RDF +

Rhizobium @ 600 g ha⁻¹ seed inoculation nitrogen (2.49 and 1.59 %), phosphorus (0.27 and 0.133 %), potassium (0.48 and 1.60 %), Fe (45.15 and 73.89 mg kg⁻¹) and Zn (32.61 and 19.69 mg kg⁻¹), 75 % RDF + PSB @ 600 g ha⁻¹ seed inoculation nitrogen (2.46 and 1.58 %), phosphorus (0.26 and 0.131%), potassium (0.47 and 1.57 %), Fe (45.15 and 73.89 mg kg⁻¹) and Zn (32.61 and 19.69 mg kg⁻¹), 75 % RDF + *Rhizobium* @ 600 g ha⁻¹ + PSB @ 600 g ha⁻¹ nitrogen (2.53 and 1.70 %), phosphorus (0.28 and 0.141 %), potassium (0.51 and 1.67 %), Fe (45.15 and 73.89 mg kg⁻¹) and Zn (32.61 and 19.69 mg kg⁻¹), 75 % RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage nitrogen (2.52 and 1.63 %), phosphorus (0.29 and 0.134 %), potassium (0.50 and 1.62 %), Fe (45.15 and 73.89 mg kg⁻¹) and Zn (32.61 and 19.69 mg kg⁻¹), 75 % RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage + *Rhizobium* @ 600 g ha⁻¹ nitrogen (3.11 and 1.77 %), phosphorus (0.31 and 0.149 %), potassium (0.53 and 1.80 %), Fe (45.15 and 73.89 mg kg⁻¹) and Zn (32.61 and 19.69 mg kg⁻¹), 75 % RDF + 0.1 %

FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage + *PSB* @ 600 g ha⁻¹ nitrogen (2.99 and 1.75 %), phosphorus (0.30 and 0.145 %), potassium (0.53 and 1.72 %) Fe (45.15 and 73.89 mg kg⁻¹) and Zn (32.61 and 19.69 mg kg⁻¹) content in seed and straw of urdbean.

The nutrient uptake by seed, straw and by urdbean crop were significantly influenced under application of Fe, Zn and Bio-fertilizers were presented in Table 3.0. The significantly higher uptake of nitrogen (49.36, 27.84 and 77.20 kg ha⁻¹), phosphorus (4.32, 2.33 and 6.65 kg ha⁻¹), potassium (7.16, 27.44 and 34.60 kg ha⁻¹), Fe (573.25, 1120.36 and 1693.60 g ha⁻¹) and Zn (413.78, 297.40 and 711.18 g ha⁻¹) by seed, straw and total by urdbean crop with application of 75% RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ foliar spray at pre flowering and pod formation stage + *Rhizobium* 600 g ha⁻¹ + *PSB* @ 600 g ha⁻¹ seed inoculation over control nitrogen (10.04, 10.99 and 21.03 kg ha⁻¹), phosphorus (1.33, 0.94 and 2.27 kg ha⁻¹), potassium (2.41, 10.68 and 13.09 kg ha⁻¹), Fe (167.06, 344.14 and 511.20 g ha⁻¹) and Zn (120.59, 95.23 and 215.82 g ha⁻¹), 75 % RDF nitrogen (14.26, 13.76 and 28.02 kg ha⁻¹), phosphorus (1.76, 1.44 and 2.90 kg ha⁻¹), potassium (3.25, 13.76 and 17.0 kg ha⁻¹), Fe (227.51, 461.25 and 688.76 g ha⁻¹) and Zn (159.97, 125.72 and 285.69 g ha⁻¹), 100 % RDF nitrogen (20.0, 16.13 and

36.13 kg ha⁻¹), phosphorus (2.04, 1.32 and 3.37 kg ha⁻¹), potassium (3.82, 16.01 and 19.83 kg ha⁻¹), Fe (278.19, 550.66 and 828.85 g ha⁻¹) and Zn (194.90, 152.17 and 347.07 g ha⁻¹), 75 % RDF + *Rhizobium* @ 600 g ha⁻¹ seed inoculation nitrogen (21.39, 17.43 and 38.83 kg ha⁻¹), phosphorus (2.33, 1.46 and 3.79 kg ha⁻¹), potassium (4.15, 17.47 and 21.63 kg ha⁻¹), Fe (310.48, 625.75 and 936.23 g ha⁻¹) and Zn (218.81, 171.05 and 389.87 g ha⁻¹), 75 % RDF + *PSB* @ 600 g ha⁻¹ seed inoculation nitrogen (21.05, 17.22 and 38.27 kg ha⁻¹), phosphorus (2.22, 1.43 and 3.64 kg ha⁻¹), potassium (4.02, 17.11 and 21.13 kg ha⁻¹), Fe (303.01, 607.60 and 910.61 g ha⁻¹) and Zn (212.47, 168.28 and 380.74 g ha⁻¹), 75 % RDF + *Rhizobium* @ 600 g ha⁻¹ + *PSB* @ 600 g ha⁻¹ nitrogen (22.93, 19.57 and 42.51 kg ha⁻¹), phosphorus (2.53, 1.62 and 4.15 kg ha⁻¹), potassium (4.62, 19.21 and 23.83 kg ha⁻¹), Fe (336.86, 685.66 and 1022.51 g ha⁻¹) and Zn (236.50, 187.88 and 424.38 g ha⁻¹), 75 % RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage nitrogen (22.20, 18.20 and 40.40 kg ha⁻¹), phosphorus (2.56, 1.50 and 4.05 kg ha⁻¹), potassium (4.35, 18.03 and 22.38 kg ha⁻¹), Fe (325.38, 643.15 and 968.53 g ha⁻¹) and Zn (226.17, 178.83 and 404.99 g ha⁻¹), 75 % RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage + *Rhizobium* @ 600 g ha⁻¹ nitrogen (31.26,

21.99 and 53.25 kg ha⁻¹), phosphorus (3.11, 1.85 and 4.96 kg ha⁻¹), potassium (5.32, 22.44 and 27.76 kg ha⁻¹), Fe (419.13, 809.61 and 1228.75 g ha⁻¹) and Zn (304.10, 226.82 and 530.92 g ha⁻¹), 75 % RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage + PSB @ 600 g ha⁻¹ nitrogen (29.72, 21.63 and 51.35 kg ha⁻¹), phosphorus (2.98, 1.79 and 4.77 kg ha⁻¹), potassium (5.23, 21.22 and 26.46 kg ha⁻¹) Fe (410.23, 804.38 and 1214.62 g ha⁻¹) and Zn (295.57, 223.41 and 518.98 g ha⁻¹) uptake by seed, straw and total by urdbean crop.

Application of Fe, Zn and Bio fertilizers significantly influenced the nutrient content and their uptake. The increase in nutrient content might be due to increased availability of native nutrients. These results supported by the findings of [18], who stated that the increase could be attributed to the synergistic effect between nutrients and bio-fertilizers which might be due to increase enzymatic activity by different nutrient sources. It is an established fact that nutrient accumulation depends upon dry matter accumulation and concentration of nutrient at cellular level. These results are in agreement with those of [19] and [20]. Significant increase in protein content has been observed in the present investigation because of increased nitrogen content in seed which might be the result of increased availability of

nitrogen to plants. An improved metabolism to greater translocation of these nutrients to reproductive organs of the crop and ultimately increased the content in seed and stover. These results are in close conformity with those of [21]. Significant increase in nutrient uptake due to application of 75% RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ foliar spray at pre flowering and pod formation stage + *Rhizobium* 600 g ha⁻¹ + PSB @ 600 g ha⁻¹ seed inoculation recorded under investigation. Since the nutrient uptake is a function of its content in crop plant and seed and straw yield of the crop. The increase in these parameters due to nitrogen and phosphorus fertilization with bio inoculants led to an increased uptake of nutrients in the present study. These results are in conformity with the findings of [22] and [23].

Soil physico-chemical properties after crop harvest

Soil status after harvesting of urdbean were measured in terms of pH, EC (dS m⁻¹), organic carbon (%), bulk density (Mg m⁻³), particle density (Mg m⁻³) and porosity (%) are presented in Table 4.0. Soil pH after harvesting of urdbean crop did not find to show any significant variation with application of Fe, Zn and Bio-fertilizers. Among the treatments, under absolute control was recorded maximum soil pH (7.32) and minimum pH

(7.19) with application of 75% RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ foliar spray at pre flowering and pod formation stage + *Rhizobium* 600 g ha⁻¹ + PSB @ 600 g ha⁻¹ seed inoculation. It is evident from data the electrical conductivity did not significantly influence with application of Fe, Zn and Bio-fertilizers. Among the treatments, application of 100% RDF recorded maximum EC (0.68 dS m⁻¹) and least EC (0.49 dS m⁻¹) under application of absolute control during course of experimentation. Under application of Fe, Zn and Bio-fertilizers did not significantly influence organic carbon content in soil after crop harvest. It is evident from data the bulk density and particle density could not reach the level of significance due to application of Fe, Zn and Bio-fertilizers. The maximum bulk density (1.33 Mg m⁻³) and particle density (2.45 Mg m⁻³) were noted under absolute control. However, minimum value of bulk density (1.31 Mg m⁻³) and particle density (2.41 Mg m⁻³) was noted under application of 75% RDF + *Rhizobium* @ 600 g ha⁻¹ seed inoculation. Soil porosity did not find to show any significant effect due to application of Fe, Zn and Bio-fertilizers. The maximum higher porosity (45.71%) was noted under control and application of 100% RDF over rest of treatments.

The results regarding residual effect of biofertilizers (*Rhizobium* and

PSB) and spray of FeSO₄ and ZnSO₄ on soil pH, EC, porosity, bulk density, particle density and organic carbon was presented. The data showed that pH, EC, particle density, bulk density of soil did not significantly influence due to individual seed treatment with *Rhizobium* and PSB levels but the lower soil pH was recorded under the treatment 75% RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ foliar spray at pre flowering & pod formation stage + *Rhizobium* 600 g ha⁻¹ + PSB @ 600 g ha⁻¹ seed inoculation, similarly higher soil pH was observed with control. [4] revealed that the effect of various treatments on soil pH with addition of organic and inorganic fertilizer is not consistent. [24] did not found any change in EC with the treatment of bio-inoculants. Similar results were also observed in bulk density and particle density which ranged from 1.31 to 1.33, 2.41 to 2.45 Mg m⁻³ and data indicated that the difference in soil bulk density and particle density values were not reach to the levels of significance due to application of Fe, Zn and biofertilizer. Similar results were also observed in per cent organic carbon and porosity.

Available nutrients of soil after crop harvest

The data pertaining to nutrient content and uptake of urdbean crop significantly influenced under application of Fe, Zn and Bio-fertilizers were

presented in Table 4.0. The significantly higher available nitrogen ($285.21 \text{ kg ha}^{-1}$), phosphorus (25.41 kg ha^{-1}), potassium ($252.61 \text{ kg ha}^{-1}$), Fe (5.75 mg kg^{-1}) and Zn (0.76 mg kg^{-1}) content in soil after crop harvest were recorded with application of 75% RDF + 0.1 % FeSO_4 + 0.5 % ZnSO_4 foliar spray at pre flowering and pod formation stage + *Rhizobium* 600 g ha^{-1} + PSB @ 600 g ha^{-1} seed inoculation over control nitrogen nitrogen ($228.21 \text{ kg ha}^{-1}$), phosphorus (17.46 kg ha^{-1}), potassium ($230.10 \text{ kg ha}^{-1}$), Fe (4.30 mg kg^{-1}) and Zn (0.59 mg kg^{-1}), 75 % RDF nitrogen ($238.88 \text{ kg ha}^{-1}$), phosphorus (20.72 kg ha^{-1}), potassium ($241.87 \text{ kg ha}^{-1}$), Fe (4.64 mg kg^{-1}) and Zn (0.62 mg kg^{-1}), 100 % RDF nitrogen ($245.84 \text{ kg ha}^{-1}$), phosphorus (21.28 kg ha^{-1}), potassium ($243.57 \text{ kg ha}^{-1}$), Fe (4.70 mg kg^{-1}) and Zn (0.65 mg kg^{-1}), 75 % RDF + *Rhizobium* @ 600 g ha^{-1} seed inoculation nitrogen ($266.92 \text{ kg ha}^{-1}$), phosphorus (21.79 kg ha^{-1}), potassium ($245.39 \text{ kg ha}^{-1}$), Fe (5.15 mg kg^{-1}) and Zn (0.67 mg kg^{-1}), 75 % RDF + PSB @ 600 g ha^{-1} seed inoculation nitrogen ($260.33 \text{ kg ha}^{-1}$), phosphorus (23.77 kg ha^{-1}), potassium ($247.51 \text{ kg ha}^{-1}$), Fe (5.07 mg kg^{-1}) and Zn (0.66 mg kg^{-1}), 75 % RDF + *Rhizobium* @ 600 g ha^{-1} + PSB @ 600 g ha^{-1} nitrogen ($272.88 \text{ kg ha}^{-1}$), phosphorus (24.0 kg ha^{-1}), potassium ($250.82 \text{ kg ha}^{-1}$), Fe (5.18 mg kg^{-1}) and Zn (0.69 mg kg^{-1}), 75 % RDF + 0.1 % FeSO_4 + 0.5 % ZnSO_4

at pre flowering and pod formation stage nitrogen ($258.54 \text{ kg ha}^{-1}$), phosphorus (22.74 kg ha^{-1}), potassium ($249.71 \text{ kg ha}^{-1}$), Fe (5.25 mg kg^{-1}) and Zn (0.71 mg kg^{-1}), 75 % RDF + 0.1 % FeSO_4 + 0.5 % ZnSO_4 at pre flowering and pod formation stage + *Rhizobium* @ 600 g ha^{-1} nitrogen ($280.17 \text{ kg ha}^{-1}$), phosphorus (23.61 kg ha^{-1}), potassium ($254.75 \text{ kg ha}^{-1}$), Fe (5.48 mg kg^{-1}) and Zn (0.74 mg kg^{-1}), 75 % RDF + 0.1 % FeSO_4 + 0.5 % ZnSO_4 at pre flowering and pod formation stage + PSB @ 600 g ha^{-1} nitrogen ($275.88 \text{ kg ha}^{-1}$), phosphorus (24.47 kg ha^{-1}), potassium ($250.15 \text{ kg ha}^{-1}$), Fe (5.44 mg kg^{-1}) and Zn (0.72 mg kg^{-1}) content in soil after crop harvest.

Application of 75% RDF + 0.1 % FeSO_4 + 0.5 % ZnSO_4 foliar spray at pre flowering & pod formation stage + *Rhizobium* 600 g ha^{-1} + PSB @ 600 g ha^{-1} seed inoculation showed the significant effect on the available nitrogen, phosphorus, Zn and Fe in soil. Significant buildup of the soil available nitrogen could be attributed to increased activity of nitrogen fixing rhizobia there by resulting in higher accumulation of nitrogen in the soil leading to better nodulation and mineralization of organic nitrogen with phosphorus application [25]. These results were in close conformity to [26], who reported that the application of microbial culture along with phosphorus enhances

the nitrogen and phosphorus content in chickpea. [27] reported that the PSB are known to have ability to solubilize phosphorus from insoluble source. The PSB secretes the different organic acids which act on insoluble phosphate to convert them in to soluble phosphate near the root of the plant and hence availability of phosphorus is increased. [28] who had indicated that the judicious integration of organic and inorganic sources of nutrition significantly improved the soil available nitrogen, phosphorus and micronutrient in soil. Further, the buildup of available phosphorus with phosphorus application can be attributed to an increase in the available soil phosphorus after satisfying phosphorus fixation capacity and other chemical reaction [29]. [30] and [31] who observed increased available phosphorus content in soil after the harvest of the crop due to phosphorus application. The increase in available phosphorus content of

CONCLUSION

It is concluded that the application of 75% RDF + Fe 0.1 % FeSO₄ + Zn 0.5 % ZnSO₄ at pre flowering and pod formation stage + *Rhizobium* 600 g ha⁻¹ +

REFERENCES

1. Anonymous. The Directorate of Pulses Development. Pulses in India

soil might be due to greater mobilization of native soil phosphorus by vigorous root proliferation and contribution through biomass. These results find support from the results of [32], [33] in soybean. The application of 75% RDF with iron and zinc application significantly increased the available nitrogen and phosphorus content in soil [34]. As expected, a linear increase in available zinc and iron content in soil was observed with application of Zn @ 0.5% and Fe @ 0.1% through ZnSO₄ and FeSO₄, after harvest of the crop. The experimental soil being low in available zinc and iron might have resulted in increased available zinc and iron application. There could be a priming effect which possibly caused solubilization of native zinc and iron with increase in the rate of zinc and iron application. The result was in conformity to those of reported by [35] and [36].

PSB @ 600 g ha⁻¹ found most suitable dose for urdbean to increase nutrient content & uptake and also beneficial for soil health.

Retrospect and Prospects. Bhopal, M. P.2017.

2. Anonymous. Agricultural Statistics at a Glance. Government of India. Ministry of Agricultural and Farmers Welfare. Department of Agriculture, Cooperation & Farmers Welfare". *Directorate of Economics and Statistics*, New Delhi, India.2020.
3. Jadhav S C, Sawant S P P, Sanap S P, Puranik U Y, Prabhudesai S S and Devmore J P. Effect of micronutrients on yield, nutrient uptake and quality of coriander (*Coriandrum sativum*) in lateritic soil Konkar region. *International Journal of Chemical Studies*.2017; 5(4): 214-216.
4. Singh D K, Lekh C, Singh K N and Singh J K. Effect of different biofertilizers in combination with chemical fertilizers on pea (*Pisum sativum* L.) under temperate Kashmir conditions". *Environment and Ecology*.2006; 24(3): 684-686.
5. Saviour M N and Stalin P. Influence of zinc and boron in residual blackgram productivity". *Indian Journal of Science and Technology*.2013; 6(8): 5105-5108.
6. Khandelwal R, Choudhary S K, Khangarot S S, Jat M K and Singh P. Effect of inorganic and biofertilizers on productivity and nutrients uptake in cowpea [*Vigna Unguiculata* (L.) walp]. *Legume Research*. 2012; 35(3): 235-238.
7. Kudi V K and Singh J K. Effect of biofertilizers and fertility levels on blackgram (*Vigna mungo* L.) under custard apple (*Annona squamosa* L.) based agri-horti system in Vindhyan region of Uttarpradesh. *International Journal of Agriculture Sciences*. 2016; 8(52): 2534-2537.
8. Ali M and Chandra S. Rhizobium inoculation of pulse crop. *Indian Farming*. 1985; 35(5): 22-25.
9. Fisher R A. *Statistical methods for research workers* Oliver and Boyd, Edinburg, London.1950.
10. Piper C S. *Soil and Plant Analysis*. International Science Publication, John Willey and Sons. Inc. New York.1950.
11. Bouyoucos H J. A hydrometer method for making practical size analysis of soils. *Agronomy Journal*. 1962; 55: 464-465.
12. Walkley A and Black I A. Estimation of soil organic carbon by the chromic acid digestion method. *Soil Science*. 1934;37: 29-38.
13. Subbiah B V and Asija G L. A rapid procedure for the estimation of available nitrogen in soils. *Current Science*. 1956; 25: 259-260.
14. Olsen S R, Cole C V, Watenable F S and Dean L A. Estimation of available phosphorous by extraction with sodium bicarbonate. United State Department of Agriculture. Circular No. 939, Washington 19-33.1954.

15. Hanway J J and Heidel H. Soil analysis method as used in Iowa State. College soil testing laboratory Bulletin. 1952;57:1-131.
16. Lindsay W L and Norvel W A. Development of a DTPA soil test for zinc, iron, manganese and copper. Soil Science Society of America Journal. 1978; 42: 421-428.
17. Richards L A. Diagnosis and Improvement of Saline and Alkaline Soils. USDA Handbook, No.60.1954.
18. Keram K S, Sharma B L and Sawarkar S D. Impact of zinc application on yield, quality, nutrients uptake and soil fertility in a medium deep black soil (Vertisol). International Journal of Science, Environment and Technology. 2012; 1(5): 563-571.
19. Choudhary R, Sadhu A C and Gegiya K M. Effect of fertility levels and biofertilizers on yield and quality of kharif green gram (*Vigna radiata* (L.) Wilczek). Haryana Journal of Agronomy. 2010; 26: 41-44.
20. Ranpariya V S and Polara K B. Effect of potassium, zinc and FYM on content and uptake of nutrients of summer green gram (*Vigna radiata* L.) at different growth stages under south saurashtra region of Gujarat. International Journal of Pure and Applied Biosciences. 2018;6(1): 997-1002.
21. Rajkhowa D J, Saikia M and Rajkhowa K M. Effect of vermicompost with and without fertilization of greengram. Legume Research.2020; 25(4): 295-306.
22. Sesode J H. Response of green gram (*Vigna radiata* L.) to fertility levels and sulphur sources application". Agriculture Science Digest.2008; 28 (1): 26-28.
23. Rathore D S, Parothit H S and Yadav B L. Integrated phosphorus management on yield and nutrientuptake of urdbean under rainfed conditions of southern Rajasthan. Journal of Food Legumes.201; 23:128-131.
24. Govindan K and Thirumurugan V. Effect of Rhizobium and PSM's in soybean- A review. Journal of Maharashtra Agricultural Univercity.2003; 28 (1): 054-060.
25. Chesti M H, Tahir A and Bhat M A. Effect of organic and inorganic phosphorus sources in quality of green gram (*Vigna radiata* L.). Legume Research.2012; 35(1): 47-49.
26. Sarawagi S K, Tiwati S K and Tripathi R S.Uptake and balance sheet of nitrogen and phosphorus in gram (*Cicer aritenium*) as influenced by phosphorus, biofertilizers and micronutrients under rainfed condition. Indian Journal of Agronomy.1999; 44(4): 768-772.

27. Kumar A, Singh S S, Kumar R, Singh A K, Kumawat N. Response of Rhizobium and different levels of molybdenum on growth, nodulation and yield of blackgram (*Vigna mungo* L.). *Environment and Ecology*.2010; 28 (3): 1728-1730.
28. Konthoujam N D, Tensubam B S, Athokpam H S, Naorem B S and Shamurailatpam D. Influence of inorganic, biological and organic manures on nodulation and yield of soybean (*Glycine max* L. Merrill) and soil properties. *Australian Journal of Crop Science* .2013; 7(9): 1407-1415.
29. Dubey B K, Sharma C P and Chatterjee C. Response of pigeonpea to applied zinc in ustifluent soils of western Uttar Pradesh. *Journal of Indian Society of Soil Science*.2001; 49(30): 471-475.
30. Duraisami M and Mani A. Agronomic evaluation of promising genotypes of mungbean under hyper arid conditions of Indian thar desert. *International Journal of Agricultural Research*.2001; 2(6): 537-544.
31. Nimje P M. Effect of phosphorus fertilization on soybean (*Glycine max* L.) based cropping sequence under rainfed conditions. *Indian Journal of Agricultural Science*.2003; 73: 191-193.
32. Kumar A and Elamathi S. Effect of nitrogen levels and rhizobium application methods on yield attributes, yield and economics of black gram (*Vigna mungo* L.). *International Journal of Agriculture Sciences*.2007; 3(1): 179-180.
33. Raja D and Takankhar V G. Effect of liquid biofertilizers (Brady rhizobium and PSB) on availability of nutrients and soil chemical properties of soybean (*Glycine Max* L.). *International Journal of Pure and Applied Bioscience*.2017; 5(5):88-96.
34. Jangir C K, Singh D and Kumar S. Yield and economic response of biofertilizer and fertility levels on black gram (*Vigna mungo* L.). *Progressive Research- An International Journal*.2017; 11(8): 5252-5254.
35. Badiyala D and Chopra P. Effect of zinc and FYM on productivity and nutrient availability in maize (*Zea mays* L.) - linseed (*Linum usitatissimum*) cropping sequence. *Indian Journal of Agronomy*.2011; 56: 88-91.
36. Kannan P, Arunacham P, Prabu K G and Prabhakaran J. Response of black gram (*Vigna mungo* L.) to multi micronutrient mixtures under rainfed Alfisol. *Journal of the Indian Society of Soil Science*.2017; 62(2): 154-160.

Table 2.0: Effect of Fe, Zn and Bio-fertilizers application on nutrient content in seed and straw of urdbean.

Treatments	Nitrogen content (%)		Phosphorous content (%)		Potassium content (%)		Iron content (mg kg ⁻¹)		Zinc content (mg kg ⁻¹)	
	Seed	Straw	Seed	Straw	Seed	Straw	Seed	Straw	Seed	Straw
T ₁ : Control	1.66	1.37	0.22	0.117	0.40	1.33	27.22	42.94	20.05	11.87
T ₂ : 75 % RDF	1.94	1.48	0.24	0.123	0.44	1.48	31.13	39.87	21.79	13.58
T ₃ : 100 % RDF	2.45	1.57	0.25	0.129	0.47	1.57	34.05	53.46	23.84	14.85
T ₄ : T ₂ + <i>Rhizobium</i> @ 600 g ha ⁻¹ seed inoculation	2.49	1.59	0.27	0.133	0.48	1.60	36.14	56.73	25.29	15.76
T ₅ : T ₂ + PSB @ 600 g ha ⁻¹ seed inoculation	2.46	1.58	0.26	0.131	0.47	1.57	35.57	55.84	24.90	15.51
T ₆ : T ₂ + <i>Rhizobium</i> @ 600 g ha ⁻¹ + PSB @ 600 g ha ⁻¹	2.53	1.70	0.28	0.141	0.51	1.67	37.58	58.99	26.30	16.39
T ₇ : T ₂ + 0.1 % FeSO ₄ + 0.5 % ZnSO ₄ at pre flowering and pod formation stage	2.52	1.63	0.29	0.134	0.50	1.62	36.72	57.65	25.70	16.01
T ₈ : T ₂ + T ₇ + <i>Rhizobium</i> @ 600 g ha ⁻¹ seed inoculation	3.11	1.77	0.31	0.149	0.53	1.80	41.82	65.66	30.27	18.24
T ₉ : T ₂ + T ₇ + PSB @ 600 g ha ⁻¹ seed inoculation	2.99	1.75	0.30	0.145	0.53	1.72	41.38	64.96	29.96	18.05
T ₁₀ : T ₂ + T ₇ + <i>Rhizobium</i> 600 g ha ⁻¹ + PSB @ 600 g ha ⁻¹ seed inoculation	3.87	1.84	0.34	0.154	0.56	1.82	45.15	73.89	32.61	19.69
SEm±	0.12	0.03	0.01	0.003	0.01	0.05	1.95	4.87	0.99	0.61
CD (p=0.05)	0.37	0.10	0.03	0.008	0.03	0.13	5.80	14.47	2.93	1.83

Table 3.0: Effect of Fe, Zn and Bio-fertilizers application on nutrient uptake by seed, straw and total by urdbean crop.

Treatments	Nitrogen uptake (kg ha ⁻¹)			Phosphorus uptake (kg ha ⁻¹)			Potassium uptake (kg ha ⁻¹)			Iron uptake (g ha ⁻¹)			Zinc uptake (g ha ⁻¹)		
	Seed	Straw	Total	Seed	Straw	Total	Seed	Straw	Total	Seed	Straw	Total	Seed	Straw	Total
T ₁ : Control	10.04	10.99	21.03	1.33	0.94	2.27	2.41	10.68	13.09	167.06	344.14	511.20	120.59	95.23	215.82
T ₂ : 75 % RDF	14.26	13.76	28.02	1.76	1.14	2.90	3.25	13.76	17.00	227.51	461.25	688.76	159.97	125.72	285.69
T ₃ : 100 % RDF	20.00	16.13	36.13	2.04	1.32	3.37	3.82	16.01	19.83	278.19	550.66	828.85	194.90	152.17	347.07
T ₄ : T ₂ + <i>Rhizobium</i> @ 600 g ha ⁻¹ seed inoculation	21.39	17.43	38.83	2.33	1.46	3.79	4.15	17.47	21.63	310.48	625.75	936.23	218.81	171.05	389.87
T ₅ : T ₂ + PSB @ 600 g ha ⁻¹ seed inoculation	21.05	17.22	38.27	2.22	1.43	3.64	4.02	17.11	21.13	303.01	607.60	910.61	212.47	168.28	380.74
T ₆ : T ₂ + <i>Rhizobium</i> @ 600 g ha ⁻¹ + PSB @ 600 g ha ⁻¹	22.93	19.57	42.51	2.53	1.62	4.15	4.62	19.21	23.83	336.86	685.66	1022.51	236.50	187.88	424.38
T ₇ : T ₂ + 0.1 % FeSO ₄ + 0.5 % ZnSO ₄ at pre flowering and pod formation stage	22.20	18.20	40.40	2.56	1.50	4.05	4.35	18.03	22.38	325.38	643.15	968.53	226.17	178.83	404.99
T ₈ : T ₂ + T ₇ + <i>Rhizobium</i> @ 600 g ha ⁻¹ seed inoculation	31.26	21.99	53.25	3.11	1.85	4.96	5.32	22.44	27.76	419.13	809.61	1228.75	304.10	226.82	530.92
T ₉ : T ₂ + T ₇ + PSB @ 600 g ha ⁻¹ seed inoculation	29.72	21.63	51.35	2.98	1.79	4.77	5.23	21.22	26.46	410.23	804.38	1214.62	295.57	223.41	518.98
T ₁₀ : T ₂ + T ₇ + <i>Rhizobium</i> 600 g ha ⁻¹ + PSB @ 600 g ha ⁻¹ seed inoculation	49.36	27.84	77.20	4.32	2.33	6.65	7.16	27.44	34.60	573.25	1120.36	1693.60	413.78	297.40	711.18
SEm±	2.01	1.12	2.32	0.13	0.09	0.15	0.25	1.09	1.12	21.34	72.82	64.02	11.66	9.11	11.87
CD (p=0.05)	5.98	3.33	6.91	0.39	0.28	0.46	0.74	3.25	3.32	63.43	216.35	190.21	34.65	27.08	35.28

Table 4.0: Effect of Fe, Zn and Bio-fertilizers application on yield, quality and monetary return of urdbean.

Treatments	pH	EC (dS m ⁻¹)	Organic carbon (%)	Bulk density (Mg m ⁻³)	Particle density (Mg m ⁻³)	Porosity (%)	Available macro nutrients (kg ha ⁻¹)			Available micro nutrients (mg kg ⁻¹)	
							N	P	K	Fe	Zn
T ₁ : Control	7.32	0.49	0.41	1.33	2.45	45.71	228.21	17.46	230.10	4.30	0.59
T ₂ : 75 % RDF	7.28	0.61	0.42	1.32	2.43	45.68	238.88	20.72	241.87	4.64	0.62
T ₃ : 100 % RDF	7.27	0.68	0.43	1.33	2.45	45.71	245.84	21.28	243.57	4.70	0.65
T ₄ : T ₂ + <i>Rhizobium</i> @ 600 g ha ⁻¹ seed inoculation	7.25	0.62	0.42	1.31	2.41	45.64	266.92	21.79	245.39	5.15	0.67
T ₅ : T ₂ + PSB @ 600 g ha ⁻¹ seed inoculation	7.22	0.61	0.44	1.31	2.41	45.64	260.33	23.77	247.51	5.07	0.66
T ₆ : T ₂ + <i>Rhizobium</i> @ 600 g ha ⁻¹ + PSB @ 600 g ha ⁻¹	7.20	0.63	0.46	1.31	2.41	45.64	272.88	24.00	250.82	5.18	0.69
T ₇ : T ₂ + 0.1 % FeSO ₄ + 0.5 % ZnSO ₄ at pre flowering and pod formation stage	7.21	0.64	0.44	1.32	2.43	45.68	258.54	22.74	249.71	5.25	0.71
T ₈ : T ₂ + T ₇ + <i>Rhizobium</i> @ 600 g ha ⁻¹ seed inoculation	7.20	0.63	0.46	1.32	2.43	45.68	280.17	23.61	254.75	5.48	0.74
T ₉ : T ₂ + T ₇ + PSB @ 600 g ha ⁻¹ seed inoculation	7.23	0.65	0.45	1.32	2.43	45.68	275.88	24.47	250.15	5.44	0.72
T ₁₀ : T ₂ + T ₇ + <i>Rhizobium</i> 600 g ha ⁻¹ + PSB @ 600 g ha ⁻¹ seed inoculation	7.19	0.64	0.48	1.32	2.43	45.68	285.21	25.41	252.66	5.75	0.76
SEm±	0.07	0.10	0.02	0.01	0.02	0.13	8.02	1.06	13.30	0.10	0.02
CD (p=0.05)	NS	NS	NS	NS	NS	NS	23.84	3.15	NS	0.30	0.06