

Short communication

EFMR Syndrome: Epilepsy and mental retardation restricted to females in Childhood

Abstract

Epilepsy with intelligence impairment, restricted to the female sex, is a rare X-linked epilepsy syndrome. It is characterized by febrile or afebrile seizures, mainly tonic-clonic, but also absence, myoclonic, and atonic beginning in the first years of life. In most cases, developmental delay and intelligence impairment of varying severity is present. Behavioral disorders, autistic traits, hyperactivity, and aggressiveness are commonly associated. This disease exclusively affects females. Male carriers are not affected despite an X-linked inheritance.

Key Words

EFMR-PCDH19-children

Introduction

An epilepsy with intelligence impairment, focused to the female sex, is a rare X-chromosomal epilepsy syndrome described as EFMR syndrome. It is based on febrile or afebrile seizures, tonic-clonic aspect, but also absental, myoclonic aspect, and atonic features are described. Seizures start in the first years of life and developmental problems and low intelligence level of varying severity is present. Behavioral disturbances with autistic features, hyperactivity signs and aggressiveness in behaviour are also commonly found. This disease exclusively is present in females. Male carriers are not ill despite an X-chromosomal inheritance. X-chromosomal epilepsy with mental retardation, also known as epilepsy and mental retardation limited to females (EFMR), was described in 1971 based on a gross family of 15 female children. In 2008, the gene focus was localized to chromosome Xq22 in further familial analysis. First seizures occur normally before 14 months of age and were often associated with fever signs. Seizure includes tonic-clonic, tonic, partial, atonic, myoclonic features, and absences. Developmental delay and intelligence development can be highly varying. The cause of X-linked epilepsy with intellectual disability is found in pathological variants in the gene for protocadherin 9 (PCDH19). Protocadherin 9 shows its expression during brain development and represents one of the first members of cadherin family to be changed in epilepsy and mental impairment. More than 200 different pathogenic PCDH19 variants have now been described. Genomic deletions are found in the Xq22.1 region with the entire PCDH19 gene or multiple affected exon regions have been found in 3% of female patients. Heterozygous carriers of pathogenic variants of the disease are affected, whereas hemizygous male carriers are frequently asymptomatic. It is an untypical mode of inheritance of an X-chromosomal disease termed cellular interference. While clinically inconspicuous male carriers only have cells with pathogenic PCDH19 variants, the presence of cells with and without pathogenic PCDH19 variants in the female organism creates mosaics through random X-inactivation, which only then become pathogenic. In the literature, isolated symptomatic male patients with a pathogenic PCDH19 variant have been described in which these were also present as mosaics. This confirms the mechanism according to which cells with and without a pathogenic variant must be present for disease development. Mutations in PCDH19 can cause developmental delays of varying severity (1-12). Developmental regression is also observed in about 50% of patients, although normal development has been described in some patients (1-30). In addition, seizures are typical of mutations in the above gene and may be convulsive, tonic-clonic, focal, atonic or myoclonic (1-12). In addition, affected women show various psychiatric manifestations such as symptoms from the autism spectrum, aggression and compulsive behaviour. The penetrance of the pathogenic PCDH19 variants is estimated to be about 90% (3,4,8,10). The non-penetrance of the remaining 10% has not yet been definitively clarified.

EFMR Syndrome

Like Dravet syndrome, it is classified as an early infantile epileptic encephalopathy (EIEE9) [MIM 30088]. Typically, an X-linked inheritance pattern is expressed by affected males and unaffected females who are heterozygous for this trait (2,3,8,9,19). In the case of the EFMR inheritance pattern, it is the other way around: the pedigree of affected patients falls through exclusively affected females into unaffected males. EFMR was first described in 1971 by Juberg and Hellman in a North American family (1-12). In this family, female members were found to have seizure onset in infancy followed by developmental regression with mild to severe intellectual impairment. The average age at seizure onset was approximately 14 months (1-30). The seizures are described as afebrile focal and generalized tonic-clonic seizures, with a gradual increase in seizure frequency (3,4,8,29,30). Scheffer et al. studied the clinical picture of 27 girls and women from four different families suffering from EFMR. They found tonic-clonic focal and generalized seizures, absences, myoclonic, tonic, and atonic seizures. Sixty-three percent of these seizures occurred with fever. At an average age of 12 years, the seizures ceased. In addition, Scheffer found that developmental regressions varied widely.

Furthermore, both status epilepticus and the occurrence of myoclonia are much less frequent in EFMR. While 92% of patients with Dravet syndrome have experienced status epilepticus, status epilepticus occurs in approximately 33% of EFMR patients. In contrast, a cluster of seizures within a short period of time is more common in EFMR, whereas this phenomenon is much less common in Dravet syndrome. Overall, the prognosis of PCDH19-positive patients appears to be more favorable than that of SCN1A-positive patients; PCDH19 patients usually experience seizure remission at puberty, whereas SCN1A patients have a high mortality rate and epileptic seizures in adulthood. EFMR patients are characterized by greater variability in their cognitive abilities, with up to one-third of girls having normal intellect and the remainder mildly to moderately impaired, in contrast to patients with Dravet syndrome, who are usually more difficult cognitively. Severe cases of intellectual impairment, however, can occur in both. Autistic features are again more common in EFMR (1-13). In addition to autistic features, compulsive and aggressive behaviors may occur. In some patients, social withdrawal plays the greatest role among the disease features, especially as patients grow older.

References:

- (1) Nagarajan L, Ghosh S, Dyke J, Lee S, Silberstein J, Azmanov D, Richard W. Epilepsy surgery in PCDH 19 related developmental and epileptic encephalopathy: A case report. *Epilepsy Behav Rep.* 2022 Jul 6;19:100560. doi: 10.1016/j.ebr.2022.100560. PMID: 35856042; PMCID: PMC9287778.
- (2) Yang C, Shi Y, Li X, Guan L, Li H, Lin J. Cadherins and the pathogenesis of epilepsy. *Cell Biochem Funct.* 2022 Jun;40(4):336-348. doi: 10.1002/cbf.3699. Epub 2022 Apr 8. PMID: 35393670.
- (3) Breuillard D, Leunen D, Chemaly N, Auclair L, Pinard JM, Kaminska A, Desguerre I, Ouss L, Nabbout R. Autism spectrum disorder phenotype and intellectual disability in females with epilepsy and PCDH-19 mutations. *Epilepsy Behav.* 2016 Jul;60:75-80. doi: 10.1016/j.yebeh.2016.04.009. Epub 2016 May 12. PMID: 27179713.
- (4) Biswas S, Emond MR, Jontes JD. Protocadherin-19 and N-cadherin interact to control cell movements during anterior neurulation. *J Cell Biol.* 2010 Nov 29;191(5):1029-41. doi: 10.1083/jcb.201007008. PMID: 21115806; PMCID: PMC2995167.
- (5) Yokota S, Hirayama T, Hirano K, Kaneko R, Toyoda S, Kawamura Y, Hirabayashi M, Hirabayashi T, Yagi T. Identification of the cluster control region for the protocadherin-beta genes located beyond the protocadherin-gamma cluster. *J Biol Chem.* 2011 Sep 9;286(36):31885-95. doi: 10.1074/jbc.M111.245605. Epub 2011 Jul 19. PMID: 21771796; PMCID: PMC3173131.
- (6) Liu YH, Cheng YT, Tsai MH, Chou IJ, Hung PC, Hsieh MY, Wang YS, Chen YJ, Kuo CY, Lin JJ, Wang HS, Lin KL. Genetics and clinical correlation of Dravet syndrome and its mimics - experience of a tertiary center in Taiwan. *Pediatr Neonatol.* 2021 Sep;62(5):550-558. doi: 10.1016/j.pedneo.2021.05.022. Epub 2021 Jun 23. PMID: 34226156.

- (7) Iqbal M, Maroofian R, Çavdarlı B, Riccardi F, Field M, Banka S, Bubshait DK, Li Y, Hertecant J, Baig SM, Dymont D, Efthymiou S, Abdullah U, Makhdoom EUH, Ali Z, Scherf de Almeida T, Molinari F, Mignon-Ravix C, Chabrol B, Antony J, Ades L, Pagnamenta AT, Jackson A, Douzgou S; Genomics England Research Consortium; Beetz C, Karageorgou V, Vona B, Rad A, Baig JM, Sultan T, Alvi JR, Maqbool S, Rahman F, Toosi MB, Ashrafzadeh F, Imannezhad S, Karimiani EG, Sarwar Y, Khan S, Jameel M, Noegel AA, Budde B, Altmüller J, Motameny S, Höhne W, Houlden H, Nürnberg P, Wollnik B, Villard L, Alkuraya FS, Osmond M, Hussain MS, Yigit G. Biallelic variants in PCDHGC4 cause a novel neurodevelopmental syndrome with progressive microcephaly, seizures, and joint anomalies. *Genet Med*. 2021 Nov;23(11):2138-2149. doi: 10.1038/s41436-021-01260-4. Epub 2021 Jul 9. PMID: 34244665; PMCID: PMC8553613.
- (8) Lim J, Ryu J, Kang S, Noh HJ, Kim CH. Autism-like behaviors in male mice with a Pcdh19 deletion. *Mol Brain*. 2019 Nov 20;12(1):95. doi: 10.1186/s13041-019-0519-3. PMID: 31747920; PMCID: PMC6864969.
- (9) Antelmi E, Mastrangelo M, Bisulli F, Spaccini L, Stipa C, Mostacci B, Mei D, Guerrini R, Tinuper P. Semiological study of ictal affective behaviour in epilepsy and mental retardation limited to females (EFMR). *Epileptic Disord*. 2012 Sep;14(3):304-9. doi: 10.1684/epd.2012.0526. PMID: 22932693.
- (10) Dell'Isola GB, Vinti V, Fattorusso A, Tascini G, Mencaroni E, Di Cara G, Striano P, Verrotti A. The Broad Clinical Spectrum of Epilepsies Associated With Protocadherin 19 Gene Mutation. *Front Neurol*. 2022 Jan 17;12:780053. doi: 10.3389/fneur.2021.780053. PMID: 35111125; PMCID: PMC8801579.
- (11) Yang L, Arafat A, Peng J, Chen C, Ma Y, Yin F. [PCDH19 gene mutations lead to epilepsy with mental retardation limited to females in 2 cases and literature review]. *Zhong Nan Da Xue Xue Bao Yi Xue Ban*. 2017 Jun 28;42(6):730-736. Chinese. doi: 10.11817/j.issn.1672-7347.2017.06.021. PMID: 28690234.
- (12) Duszyc K, Terczynska I, Hoffman-Zacharska D. Epilepsy and mental retardation restricted to females: X-linked epileptic infantile encephalopathy of unusual inheritance. *J Appl Genet*. 2015 Feb;56(1):49-56. doi: 10.1007/s13353-014-0243-8. Epub 2014 Sep 10. PMID: 25204757.
- (13) Scheffer IE, Turner SJ, Dibbens LM, Bayly MA, Friend K, Hodgson B, Burrows L, Shaw M, Wei C, Ullmann R, Ropers HH, Szepietowski P, Haan E, Mazarib A, Afawi Z, Neufeld MY, Andrews PI, Wallace G, Kivity S, Lev D, Lerman-Sagie T, Derry CP, Korczyn AD, Gecz J, Mulley JC, Berkovic SF. Epilepsy and mental retardation limited to females: an under-recognized disorder. *Brain*. 2008 Apr;131(Pt 4):918-27. doi: 10.1093/brain/awm338. Epub 2008 Jan 29. PMID: 18234694.
- (14) Kolc KL, Sadleir LG, Scheffer IE, Ivancevic A, Roberts R, Pham DH, Gecz J. A systematic review and meta-analysis of 271 PCDH19-variant individuals identifies psychiatric comorbidities, and association of seizure onset and disease severity. *Mol Psychiatry*. 2019 Feb;24(2):241-251. doi: 10.1038/s41380-018-0066-9. Epub 2018 Jun 11. PMID: 29892053; PMCID: PMC6344372.
- (15) Zhao X, Wang Y, Mei S, Kong X. A novel PCDH19 missense mutation, c.812G>A (p.Gly271Asp), identified using whole-exome sequencing in a Chinese family with epilepsy female restricted mental retardation syndrome. *Mol Genet Genomic Med*. 2020 Jun;8(6):e1234. doi: 10.1002/mgg3.1234. Epub 2020 Apr 21. PMID: 32314541; PMCID: PMC7284031.
- (16) Hudson JD, Tamilselvan E, Sotomayor M, Cooper SR. A complete Protocadherin-19 ectodomain model for evaluating epilepsy-causing mutations and potential protein interaction sites. *Structure*. 2021 Oct 7;29(10):1128-1143.e4. doi: 10.1016/j.str.2021.07.006. Epub 2021 Sep 13. PMID: 34520737.
- (17) Chemaly N, Losito E, Pinard JM, Gautier A, Villeneuve N, Arbues AS, An I, Desguerre I, Dulac O, Chiron C, Kaminska A, Nabbout R. Early and long-term electroclinical features of

patients with epilepsy and PCDH19 mutation. *Epileptic Disord.* 2018 Dec 1;20(6):457-467. doi: 10.1684/epd.2018.1009. PMID: 30530412.

(18) Kozina AA, Okuneva EG, Baryshnikova NV, Fedonyuk ID, Kholin AA, Il'ina ES, Krasnenko AY, Stetsenko IF, Plotnikov NA, Klimchuk OI, Surkova EI, Ilinsky VV. Two novel PCDH19 mutations in Russian patients with epilepsy with intellectual disability limited to females: a case report. *BMC Med Genet.* 2020 Oct 21;21(1):209. doi: 10.1186/s12881-020-01119-6. PMID: 33087045; PMCID: PMC7579871.

(19) Brown A, Arpone M, Schneider AL, Micallef S, Anderson VA, Scheffer IE. Cognitive, behavioral, and social functioning in children and adults with Dravet syndrome. *Epilepsy Behav.* 2020 Nov;112:107319. doi: 10.1016/j.yebeh.2020.107319. Epub 2020 Aug 25. PMID: 32858363.

(20) Chen ZS, Hsieh A, Sun G, Bergey GK, Berkovic SF, Perucca P, D'Souza W, Elder CJ, Farooque P, Johnson EL, Barnard S, Nightscales R, Kwan P, Moseley B, O'Brien TJ, Sivathamboo S, Laze J, Friedman D, Devinsky O; MS-BioS Study Group. Interictal EEG and ECG for SUDEP Risk Assessment: A Retrospective Multicenter Cohort Study. *Front Neurol.* 2022 Mar 18;13:858333. doi: 10.3389/fneur.2022.858333. PMID: 35370908; PMCID: PMC8973318.

(21) Krüger J, Schubert J, Kegele J, Labalme A, Mao M, Heighway J, Seebohm G, Yan P, Koko M, Aslan-Kara K, Caglayan H, Steinhoff BJ, Weber YG, Keo-Kosal P, Berkovic SF, Hildebrand MS, Petrou S, Krause R, May P, Lesca G, Maljevic S, Lerche H. Loss-of-function variants in the *KCNQ5* gene are implicated in genetic generalized epilepsies. *EBioMedicine.* 2022 Oct;84:104244. doi: 10.1016/j.ebiom.2022.104244. Epub 2022 Sep 9. PMID: 36088682; PMCID: PMC9471468.

(22) Chouery E, Makhlof J, Daoud Khatoun W, Mehawej C, Megarbane A. *PCDH19* in Males: Are Hemizygous Variants Linked to Autism? *Genes (Basel).* 2023 Feb 27;14(3):598. doi: 10.3390/genes14030598. PMID: 36980870; PMCID: PMC10048232.

(23) Chen Y, Yang X, Chen J, Yang X, Yang Y, Liu A, Zhang X, Wu W, Sun D, Yang Z, Jiang Y, Zhang Y. *PCDH19*-related epilepsy in mosaic males: The phenotypic implication of genotype and variant allele frequency. *Front Neurol.* 2022 Nov 3;13:1041509. doi: 10.3389/fneur.2022.1041509. PMID: 36408521; PMCID: PMC9669318.

(24) Pancho A, Mitsogiannis MD, Aerts T, Dalla Vecchia M, Ebert LK, Geenen L, Noterdaeme L, Vanlaer R, Stulens A, Hulpiau P, Staes K, Van Roy F, Dedecker P, Schermer B, Seuntjens E. Modifying *PCDH19* levels affects cortical interneuron migration. *Front Neurosci.* 2022 Oct 25;16:887478. doi: 10.3389/fnins.2022.887478. PMID: 36389226; PMCID: PMC9642031.

(25) Gecez J, Thomas PQ. Disentangling the paradox of the *PCDH19* clustering epilepsy, a disorder of cellular mosaics. *Curr Opin Genet Dev.* 2020 Dec;65:169-175. doi: 10.1016/j.gde.2020.06.012. Epub 2020 Jul 26. PMID: 32726744.

(26) Hoshina N, Johnson-Venkatesh EM, Hoshina M, Umemori H. Female-specific synaptic dysfunction and cognitive impairment in a mouse model of *PCDH19* disorder. *Science.* 2021 Apr 16;372(6539):eaaz3893. doi: 10.1126/science.aaz3893. PMID: 33859005; PMCID: PMC9873198.

(27) Moncayo JA, Vargas MN, Castillo I, Granda PV, Duque AM, Argudo JM, Matcheswalla S, Lopez Dominguez GE, Monteros G, Andrade AF, Ojeda D, Yopez M. Adjuvant Treatment for Protocadherin 19 (*PCDH19*) Syndrome. *Cureus.* 2022 Jul 22;14(7):e27154. doi: 10.7759/cureus.27154. PMID: 36004035; PMCID: PMC9392850.

(28) Smith L, Singhal N, El Achkar CM, Truglio G, Rosen Sheidley B, Sullivan J, Poduri A. *PCDH19*-related epilepsy is associated with a broad neurodevelopmental spectrum. *Epilepsia.* 2018 Mar;59(3):679-689. doi: 10.1111/epi.14003. Epub 2018 Jan 28. Erratum in: *Epilepsia.* 2018 Jun;59(6):1272. PMID: 29377098; PMCID: PMC6264912.

(29) Sadleir LG, Kole KL, King C, Mefford HC, Dale RC, Gecz J, Scheffer IE. Levetiracetam efficacy in PCDH19 Girls Clustering Epilepsy. *Eur J Paediatr Neurol*. 2020 Jan;24:142-147. doi: 10.1016/j.ejpn.2019.12.020. Epub 2020 Jan 3. PMID: 31928905.

(30) Gerosa L, Mazzoleni S, Rusconi F, Longaretti A, Lewerissa E, Pelucchi S, Murru L, Giannelli SG, Broccoli V, Marcello E, Kasri NN, Battaglioli E, Passafaro M, Bassani S. The epilepsy-associated protein PCDH19 undergoes NMDA receptor-dependent proteolytic cleavage and regulates the expression of immediate-early genes. *Cell Rep*. 2022 May 24;39(8):110857. doi: 10.1016/j.celrep.2022.110857. PMID: 35613587; PMCID: PMC9152703.

UNDER PEER REVIEW