

Case study

Primary Splenic Diffuse Large B-Cell Lymphoma with Massive Splenomegaly

Abstract

Primary malignancies of the spleen are very rare, diffuse large B-cell lymphoma of the spleen is a rare primary neoplasm. Primary splenic diffuse large B-cell lymphoma is a type of non-Hodgkin's lymphoma that constitute, less than 1% of non-Hodgkin's lymphoma and less than 2% of all lymphomas. Primary splenic lymphoma has been associated with the hepatitis C and HIV virus. It affects men more than women and present in late middle age.

Patient with diffuse large B cell lymphoma typically present with fever, fatigues weight loss, palpable splenomegaly and left upper quadrant pain. Diffuse large B cell lymphoma stage I being limited to the spleen, whereas stage II spleen and hilar Lymph-node involvement and stage III is distant metastasis including liver and surrounding organs involvements, stomach, pancreas, kidney and colon. Stage I disease, a splenectomy is the standard treatment for DLBCL. Radiation and chemotherapy can be used. When the cancer is confined to the spleen, the prognosis is quite favourable with a median survival of 7.48 years.

We are reporting an extremely rare case of primary splenic lymphoma in a 45 years old female patient, presented with abdominal pain and huge splenomegaly. The CT revealed a bulky, low density, irregular mass in the spleen and clinically disease limited to spleen only. So we performed open total splenectomy.

Keywords

Diffuse large B. cell lymphoma, Primary splenic lymphoma, Splenomegaly

Introduction

Lymphoma is a type of blood cancer that develops from white blood cells called lymphocytes grow out of control. There are two types of lymphocytes. B-lymphocytes (B-cell) and T-lymphocytes (T-cell). Lymphoma can be grouped as Hodgkin's lymphoma or Non-Hodgkin's lymphoma. Primary splenic diffuse large B-cell lymphoma DLBCL is a rare type of Non-Hodgkin's lymphoma. It commonly present with abdominal pain, splenic mass and an elevated level of lactate dehydrogenase level. Primary splenic lymphoma are again subtypes. (2,3,4)

1. Marginal zone lymphoma
2. Diffuse large B-cell lymphoma
3. T-cell lymphoma

Diagnosis is bases on imaging findings of splenomegaly with hypoechoic masses on ultrasonography and CT scan. Treatment option includes total splenectomy, chemotherapy and splenic radiation. (2,3,4)

Case Report

A 45 years old female was admitted to our centre on 10/02/2011, with complaints of weight loss, abdominal pain and distention of abdomen. Physical examination revealed, she had a protuberant abdomen with firm, huge palpable splenomegaly extending from xiphisternum to below the umbilicus. There was no ascites or hepatomegaly or lymphadenopathy. Abdominal ultrasonography

and CT showed massive splenomegaly size 27x15x10 cm with a huge heterogeneous hypoechoic masses in the spleen. Haematological examination revealed pancytopenia. WBC 1.9×10^9 /L (neutrophils 48.0%, eosinophils 0%, basophils 0%, monocytes 4.0%, lymphocytes 48.0%), Hb 10.3 g/dL, platelet count 99×10^9 /L, LDH 127 U/L (normal range 100-220 U/L). Liver and renal functions were within normal limits. Bone marrow test was normal. A final diagnosis of a primary splenic malignant tumour was reached and patient was prepared for surgery.

The patient underwent total splenectomy, through a mid-line surgical approach. During the operation, Patient had huge splenomegaly with multiple nodular lesions on the surface of the spleen. The liver, mesenteric or retro peritoneal lymph nodes were not involved. The spleen was adherent to stomach, omentum and sub-diaphragmatic area and was separated carefully. Total splenectomy done, achieving good haemostasis, drainage tube kept and abdomen closed in layers.

Gross specimen of spleen weighing 4 kg. Massive splenomegaly size 27x15x10 cm. The Cut surface was almost totally effaced by a huge grey-white homogenous tumour. Microscopically revealed diffuse proliferation of large neoplastic B-cell lymphoma –stage I. Splenic lymphoma was diagnosed and there were no hilar lymph-nodes. Eight days after the operation, the patient revealed from the pancytopenia. WBC of 7.6×10^9 /L, Hb 13.0 g/dL and platelet count 330×10^9 /L although CRP increased to 8.21 mg/dL. Patient recovery was uneventful and patient discharged on 10th post-operative days. The patient was advised chemotherapy with CHOP regimen and after one year follow up, patient is found to be symptom free and healthy. (Fig 1-8)




Fig-1 CT Abdomen showing 27x15x10 cm splenomegaly with hypoechoic lesion in spleen




Fig-2 Intra-operative photograph showing huge splenic mass at of lower pole of spleen




Fig-3 Intra-operative photograph showing huge splenic mass with splenomegaly




Fig-4 Intra-operative photograph showing huge splenic mass with splenic pedicle




Fig-5 Intra-operative photograph showing huge splenic mass of size 27x15x10

Fig-6 showing huge Splenomegaly with huge splenic mass of- weighing 4 kg





Fig-7 Cut -section showing gray – white fleshy tumor

Fig-8 Microscopic -diffuse Large B-cell lymphoma

Primary splenic diffuse large B Cell lymphoma is very rare as it occurs in less than 1% of non-Hodgkin's lymphoma. It is most commonly found in females and older males. Symptoms can include splenomegaly, left upper quadrant pain, fever and weight loss. It can also be associated with HIV virus and can present with metastasis to hilar and retroperitoneal lymph-nodes. It also causes cystopenia. (2,5)

Ahmann's classification.

Primary splenic diffuse large B-cell lymphoma is classified in three stage.

1. Stage I – Disease confined to the spleen.
2. Stage II – Involvement of the spleen and hilar lymph-nodes.
3. Stage III - Extra splenic nodule or hepatic mesenteric or retroperitoneal lymph-nodes. (2,3,4)

Imaging studies, ultrasonography and CT abdomen shows hypodense or hypoechoic lesion in the spleen. Primary splenic lymphomas histological subtypes including.

1. Diffuse large B-cell lymphoma
2. Marginal zone lymphoma
3. Anaplastic large cell lymphoma (ALCL)
4. Follicular lymphoma. (2,3)

The most common variant is diffuse large B-cell lymphoma accounting for approximately 31% of all non-Hodgkin's lymphoma cases, anaplastic variant of diffuse large B-cell lymphoma was first defined in the 2017. A rare morphological variant of DLBCL and constitutes less than 3.4% of all DLBCL cases. AV –DLBCL affect nodal and extra nodal sites. Patient present with fever, weight loss and night sweats. (2,4)

Splenectomy is considered the first line of procedure and has shown high efficacy in both diagnosis and treatment of splenic lesions. Several studies have demonstrated the role of splenectomy in improving the prognosis of primary splenic lymphoma. Adjutant chemotherapy including R-CHOP regimen i.e. Rituximab, cyclophosphamide, doxorubicin, vincristine and prednisolone.

Bairey et al studies 87 patient with primary splenic DLBCL, Splenectomy and chemotherapy had a better free survival. Brox et al studies 9 patients with primary splenic lymphoma and median survival time of 7.48 years who underwent splenectomy with chemotherapy. Splenic irradiation is also treatment option for inoperable cases, in order to reduce the size of the spleen. (2,5)

Conclusion

In cases of massive splenomegaly, primary splenic diffuse large B-cell lymphoma a total splenectomy is the gold standard procedure. Radiation and chemotherapy can be used. When the cancer is confined to the spleen, the prognosis is quite favourable with a median survival of 7.48 years.

References

1. Jillian L. VanVliet, Primary Diffuse Large B Cell Lymphoma of the Spleen, Journal of Diagnostic Medical Sonography 26(3) 147–149 © The Author(s) 2010 Reprints and permission: <http://www.sagepub.com/journalsPermissions.nav> DOI: 10.1177/8756479310368431
2. Khalid S, Daw JH, Daw H, Haddad A. Primary Splenic Diffuse Large B-Cell Lymphoma: A Rare Case of Massive Splenomegaly and Thrombocytopenia. Cureus. 2018 Jul 22;10(7):e3026. doi: 10.7759/cureus.3026. PMID: 30254815; PMCID: PMC6150744.
3. Ismail S, Ali F, Ajamieh H, Kanaan S, Issa R, Daoud A, Alshehabi Z. Primary splenic anaplastic variant of diffuse large B-cell lymphoma: a case report. J Med Case Rep. 2021 May 6;15(1):227. doi: 10.1186/s13256-021-02846-x. PMID: 33952307; PMCID: PMC8101127.
4. Bairey, O., Shvidel, L., Perry, C., Dann, E.J., Ruchlemer, R., Tadmor, T. and Goldschmidt, N. (2015), Characteristics of primary splenic diffuse large B-cell lymphoma and role of splenectomy in improving survival. Cancer, 121: 2909-2916. <https://doi.org/10.1002/cncr.29487>

5. Taha A. Baiomy, Ola A. Harb, Ahmed A. Obaya, Loay M. Gertallah, Primary splenic non-Hodgkin lymphoma of diffuse large B cell type; a case report and review of the literature, *Human Pathology: Case Reports*, <https://doi.org/10.1016/j.ehpc.2020.200459>.
6. Rajendran T, Kini JR, Abna A, et al Clinical benefit of R-CHOP without splenectomy in stage I primary splenic diffuse large B-cell lymphoma *BMJ Case Reports CP* 2022;15:e246610.
7. Hui Li and others, Unusual Presentation of Primary Splenic Diffuse Large B Cell Lymphoma of Red Pulp, *American Journal of Clinical Pathology*, Volume 144, Issue suppl_2, 1 October 2015, <https://doi.org/10.1093/ajcp/144.suppl2.117>
8. Chen, Yue-Ren, Hwa-Koon Wu, and Yang-Yuan Chen. 2023. "Unusual Rapid Growth of Primary Splenic Diffuse Large B-Cell Lymphoma with Extensive Necrosis" *Diagnostics* 13, no. 1: 35. <https://doi.org/10.3390/diagnostics13010035>
9. Maia Neves N, Lopes AP, Carvalho Coelho S, Raimundo A, Mafra MM, Bayão Horta A. Primary splenic diffuse large B-cell lymphoma: a case report. *EJCRIM* 2023;10 doi:10.12890/2023_003932.
10. Maia Neves N, Lopes AP, Carvalho Coelho S, Raimundo A, Mafra MM, Bayão Horta A. Primary splenic diffuse large B-cell lymphoma: a case report. *EJCRIM* 2023;10 doi:10.12890/2023_003932.